



Net Weight: 25 lbs.

# Dusting Sulphur

**ACTIVE INGREDIENT:**  
Sulphur 93.0

**INERT INGREDIENTS:** 7.0

Total 100.0%

(Sulphur fineness 93% through 325 mesh screen)  
EPA Reg. No. 769-478 EPA Est. 769-AL-1

## CAUTION

KEEP OUT OF REACH OF CHILDREN.

### PRECAUTIONARY STATEMENTS

**HAZARDS TO HUMANS:** Harmful if swallowed. Avoid breathing dust. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not use for medical purposes or as a feed supplement.

**FIRST AID IF IN EYES:** Immediately flush with plenty of water for at least 15 minutes. Call a physician if irritation persists. **FOR SKIN CONTACT:** Wash with soap and water. **IF INHALED:** and overcome by vapor or dust, remove immediately from exposure and call a physician. Administer artificial respiration if breathing is irregular or has stopped. **IF SWALLOWED:** Drink 1 to 2 glasses of water and induce vomiting by sticking finger in back of throat. Call a physician.

**ENVIRONMENTAL HAZARDS:** Do not contaminate water by cleaning of equipment or disposal of waste. Apply the product only as specified on this label.

### DIRECTIONS FOR USE

Read entire label before use. This product is a sulfur insecticide with its label.

**USE PRECAUTIONS:** CAUTION - Some fruits and other plants are susceptible to injury from sulphur. Do not apply to plants susceptible to sulphur injury. Do not apply to plants susceptible to sulphur injury. Do not apply to plants susceptible to sulphur injury. Do not apply to plants susceptible to sulphur injury.

CITRUS

**PEACHES, PLUMS:** To control BROWN ROT, POWDERY MILDEW and SCAB, apply 25 to 45 pounds per acre. Follow locally issued spray schedule for number and frequency of applications. Usually 6 applications are made: no. 1 - when petals have fallen; no. 2 - when shucks have shed; no. 3 - two weeks later; no. 4 - two weeks after no. 3; no. 5 - four weeks before harvest; and no. 6 - two weeks before harvest.

**APPLES:** To control APPLE SCAB, POWDERY MILDEW, apply 25 to 45 pounds per acre. Dust thoroughly between periods of rainfall. Repeat as necessary to cover new foliage growth during pink, petal fall and cover spray period. MacIntosh, Golden Delicious, Jonathan and certain other varieties may be controlled by sulphur applications under certain conditions.

**OUTDOOR FLOWER, ORNAMENTALS:** To control POWDERY MILDEW, LEAF SPOT and BLACK SPOT on ROSES, BOTRYTIS BLIGHT on HYDRANGEA, and RED SPIDERS, dust thoroughly and repeat application as necessary. For disease control, repeat applications on a 2-week basis.

**BEANS:** To control RUST and RED SPIDERS, dust thoroughly at the rate of 15 to 30 pounds per acre. Be sure dust is directed to underside of leaves. Repeat as necessary. For rust control, applications should be made at 7 to 10 day intervals. Do not apply to varieties of beans susceptible to sulphur injury. Do not apply if the temperature during or within 3 days after application is expected to exceed 90° F.

**PEANUTS:** To control LEAF SPOT, apply 20 to 25 pounds per acre at 7 to 10 day intervals or as necessary. Begin dusting when first appearance of leaf spot is observed at harvest.

**POULTRY (Chickens, Turkeys, Ducks, Geese, Game Birds and Pigeons):** To control DEPLUMING MITES, dust poultry with sulphur. Apply sulphur to poultry at least once a week. Do not apply to poultry if the temperature is expected to exceed 90° F.

### DISPOSAL

Do not contaminate water by cleaning of equipment or disposal of waste.

### NOTICE

This product is a sulfur insecticide with its label.

**Woolfolk Chemical Works, Inc.**  
Manufacturers Atmore, Alabama