

SECURITY

Nutonex Flowable Sulphur

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 769-484

ACTIVE INGREDIENT: Sulphur 52.5%
INERT INGREDIENTS: 47.5%
EPA Reg No 769-484 **EPA Est 769-GA-1** **TOTAL 100.0%**
 This product contains 6 pounds of Sulphur per gallon

KEEP OUT OF REACH OF CHILDREN.

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat, or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed or foodstuffs.

FIRST AID: If swallowed get medical attention. In case of contact immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

ENVIRONMENTAL HAZARDS: Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

USE PRECAUTIONS: CAUTION - Some fruits and other plants are susceptible to injury from sulphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality. Do not allow spray drift to reach cucumbers, squash or melons. During periods of high temperature, sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use SECURITY Nutonex Flowable Sulphur with oil or within 4 weeks of an oil application.

DIRECTIONS FOR MIXING: Shake container before pouring. For larger sizes, rock can back and forth several times before opening. The nature of this product is to thicken upon standing, and shaking will reverse this and allow it to pour smoothly. Pour the recommended amount of product into the water in the sprayer and agitate or stir until well mixed. If this mixture is allowed to stand for several hours, it may be necessary to agitate mixture again to redispense product before spraying. It is recommended that equipment be thoroughly cleaned after each day's use.

OUTDOOR FLOWERS AND ORNAMENTALS: To control POWDERY MILDEW, LEAF SPOT, BLACK SPOT ON ROSES, BOTRYTIS ON HYDRANGEA, and RED SPIDERS, apply SECURITY Nutonex Flowable Sulphur at the rate of 1 gallon per 100 gallons of water (2 1/2 tablespoons per 1 gallon) and repeat applications at 5 to 7 day intervals beginning when disease first appears and continuing until all danger from the disease has passed. During periods of humid or rainy weather, it may be necessary to shorten the application intervals.

BEANS: To control RUST and RED SPIDERS on beans, apply at the rate of 1 gallon per 100 gallons of water (2 1/2 tablespoons per 1 gallon). For RED SPIDERS, begin application at first sign of pests. Direct spray to undersides of leaves and repeat as necessary for control. For RUST begin applications when first leaves form or at first sign of disease. Repeat at 7 to 10 day intervals or as necessary for control. Sulphur may injure certain varieties of beans.

PEACHES AND PLUMS: To control BROWN ROT and SCAB, apply 2 to 3 quarts per 100 gallons water (2 tablespoons per 1 gallon). Follow locally issued spray schedules for number and frequency of application. Usually 6 applications are made: no. 1 - when 2/3 petals have fallen; no. 2 - when 1/2 shucks have shed; no. 3 - two weeks later; no. 4 - two weeks after no. 3; no. 5 - four weeks before harvest; and no. 6 - two weeks before harvest.

APPLES AND PEARS: To control SCAB and POWDERY MILDEW, apply 2 1/2 to 3 pints per 100 gallons water (1 tablespoon per gallon) during pre-bloom through calyx sprays and 2 pints per 100 gallons water (2 teaspoons per 1 gallon) during cover sprays. Spray for thorough coverage. Macintosh, Golden

Delicious, Jonathan and certain other apple varieties may be injured by sulphur applications under certain conditions.

CITRUS (grapefruit, lemons, limes, oranges, tangelos and tangerines): To control RUST MITE, apply 1 to 4 pints per 100 gallons water (1-4 teaspoons per gallon) depending on tree size. Spray for thorough coverage. Do not apply within 4 weeks of an oil application.

PEANUTS: To control RUST, apply 2 to 4 pints per acre (2-4 teaspoons per gallon of water per 100 feet of row) in sufficient water for thorough coverage. Begin applications approximately 6 weeks after planting. Repeat at 7 to 10 day intervals as long as rust control is required. Do not apply within 4 weeks on an oil application.

POTATOES: To control POWDERY MILDEW on commercial potatoes, apply 3 to 4 pints in 8 gallons of water per acre by air. For home garden, apply 3 to 4 teaspoons per gallon of water and apply to 100 feet of row. Apply at the first sign of infection and repeat at 3 week intervals.

STRAWBERRIES: To control POWDERY MILDEW, apply 1 pint per 100 gallons water (1 teaspoon per gallon). Apply at first sign of infection and repeat at 3 week intervals. To control RED SPIDERS, apply at the rate of 1/2 to 1 gallon per acre (12 1/2 to 5 1/2 tablespoons per gallon of water applied to 100 feet of row). Direct spray to undersides of leaves and repeat as necessary for control. For Strawberries that will be canned, discontinue application of sulphur well before harvest in accordance with local canner's recommendations.

TOMATOES: To control TOMATO RUSSET MITE, apply 1/2 to 1 gallon per acre (2 1/2 to 5 1/2 tablespoons per gallon of water) depending upon size of plants. Use full coverage sprays.

DISPOSAL

Do not reuse empty container. Wrap container and put in trash collection. For larger sizes, triple rinse container and pour rinsate into sprayer. Empty container should be disposed of in a safe place.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label. 086211

WOOLFOLK CHEMICAL WORKS, INC.
MANUFACTURERS

FORT VALLEY, GEORGIA 43