

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately. DO NOT induce vomiting.
IF INHALED: Remove victim to fresh air. Apply respiration if indicated.
IF ON SKIN: Wash immediately with soap and water.
IF IN EYES: Flush eyes for at least 15 minutes with water. Call a physician immediately.
NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

WARNING: Keep out of reach of children. May be fatal if swallowed. May be absorbed through skin. May be injurious to the eyes and skin. Do not get in eyes or on skin. Do not smoke while using. Wash thoroughly after handling. Wash contaminated clothing before reuse. Flush contaminated eyes with plenty of water and get medical attention. Avoid breathing spray mist or vapors. Do not spray vegetable gardens or other plants (except peach trees) used for food purposes. Keep away from food, feedstuffs, and domestic water supplies. Keep container closed. Keep children and pets off treated areas until spray has dried.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Keep out of any body of water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. This product is highly toxic to bees exposed to drift treatment or residues on plants. Protective information may be obtained from your Cooperative Agricultural Extension Service. Apply this product only as specified on this label.

PHYSICAL OR CHEMICAL HAZARDS

Irresistible. Do not use, pour, spill or store near heat or open flame. Handle concentrate in ventilated area.

DIRECTIONS FOR USE

is a violation of federal law to use this product in a manner inconsistent with its labeling.
USE ON TURF: Use SECURITY Dursban 2-E Lawn and Ornamental Spray to control fleah bugs, and weevils (lawn moths), ants, brown dog ticks, crickets, cutworms, burrows, and grasshoppers infesting home lawns and other ornamental and recreational turf areas. Mix SECURITY Dursban 2-E Lawn and Ornamental Spray with water according to directions given in the following table and apply as a coarse large droplet spray using a low pressure sprayer such as a knapsack type, hose-attached applicator or other similar equipment. If it is impractical to use the quantity of water listed in the table, apply in at least one gallon of water per 1,000 square feet and sprinkle with the recommended amount of water after application. For best results, lawn should be moist at the time of application and care should be taken to obtain complete and uniform coverage of pest infested areas. Spray from plants first appear, re-treat when needed.

DOSE AND DILUTIONS FOR USE ON TURF

Area to be treated	Amount of Dursban 2-E	
	Lawn and Ornamental Spray	Mix with Water
500 sq ft	1/4 fl. oz.	15 gallons
1,000 sq ft	1/2 fl. oz.	30 gallons
5,350 sq ft	7 1/2 fl. oz.	150 gallons
10,700 sq ft	1 pint	325 gallons
One Acre	2 quarts	1,300 gallons

fluid ounce = 2 Tablespoons = 6 teaspoons

Control certain white grubs (larvae of European chaffer and Japanese beetle), use two to six times the amount of SECURITY Dursban 2-E Lawn and Ornamental Spray shown in the table, i.e. 3 to 6 fl. oz. per 1,000 sq. ft. or 1 to 2 gallons per acre. Apply in enough water to obtain uniform coverage of the area treated followed immediately by thorough watering to wash the insecticide into the turf. For best results, treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service specialist for proper time to treat in your area. Control Hypodermis (cutworms) weevils, use twice the amount of SECURITY Dursban 2-E



Dursban 2-E

Lawn and Ornamental Spray

NOV 21 1980
 Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 769-485

ACTIVE INGREDIENTS:

Chlorpyrifos [0,0-dimethyl 0-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 24.60%
 Xylene range aromatic solvent 68.80%

INERT INGREDIENTS: 6.60%

EPA Reg. No. 769-485 EPA Est. 769-GA-1 TOTAL 100.00%

* Dursban is a registered trademark of the Dow Chemical Company.

This product contains 2 pounds chlorpyrifos per gallon.

KEEP OUT OF REACH OF CHILDREN.

WARNING

See side panel for additional precautionary statements.

ONE FL. QUART

WOOLFOLK CHEMICAL WORKS, INC.

Manufacturers

Fort Valley, Georgia 31030

Lawn and Ornamental Spray shown in the table, i.e. 1 gallon diluted in 1,300 gallons of water per acre. Spray suspected grubbing areas in mid-April and again in mid-May. Consult your local Agricultural Extension Service specialist for best time to treat in your area.
NOTE: To help prevent infestation of buildings by ants, crickets, earwigs, and cockroaches (pillbugs), treat a band of soil 6 to 10 feet wide around and adjacent to the building, and the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use the same dosage and dilution recommended for control of these pests on turf as shown in the table. Apply at a coarse spray to thoroughly and uniformly wet the band area.
FOR USE ON ORNAMENTAL PLANTS: Use SECURITY Dursban 2-E Lawn and Ornamental Spray to control aphids, bagworms, cutworms, grasshoppers, leafhoppers, mealybugs, beetles, spiderbugs, thrips (exposed), and whiteflies on flowers, shrubs, vines, shade and flowering trees and evergreens found to be infested with these pests. Dilute with water according to directions given in the following table and apply using a suitable low pressure sprayer such as a knapsack type or hose-attached applicator in a manner to provide complete and uniform coverage of infested plants. Avoid over-spraying to the point of excessive runoff. Treat when pests appear and repeat at 7 to 10 day intervals if needed.
NOTE: Do not use on azaleas, camellias, poinsettias, rose bushes or variegated ivy because of possible injury to these plants.

DOSE AND DILUTIONS FOR USE ON ORNAMENTAL PLANTS

PEST CONTROLLED	Amount of Dursban 2-E Lawn and Ornamental Spray in Water to Make		
	3 Gallons	12 Gallons	150 Gallons
Aphids, Bagworms, Grasshoppers, Meas., Spiderbugs, and Whiteflies	0.5 fl. oz.	1.6 fl. oz.	1 pint*
Cutworms, Mealybugs, Leafhoppers, and Thrips (exposed)	1 fl. oz.	3.2 fl. oz.	1 quart**

* Equivalent to 4 ounces of chlorpyrifos per 100 gallons

** Equivalent to 8 ounces of chlorpyrifos per 100 gallons

NOTE: Mosquitoes coming to rest on areas treated for control of turf and ornamental plant pests will be killed for varying periods of time after treatment, depending on exposure of treated areas to weathering conditions.

DOGWOOD BORER: To control DOGWOOD BORER, apply SECURITY Dursban 2-E to trunk and lower limbs in May or June before borers enter the trunk. Mix 2 ounces SECURITY Dursban 2-E with 1 gallon of water (1 pint to 8 gallons) and apply as a coarse, low pressure spray. Apply thoroughly, particularly to injured areas and old borer wounds.

PINES: To control infestations and help reduce damage to pines from SOUTHERN PINE BEETLE, apply SECURITY Dursban 2-E as a 1% spray by application to individual trees using suitable hand or power operated ground spray equipment. Mix 5% ounces of SECURITY Dursban 2-E with water to make up a total of 1 gallon of mixture (equivalent to 4.2 gallons in 100 gallons of water) and spray to the point of runoff. Do not mix with oil. For preventative treatment, apply the spray to the main trunk of pines in the early spring or when threat of attack exists from nearby infested trees. Repeat treatment as needed for continued protection. For remedial treatment, apply the spray to the main trunk of infested trees (or to infested logs) when damage occurs but before beetles begin to emerge.

PEACHES: To control PEACH TREE BORERS, apply SECURITY Dursban 2-E as a trunk spray before newly hatched borers enter the trees. Mix 2 ounces SECURITY Dursban 2-E with 1 gallon of water (1 pint to 8 gallons) and apply as a coarse, low pressure spray to give uniform coverage of tree trunks. Thoroughly wet all bark areas from ground level to scaffold limbs. Do not allow spray to contact fruit. Consult your State Agricultural Experiment Station or Extension Service specialist for proper time to treat in your area.

RESTRICTIONS: Make only one application per season. Do not apply within 14 days of harvest. Do not allow meat or dairy animals to graze in treated orchards.

DISPOSAL

Do not reuse empty container. Wrap container and place in trash collection.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

08500