

SECURITY® Fungi-Gard™

**BROAD SPECTRUM LIQUID FUNGICIDE
FOR USE ON ROSES, FLOWERS,
LAWNS AND VEGETABLES**

CONTAINS DICONIL 2787*

ACTIVE INGREDIENT
Chlorothalonil (p,p'-dichloro-3,4,5-trichlorobenzene) 11.24%
INERT INGREDIENTS: 88.76%
EPA Reg. No. 769-489 EPA Est. 769-GA-1 TOTAL 100.00%
*Covered under U. S. Patent No. 3,290,353 and No. 3,331,735
*DICONIL 2787 is a Reg. Trademark of Durrum Chemicals
This product contains 1.0 lb. Chlorothalonil per gallon.

KEEP OUT OF REACH OF CHILDREN.

DANGER

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Call a physician immediately.
IF INHALED: Remove victim to fresh air. Apply respiration if indicated.
IF ON SKIN: Wash immediately with soap and water.
IF IN EYES: Immediately flush eyes with plenty of water for 15 minutes. Seek medical attention for eyes immediately.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Corrosive. Causes severe eye damage. Do not get in eyes. Wear goggles or eye shield when handling this product. Keep out of reach of children. May be harmful if swallowed, inhaled, or absorbed through the skin. Avoid contact with skin, eyes and clothing. Do not breathe spray mist. Wash hands, arms and face thoroughly with soap and water before eating or smoking. Wash all contaminated clothing with soap and hot water before reuse. Do not contaminate feed or foodstuffs. Keep children and pets off treated areas until completely dry. Apply this product only as specified on this label.

NOTE: This product may produce temporary allergic side effects characterized by redness of the eyes, mild bronchial irritation and redness or rash on exposed skin areas. Persons having allergic reactions should contact a physician. Affected persons respond to treatment with antihistamines or steroid creams and/or systemic steroids.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Do not apply directly to lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste.

STORAGE AND DISPOSAL

STORAGE: Do not store where temperatures reach 32°F or product will freeze.
DISPOSAL: Do not reuse empty container. Wrap empty container and place in trash collection.

**NET WEIGHT
½ GALLON**

769-489

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
LAWN, ORNAMENTAL TURF:
SECURITY Fungi-Gard will control many turf diseases when used according to the rates below. Begin application when disease threatens or when conditions are favorable for disease development. Do not mow or water turf for 24 hours following treatment. Apply SECURITY Fungi-Gard in an adequate amount of water to provide complete coverage. This may vary from 5 to 15 gallons per 1000 square feet. Under severe disease conditions, use the curative rates and spray on a 7 day schedule. Do not graze treated areas. Do not feed clippings to livestock.

Disease	Spray Interval	Preventative*	Curative**
Copper Spot	7-10 days	1 1/2 to 2 1/2 pts	2 1/2 to 2 1/2 pts
Curculena Leaf Spot	7-10 days	1/4 to 1 1/2 pts	2 1/2 to 2 1/2 pts
Dollar Spot	7-14 days	1/4 to 1 1/2 pts	1 1/2 to 2 1/2 pts
Fairy Leaf Spot	7-10 days	1/4 to 1 1/2 pts	2 1/2 to 2 1/2 pts
Helminthosporium Leaf Spot and Melting Out	7-10 days	1/4 to 1 1/2 pts	1 1/2 to 2 1/2 pts
Large Brown Patch	7-10 days	1/4 to 1 1/2 pts	2 1/2 to 2 1/2 pts
Red Thread	7-10 days	1/4 to 1 1/2 pts	2 1/2 to 2 1/2 pts
Stem Rust of Bluegrass	7-14 days	1 1/2 to 2 1/2 pts	2 1/2 to 2 1/2 pts

*Recommended rates for preventing disease establishment. Use lower rates when disease conditions are light to moderate, higher indicated rates when conditions are severe.
**Rates for use on a 7 day schedule when disease is present. Higher indicated rates should be applied under severe conditions.

ORNAMENTAL FLOWERS

SECURITY Fungi-Gard will control a number of diseases of ornamentals. Apply 3 tablespoons per gallon of water (except ROSES, apply 2 tablespoons per gallon) beginning at first sign of disease and repeat at 7 to 14 day intervals throughout the season. Make applications on a 7 day schedule during periods of wet weather or severe disease conditions. Application should be thorough to obtain adequate coverage of the foliage.
Almond (flowering) Monarda blossom and branch blight
Almond (flowering) Monarda blossom and branch blight
Aster Leaf spots
Aster*** (RHODODENDRON SPP.) Outhwaite flower blight and phytophthora die back
Carnation Alternaria leaf spot and botrytis blight
Cherry Laurel Shot hole leaf spot
Chrysanthemum and Daisy Ascochyta ray blight, gray mold, and septoria leaf spot
Crabapple Cedar apple rust, sphaeria leaf spot, and scab
Dogwood Septoria leaf spot
Firethorn Scab
Geranium Botrytis blight and rust
Gradiolus Botrytis blight and curculena leaf spot
Hyacinth Rust
Hollyhock Rust
Iris Botrytis blight and leaf spot
Japanese Andromeda (Persea spp.) Phytophthora die back
Leather Leaf Fern (Florida only) Ascochyta blight, cercospora leaf spot, cyrtodactylus leaf spot, and rhizoctonia blight
Lily Botrytis blight
Mahonia Rust
Mountain Laurel Leaf Spot
Oyster Plant (RHODODENDRON) (Florida only) Tan leaf spot
Pelargonium*** Botrytis and phytophthora blight
Phloxendron (Florida only) Phytophthora blight and dactylospora leaf spot
Privet Cercospora leaf spot
Purple Leaf Sand Cherry Monarda blossom and branch blight
Quince (flowering) Monarda blossom and branch blight
Rose Apply 2 tablespoons per gallon of water for control of black spot and blossom blight
Verbena Powdery mildew
Zinnia Powdery mildew

***Discoloration of blooms has been noted, at least on certain varieties.

VEGETABLES

In the garden, apply SECURITY Fungi-Gard when conditions favor disease development. Follow the recommended application timing and intervals for the vegetables listed. Do not apply to any crop not listed on this label. Rates are expressed in tablespoons per gallon of water. Consult the chart below for the amount of dilute spray to apply for specific row spacing. All rates are based on a broadcast rate of 150 gallons per acre. During early stages of growth, proportionately less spray will be required. Always spray for thorough coverage. Follow the time interval between last application and harvest indicated by the number in () following each vegetable. A (0) indicates product can be used through day of harvest. Do not graze treated areas or feed treated plant parts or ensilage to livestock.

AMOUNT OF DILUTE SPRAY PER 100 FEET OF ROW WHEN ROW SPACING IS:

12 inches 1 gallon	24 inches 1/2 gallon	36 inches 1 gallon	48 inches 1 1/2 gallons
-----------------------	-------------------------	-----------------------	----------------------------

SNAP BEANS (7): To control RUST, apply 3 1/2 tablespoons per gallon of water. Begin application during early bloom or when disease first threatens. Repeat weekly or as necessary to maintain control.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS (8): To control ALTERNARIA LEAF SPOT (Southern states only) and DOWNY MILDEW, apply 2 tablespoons per gallon of water. Begin application after transplants are set in the field or shortly after emergence of field seeded crop or when conditions favor disease development. Repeat at 7 to 10 day intervals or as necessary to maintain control.

CANTALOUPE, HONEYDEW MELON, MUSKMELON, WATERMELON, PUMPKIN, SQUASH (9): To control ANTHRACNOSE and DOWNY MILDEW, apply 2 to 2 1/2 tablespoons per gallon of water. To control GUMMY STEM BLIGHT, LEAF BLIGHT, SCAB, CERCOSPORA LEAF SPOT and POWDERY MILDEW (in cantaloupes, muskmelon, honeydew, except in Southwestern states) apply 2 1/2 to 3 1/2 tablespoons per gallon of water. Begin when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat at 7 day intervals under severe disease conditions, shorten spray interval.

CARROTS (8): To control EARLY BLIGHT and LATE BLIGHT, apply 2 to 2 1/2 tablespoons per gallon of water. Start when disease threatens. Repeat at 7 to 10 day intervals or as necessary to maintain control.

SWEET CORN (14): To control HELMINTHOSPORIUM LEAF BLIGHT, apply 2 to 2 1/2 tablespoons per gallon of water. Begin when conditions favor disease development and repeat at 4-7 day intervals or as required to maintain control. For use on fresh market crop only. Do not apply to sweet corn to be commercially processed.

CUCUMBERS (8): To control TARGET SPOT (Florida only), ANTHRACNOSE and DOWNY MILDEW, apply 2 to 2 1/2 tablespoons per gallon of water. To control POWDERY MILDEW (except Southwestern states), GUMMY STEM BLIGHT, LEAF BLIGHT and SCAB, apply 2 1/2 to 3 1/2 tablespoons per gallon of water. For suppression of FRUIT ROT, apply 3 1/2 tablespoons per gallon of water. Begin applications when plants are in first true leaf stage or when conditions are favorable for disease development. Repeat applications at 7 day intervals. Under severe disease conditions, shorten spray interval.

ONIONS (7 days for dry bulb onions, 14 days for green onions): To control BOTRYTIS LEAF BLIGHT (blast), apply 2 to 3 1/2 tablespoons per gallon of water. Begin application before disease appears and repeat at 7 to 10 day intervals. Under severe conditions, use higher rate and shorten spray interval. Do not apply on sweet Spanish onions. Do not apply more than 3 times per season on green onions.

PEANUTS (14): To control CERCOSPORA LEAF SPOTS, apply 1 1/2 to 1 4/5 tablespoons per gallon of water. To control LEAF RUST, apply 1 4/5 tablespoons per gallon of water. Begin application when disease first appears. Repeat at 10 to 14 day intervals or as necessary to maintain control. Under severe disease conditions, use the higher rate.


POTATOES (8): To control EARLY BLIGHT, LATE BLIGHT and BOTRYTIS VINE ROT, apply 1 1/2 to 1 4/5 tablespoons per gallon of water. Begin applications when plants are 6 inches high or when disease threatens. Repeat at 7 to 10 day intervals or as necessary to maintain control. Under severe disease conditions, use the higher rate and shorten spray intervals.

TOMATOES (8): To control EARLY BLIGHT, LATE BLIGHT, GRAY LEAF SPOT and GRAY LEAF MOLD, apply 2 to 2 1/2 tablespoons per gallon of water. To control ANTHRACNOSE, apply 2 1/2 to 3 1/2 tablespoons per gallon of water. To control RHIZOCTONIA FRUIT ROT and BOTRYTIS GRAY MOLD, apply 3 1/2 tablespoons per gallon of water. Begin when disease threatens and repeat at 7 to 10 day intervals. Under severe disease conditions, shorten spray intervals.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties other than beyond the description on this label.

Manufactured by:
WOOLFOLK CHEMICAL WORKS, INC.

Fort Valley, GA 31030



SECURITY

Fungi-Gard

**BROAD SPECTRUM LIQUID FUNGICIDE
FOR USE ON ROSES, FLOWERS,
LAWNS AND VEGETABLES**

CONTAINS DICONIL 2787

ACTIVE INGREDIENT

INERT INGREDIENTS

1. Fungi-Gard is a broad spectrum liquid fungicide for use on roses, flowers, lawns and vegetables. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

KEEP OUT OF REACH OF CHILDREN.

DANGER

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink water or milk. Do not induce vomiting. IF INHALED: Move to fresh air. IF ON SKIN: Wash with soap and water. IF IN EYES: Flush with water for 15 minutes.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

DANGER: Fungi-Gard is a broad spectrum liquid fungicide for use on roses, flowers, lawns and vegetables. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

NOTE

Fungi-Gard is a broad spectrum liquid fungicide for use on roses, flowers, lawns and vegetables. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

ENVIRONMENTAL HAZARDS

STORAGE AND DISPOSAL

STORAGE: Store in a cool, dry place. DISPOSAL: Dispose of in accordance with local regulations.

NET WEIGHT
1 GALLON

769-487

DIRECTIONS FOR USE

LAWN, ORNAMENTAL, TURF

Fungi-Gard is a broad spectrum liquid fungicide for use on lawns, ornamentals, and turf. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Disease	Spray Interval	Rate Per 1000 Square Feet	Cartridge
Ascochyta Blight	14 days	1.0	1.0
Anthracnose	14 days	1.0	1.0
Botrytis	14 days	1.0	1.0
Downy Mildew	14 days	1.0	1.0
Leaf Spot	14 days	1.0	1.0
Root Rot	14 days	1.0	1.0
Rhizoctonia	14 days	1.0	1.0
Stem Rot	14 days	1.0	1.0
Take-All	14 days	1.0	1.0
Thielaviopsis	14 days	1.0	1.0
Verticillium	14 days	1.0	1.0
Wilt	14 days	1.0	1.0

ORNAMENTAL FLOWERS

Fungi-Gard is a broad spectrum liquid fungicide for use on ornamental flowers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Fungi-Gard is a broad spectrum liquid fungicide for use on ornamental flowers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Fungi-Gard is a broad spectrum liquid fungicide for use on ornamental flowers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Fungi-Gard is a broad spectrum liquid fungicide for use on ornamental flowers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Fungi-Gard is a broad spectrum liquid fungicide for use on ornamental flowers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Fungi-Gard is a broad spectrum liquid fungicide for use on ornamental flowers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

VEGETABLES

Fungi-Gard is a broad spectrum liquid fungicide for use on vegetables. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

AMOUNT OF DILUTE SPRAY PER 100 FEET OF ROW WHEN ROW SPACING IS

Feet	Spacing	Rate
18 inches	1.0	1.0
24 inches	1.0	1.0
30 inches	1.0	1.0
36 inches	1.0	1.0

SNAP BEANS (14): Fungi-Gard is a broad spectrum liquid fungicide for use on snap beans. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

CABBAGE, CAULIFLOWER, BROCCOLI, BRUSSELS SPROUTS (14): Fungi-Gard is a broad spectrum liquid fungicide for use on cabbage, cauliflower, broccoli, and Brussels sprouts. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

CANTALOUPE, HONEYDEW MELON, MUSHMELON, WATERMELON, PUMPKIN, SQUASH (14): Fungi-Gard is a broad spectrum liquid fungicide for use on cantaloupe, honeydew melon, muskmelon, watermelon, pumpkin, and squash. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

CARROTS (14): Fungi-Gard is a broad spectrum liquid fungicide for use on carrots. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

SWEET CORN (14): Fungi-Gard is a broad spectrum liquid fungicide for use on sweet corn. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

CUCUMBERS (14): Fungi-Gard is a broad spectrum liquid fungicide for use on cucumbers. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

ONIONS: 7 days for dry bulb onions, 14 days for green onions. Fungi-Gard is a broad spectrum liquid fungicide for use on onions. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

PEANUTS (14): Fungi-Gard is a broad spectrum liquid fungicide for use on peanuts. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

POTATOES (14): Fungi-Gard is a broad spectrum liquid fungicide for use on potatoes. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

TOMATOES (14): Fungi-Gard is a broad spectrum liquid fungicide for use on tomatoes. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

NOTE: Fungi-Gard is a broad spectrum liquid fungicide for use on vegetables. It contains Diconil 2787, which is a fungicide. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi. It is a liquid and is applied by spraying. It is a broad spectrum fungicide and is effective against a wide range of fungi.

Manufactured by
WOOLFOLK CHEMICAL WORKS, INC. Fort Valley, GA 31030