

SECURITY

SEVIN® 4 FLOWABLE

Controls Insects on Vegetables, Lawns, Ornamentals and Fruits.

ACTIVE INGREDIENT:

Carbaryl (1-naphthyl methyl carbamate) 42.85%

INERT INGREDIENTS: 57.15%

EPA Reg. No. 769-493 EPA Est. 769-GA-1 Total 100.00%

SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide

THIS PRODUCT CONTAINS 4 POUNDS CARBARYL PER GALLON

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air.

IF ON SKIN: Wash with soap and water. Remove any contaminated clothing.

IF IN EYES: Flush eyes with water.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS - HARMFUL IF SWALLOWED OR INHALED. Avoid breathing spray mist. Do not take internally. Avoid contact with skin and eyes. Wear regular long sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidotal. **DO NOT USE 2-PAM, OPIATES OR CHOLINESTERASE INHIBITING DRUGS.**

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic organisms. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

NET CONTENTS 2½ GALLONS

Manufactured by
WOOLFOLK CHEMICAL WORKS, INC.
Fort Valley, Georgia 31030

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

MIXING INSTRUCTIONS

To insure a uniform product, gently shake contents before pouring. Be sure sprayer tank and spray tips and screens are clean before use. Fill the sprayer with 1/2 to 3/4 the desirable amount of clean water. Pour the recommended amount of product into the tank and complete filling to desired level with water. Before and frequently during use, agitate the spray mixture in the tank to prevent settling. It is recommended that spray equipment be thoroughly cleaned after each day's use.

VEGETABLES

SECURITY Sevin 4 Flowable will control a wide range of vegetable insect pests. Begin applications when insects first appear and repeat applications to maintain control, usually a 7 day schedule. Do not apply to any crop not listed on this label. Coverage of both upper and lower leaf surfaces is essential for good control. Use sufficient gallonage to obtain full

Apply 1 quart per acre to control:

Alfalfa Caterpillar	Harlequin Bug	Squash Bug
Bean Leaf Beetle	Japanese Beetle	Three Cornered
Blister Beetle	Leafhopper	Alfalfa Hopper
Cucumber Beetle	Melonworm	Thrips
Colorado Potato Beetle	Mexican Bean Beetle	Velvetbean Caterpillar
Flea Beetle	Red-Necked Peanutworm	Western Bean Cutworm
Green Cloverworm		

Apply 1½ quarts per acre to control:

Armyworms	Cutworms	Pea Weevil
Aster Leafhopper	Grasshopper	Spittlebug
Corn Earworm	Pea Leaf Weevil	Yellow Striped Armyworm

Apply 2 quarts per acre to control:

Asparagus Beetle	Imported Cabbageworm	Southwestern Corn Borer
Chinch Bug	Lace Bug	Tarnished Plant Bug
Corn Rootworm	Lima Bean Pod Borer	Tomato Fruitworm
Cowpea Curculio	Lygus Bug	Tomato Hornworm
European Corn Borer	Stink Bug	Tomato Pinworm
Fall Armyworm	Stink Beetle	

SECURITY Sevin 4 Flowable may be applied to the vegetables listed below. Follow the time intervals between last application and harvest indicated by the number in () following each vegetable. A (0) indicates this product may be applied up to and including the day of harvest.

ASPARAGUS (1)	KOHLRABI (3)
BEANS (0) including dry beans, green beans, lima beans, navy beans, pinto beans	LETTUCE, head (3), leaf (14)
BROCCOLI (3)	MUSTARD GREENS (14)
BRUSSELS SPROUTS (1)	MELONS (0)
BEEFS (roots) (3)	OKRA (0)
CABBAGE (3)	PEANUTS (0)
CAULIFLOWER (3)	PEAS, SOUTHERN (0) including black-eyed, cowpeas and crowders
CHINESE CABBAGE (1)	PEAS, GARDEN or ENGLISH (0)
COLLARDS (14)	PEPPERS (0)
CARROTS (0)	POTATOES (0)
CORN (0)	PUMPKIN (0)
CUCUMBERS (0)	RADISH (3)
DANDELION (14)	RUTABAGA (3)
ENDIVE (ESCAROLE) (14)	SQUASH (0)
EGGPLANT (0)	SALSIFY (roots) (3), tops (14)
HANDS (0)	SPINACH (14)
	SWISS CHARD (14)
	WATERMELON (0)

ACCEPTED
FEB 12 1981
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for this pesticide registered under EPA Reg. No. 769-493

SPECIFIC VEGETABLE DIRECTIONS:

ASPARAGUS Treat ferns or brush growth. Do not treat more than once every 3 days.
CORN For insects attacking silks and ears, apply at 1-6 day intervals starting when first silks appear and continuing until silks begin to dry. For larvae in whorl and foliage feeders, apply as necessary. For chinch Bugs, use high gallonage application directed at the base of plants.

CUCUMBERS, MELONS, PUMPKINS, SQUASH Avoid excessive applications.

ORNAMENTAL PLANTS, ROSES, FLOWERS, SHRUBS AND SHADE TREES

SECURITY Sevin 4 Flowable will control the insects listed below. Apply 1 quart per 100 gallons of water and spray thoroughly covering both upper and lower leaf surfaces. Begin applications when insects or their damage first appear and repeat in 7-10 days or as necessary to maintain control. The addition of **SECURITY** Spreader-Sticker may improve residual control.

Ants	Grasshoppers	Plant Bugs
Apple Aphid	Gypsy Moth	Poinsettia Hornworm
Armyworm	Hackberry Nipple	Psyllids
Azalea Leafminer	Gall Maker	Puss Caterpillar
Bagworms	Holly Bud Moth	Redumped Oakworm
Birch Leafminer	Holly Leafminer	Rose Aphid
Blister Beetle	Jackpine Budworm	Rose Chaffer
Boxelder Bug	Japanese Beetle	Rose Slug
Brown and Leaf miner	Jeffery Pine	Saddled Prominent
Brown-tail Moth	Needleminer	Sawflies (exposed)
Cankerworms	June Beetles	Scale Insects
Catalpa Sphinx	Lace Bug	Scabbards
Chiggers	Leafhoppers	Spiny Elm Caterpillar
Cooley Spruce	Leafrollers	Springtails
Gall Aphid	Locust Borer	Spruce Budworm
Cutworms	Maple Leafcutter	Spruce Needleminer
Cypress Tip Moth	Mealy Bugs	Subtropical Pine
Douglas Fir	Mimosa Webworm	Tip Moth
Tussock Moth	Nantucket Pine	Tent Caterpillars
Eastern Spruce	Tip Moth	Thorn Bug
Gall Aphid	Oak Leafminers	Thrips (exposed)
Elm Leaf Aphid	Oak Leaf Skeletonizer	Ticks
Elm Leaf Beetle	Oakworm Complex	Walnut Caterpillars
Elm Spanworm	Oleander Caterpillar	Webworms
Eriophid Mites	Olive Ash Borer	Western Hemlock
European Pine	Orange Striped	Looper
Shoot Moth	Oakworm	Western Spruce
Fall Armyworm	Orange Tortrix	Budworm
Leaf Beetles	Periodical Cicada	Willow Leaf Beetles
Fly of Rose Beetle	Pine Sawfly	Yellow Poplar
Gall Midges	Pine Spittlebug	Weevils
Gall Wasps	Pitch Pine Tip	
Green Striped	Moth	
Mapleworm		

LAWN AND RECREATIONAL AREAS

SECURITY Sevin 4 Flowable will control the following insects on lawn and recreational areas:

Ants	European Chaffer	Meadow Spikes
Armyworms	Fall Armyworm	Side Webworms
Bluegrass Billbug	Fiery Skipper	Lawn Moths
Centipedes	Fleas	Scabbards
Chiggers	Grasshoppers	Springtails
Cutworms	June Beetles	Ticks
Earwigs	Leafhoppers	Yellow-striped Armyworm
	Larval Moth	Plant Bugs
	Moth	

SECURITY
SEVIN® 4
FLOWABLE

Controls Insects on Vegetables, Lawns, Ornamentals and Fruits.

ACTIVE INGREDIENT: Carbaryl (1-naphthyl methylcarbamate) 42.85%
INERT INGREDIENTS: 57.15%
 EPA Reg. No. 769-493 EPA Est. 769-GA-1 Total 100.00%
 SEVIN is the registered trademark of Union Carbide Corporation for carbaryl insecticide.
THIS PRODUCT CONTAINS 4 POUNDS CARBARYL PER GALLON.

STATEMENT OF PRACTICAL TREATMENT
IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.
IF INHALED: Remove victim to fresh air.
IF ON SKIN: Wash with soap and water. Remove any contaminated clothing.
IF IN EYES: Flush eyes with water.

KEEP OUT OF REACH OF CHILDREN.
CAUTION

PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS - HARMFUL IF SWALLOWED OR INHALED. Avoid breathing spray mist. Do not take internally. Avoid contact with skin and eyes. Wear regular long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. **NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidote. **DO NOT USE 2-PAM, OPIATES OR CHOLINESTERASE INHIBITING DRUGS.**

ENVIRONMENTAL HAZARDS
 This pesticide is toxic to aquatic organisms. Keep out of lakes, streams and ponds. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste.
PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or to residues remaining on the treated area. Do not apply when bees are actively visiting the crop cover crop or weed blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

NET CONTENTS 2 1/2 GALLONS
 Manufactured by
WOOLFOLK CHEMICAL WORKS, INC.
 Fort Valley, Georgia 31030

it is a violation of federal law

To insure a uniform product, get spray tips and screens are the amount of clean water. Pour to bring to desired level with water in the tank to prevent settling. Cleaned after each day's use.

SECURITY Sevin 4 Flowable applications when insects first appear on a 5-7 day schedule. Do not apply to and lower leaf surfaces is essential.

- Apply 1 quart per acre to control:
- Ailanthus Caterpillar
 - Bean Leaf Beetle
 - Blister Beetle
 - Cucumber Beetle
 - Colorado Potato Beetle
 - Flea Beetle
 - Green Cloverworm

- Apply 1 1/2 quarts per acre to control:
- Armyworms
 - Aster Leafhopper
 - Corn Earworm

- Apply 2 quarts per acre to control:
- Asparagus Beetle
 - Cinch Bug
 - Corn Rootworm
 - Cowpea Curculio
 - European Horn Borer
 - Fall Armyworm

SECURITY Sevin 4 Flowable intervals between last application on vegetable. A (D) indicates that **ASPARAGUS (1)**, **BEANS (C)** including dry beans, lima beans, navy beans.

- BROCCOLI (3)
- BRUSSELS SPROUTS (3)
- BEETS (roots) (3)
- CABBAGE (1)
- CARROT FLOWER (3)
- CHINESE CABBAGE (3)
- COLLARDS (1)
- CARROTS (C)
- CORN (B)
- CUCUMBERS (3)
- CANTALOUPE (1)
- EGGPLANT (1)
- FROGLIANT (1)
- HANOVER SAGE (3)

ACCEPTED