ROSES, FLOWERS, ORNAMENTALS

SECURITY Sevin Maneb Spray [illegible] below or ornamental plants, roses, flowers, [illegible] shade trees and also will control [illegible] plants listed. Apply SECURITY Sevin Maneb Spray [illegible] of 2½ tablespoons per gallon of water [illegible] gallons of water and spray [illegible] and lower leaf surfaces. For insects [illegible] insects or their damage first appear [illegible] intervals as necessary to maintain control [illegible] follow suggested timing of spraying [illegible] The addition of [illegible] SECURITY [illegible] per gallon of water [illegible] residual insecticide.

INSECTS CONTROLLED

Ants	[illegible]
Apple Aphid	[illegible]
Armyworm	[illegible]
Azalea Leafminer	[illegible]
Bagworms	[illegible]
Birch Leafminer	[illegible]
Blister Beetle	[illegible]
Blister Bug	[illegible]
Boxwood Leafminer	[illegible]
Brownta Moth	[illegible]
Cankerworms	[illegible]
Catalpa Sphinx	[illegible]
Chiggers	[illegible]
Cooley Spruce	[illegible]
Gall Aphid	[illegible]
Cutworms	[illegible]
Cypress Tip Moth	[illegible]
Douglas fir	[illegible]
Tussock Moth	[illegible]
Eastern Spruce	[illegible]
Gall Aphid	[illegible]
Elm Leaf Aphid	[illegible]
Elm Leaf Beetle	[illegible]
Elm Spanworm	[illegible]
Fall Webworm	[illegible]
European Pine	[illegible]
Shoot Moth	[illegible]
Fall Armyworm	[illegible]
Flea Beetles	[illegible]
Fuller Rose Beetle	Rose Chafer
Gall Midges	[illegible]
Gall Wasps	[illegible]
Green Stinkbug	[illegible]
Mealybugs	[illegible]
Mites	[illegible]
Leafhoppers	[illegible]
Gypsy Moth	[illegible]
Hickory Shuckworm	[illegible]
Japanese Beetle	[illegible]
Hickory Bark Moth	[illegible]
Hickory Horn	[illegible]
Hickory Shuckworm	[illegible]
Leaf Beetle	[illegible]
Leafhopper	[illegible]
Leafminer	[illegible]
Tent Caterpillar	[illegible]
Tussock Moth	[illegible]
Western Spruce Budworm	[illegible]
Bagworm	[illegible]
Willow Leaf Beetle	[illegible]
Mites	[illegible]
Aphids	[illegible]

DISEASES CONTROLLED

[illegible] RUST and STEM RUST [illegible] appear [illegible] days.

[illegible] ANTHRACNOSE [illegible] and BOTRYTIS BLIGHT [illegible] spray when new growth [illegible] and repeat weekly.

[illegible] To control ALTERNARIA BLIGHT or LEAF SPOT, ANTHRACNOSE and BOTRYTIS BLIGHT [illegible] spray when new growth [illegible] and repeat weekly.

[illegible] To control ANTHRACNOSE: spray tree [illegible] Begin when buds open. Repeat when [illegible] 4 weeks later and [illegible] summer.

[illegible] BOTRYTIS [illegible] before [illegible] are developing. Repeat [illegible] weekly intervals.

[illegible] Aer [illegible]

DISPOSAL

[illegible]

NOTICE [illegible]

MANUFACTURERS
WOOLFOLK CHEMICAL WORKS, INC.
Fort Valley, Georgia 31030

SECURITY.

Sevin-Maneb Spray

Controls Insects & Diseases on Vegetables, Fruits, Roses, Flowers and Ornamentals.

ACCEPTED MAR 25 1963

ACTIVE INGREDIENTS:

Carbaryl (1-napthyl methylcarbamate)	11.10%
Maneb (Manganese ethylene bisdithiocarbamate)	13.00%
(Total Manganese as metallic 2.68%)	
INERT INGREDIENTS:	75.90%
EPA Reg. No. 769-499 EPA Est. 769-GA-1	Total 100.00%

SEVIN is a registered trademark of Union Carbide Corporation for carbaryl insecticide.

This product contains 1 pound Carbaryl and 1.2 pounds Maneb per gallon.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air.

IF ON SKIN: Wash with soap and water. Remove any contaminated clothing.

IF IN EYES: Flush eyes with water.

KEEP OUT OF REACH OF CHILDREN.

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS: HARMFUL IF SWALLOWED OR INHALED. May cause irritation of eyes, nose, throat and skin. Avoid breathing spray mist. Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water. For eyes, get medical attention. Wear regular long-sleeved work clothing while spraying. Wash hands and face before eating. Wash thoroughly after handling. NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. DO NOT USE 2-PAM. OPIATES OR CHOLINESTERASE INHIBITING DRUGS.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of waste. Apply this product only as specified on this label.

PROTECT BEES: This product is highly toxic to bees exposed to direct treatment or to residues remaining on treated areas. Do not apply when bees are actively visiting the crop, cover crop, or weeds blooming in the treatment area. Applications should be timed to provide the maximum possible interval between treatment and the next period of bee activity.

ACCEPTED
APR 25 1963

bles, Lawns, Roses,

..... 11.10%
nate) 13.00%

..... 75.90%
A-1 Total 100.00%
de Corporation for carbaryl

1 1.2 pounds Maneb per

TREATMENT

er and induce vomiting by
induce vomiting or give
get medical attention.

Remove any contaminated

CHILDREN.

!

EMENTS

ANIMALS HARMFUL IF
on of eyes nose, throat and
ontact with skin, eyes or
ty of water For eyes, get
ed work clothing while
ng Wash thoroughly after
is a moderate, reversible
tal DO NOT USE 2-PAM,
RUGS

ZARDS

streams or ponds Do not
nly when weather conditions
minate water by cleaning of
product only as specified on

to bees exposed to direct
d areas Do not apply when
or weeds blooming in the
d to provide the maximum
ext period of bee activity

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

NOTICE: When mixing or loading this product all workers must wear the following protective clothing: long pants, long sleeved shirt, gloves, hat and boots.

MIXING INSTRUCTIONS

To insure uniform product, gently shake [illegible] SECURITY Sevin Maneb Spray [illegible] pressure sprayer such as a knapsack type, [illegible] hose [illegible] or other similar [illegible] equipment.

For HUSE 5% sprayers, it is recommended that this product be diluted 1 part product to 1 part water before filling applicator bottle. This will require setting the hose sprayer to apply twice the recommended label amount. If sample [illegible] apply 3½ tbsp. per gallon, the sprayer should be set to apply 7 tbsp. Check sprayer to see that it delivers the proper amount of material before using. Be sure the sprayer tank and spray plumbing are clean before use. For sprayers other than hose-on, fill the spray tank with ½ to ¾ the desired amount of clean water. Pour the recommended amount of product into the tank and complete filling to the [illegible] level with water. Before and frequently during use, shake or agitate the spray mixture in the tank to prevent settling. It is recommended that spray equipment be thoroughly cleaned after each day's use.

CONVERSION TABLE: 1 oz. = 2 tbsp. = 6 tsp.

VEGETABLES

Apply in sufficient volume of water to provide good coverage. In the garden, approximately one gallon of diluted spray will be required per 100 feet of row. This is equivalent to 193 gallons per acre. Direct spray to cover both upper and lower leaf surfaces. Apply this product at the rate of 2 gallons per acre or 3 1/3 tbsp. per 1 gal. of water. For insect control, begin application when insects or their damage first appear and repeat as necessary (except on harvest on asparagus only). For disease control follow the same directions for each crop. Do not feed forage or hay from treated crops to livestock.

PREHARVEST INTERVAL: Follow the [illegible] between application and harvest indicated by the [illegible] for each crop. A (0) indicates this product may be applied [illegible] including the day of harvest.

INSECT CONTROL

Alfalfa Caterpillar	[illegible]
Armyworm	[illegible]
Asparagus Beetle	[illegible]
Aster Leafhopper	[illegible]
Bean Leaf Beetle	[illegible]
Blister Beetle	[illegible]
Chinch Bug	[illegible]
Colorado Potato Beetle	[illegible]
Corn Earworm	[illegible]
Corn Flea Hopper	[illegible]
Cowpea Curculio	[illegible]
Cucumber Beetle	[illegible]
Cutworm	[illegible]
European Earwig	[illegible]
Fall Armyworm	[illegible]
Flea Beetle	[illegible]
Grasshopper	[illegible]
Green Cloverworm	[illegible]
Three Cornered Alfalfa Hopper	[illegible]
[illegible]	[illegible]
[illegible]	[illegible]

DISEASE CONTROL

[illegible]

LAWNS

[illegible]

INSECTS CONTROLLED [illegible] their [illegible]

Ants	Grasshoppers
Armyworm	June Beetles
[illegible]	Leafhoppers
[illegible]	Lucerne Moth
[illegible]	Mites
[illegible]	Mosquitoes
[illegible]	Sod Webworm
[illegible]	Springtails
[illegible]	Ticks
[illegible]	Yellow Striped Armyworm

DISEASE CONTROLLED [illegible] BROWN PATCH [illegible] Begin at first [illegible] 7 to 10 day intervals. To control [illegible] PUFFS begin when first pustules [illegible] and repeat at 7 to 14 day intervals.

AMOUNT TO USE

AREA TO TREAT	AMOUNT TO USE
[illegible]	[illegible]

ADDITIONAL DIRECTIONS FOR USE ON THE OTHER SIDE PANEL

NET CONTENTS
ONE GALLON