

**PRECAUTION STATEMENTS
HAZARDS TO HUMANS & DOMESTIC ANIMALS
DANGER**

Fatal if swallowed. Do not breathe dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly after using. Wash equipment and clothing after application. Keep children and pets off treated area until this material has been watered in. Air and grass dry. Do not allow domestic animals to graze treated area.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to wild life. Use with care when spraying in areas frequented by wild life. Keep out of wells, ponds, and streams. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with labeling.

SECURITY TURF-CAL FLOWABLE is for use as a pre-emergence control of crabgrass in established turf, such as lawns, parks, golf courses, and lawns located on industrial sites. (Do not use on Home Lawns.) **SECURITY TURF-CAL FLOWABLE** will also control **POA ANNUA** (Annual Bluegrass), chickweed and soil insect grubs (Japanese Beetle, Asiatic Beetle and white grub). The recommended rate for control of these soil insect grubs is 3 $\frac{1}{2}$ qts. per 1000 sq. ft. Consequently, when a rate of either 5 $\frac{1}{2}$ or 7 $\frac{1}{2}$ qts. per 1000 sq. ft. is used for weed control (refer to table below), control of grubs is effected.

TIME OF APPLICATION

SPRING Apply after ground is free from frost and well before crabgrass and Poa Annua germinate. March and April are the best months for treatment. Applications made later may not be fully effective that year.

EARLY FALL To control crabgrass and Poa Annua the following year, apply well before ground freezes.

APPLICATION AND AMOUNT TO USE Rake turf area free of leaves, mowing if necessary, so spray reaches soil surface. Measure area to be treated and uniformly spray required amount in sufficient water for thorough coverage. (Less treating with one half required amount in each direction will give best results.) Rain is not expected immediately after application. Sprinkle turf thoroughly. Apply as follows:

CRABGRASS AND POA ANNUA CONTROL

Type of Grass	Type of Soil	Amount per 1000 sq. ft.	When to apply
Kentucky Bluegrass	Well-drained	5 $\frac{1}{2}$ qts.	March to April
Bermuda, St. Augustine, and other warm season grasses	Well-drained	7 $\frac{1}{2}$ qts.	March to April

SECURITY
TURF-CAL FLOWABLE

ACTIVE INGREDIENT

Triphenyl Arsenate


INERT INGREDIENTS

E.P.A. Reg. No. 769-100 E.P.A. Reg. No. 769-100

This Arsenic is not for use in food or feed.

Arsenic in water should not be exposed to air or sunlight.

This product contains no lead and no other toxic substances.



DANGER - POISON

**KEEP OUT OF REACH OF CHILDREN.
STATEMENT OF PRACTICAL TREATMENT**

IF SWALLOWED Give several glasses of water and induce vomiting by administering Syrup of Ipecac or by touching back of throat with finger or blunt object. Repeat until vomit is clear. Then give 2 tablespoons Epsom Salts or Milk of Magnesia in water and plenty of milk and water. Have victim lie down and keep quiet. **CALL A PHYSICIAN IMMEDIATELY.** Symptoms of poisoning possible poisoning includes gaseous breath odor, difficulty in swallowing, burning in mouth, nose, throat and stomach, and profuse painful diarrhea.

IF IN EYES OR SKIN Flush with plenty of water. Get medical attention if irritation persists.

SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

182150

RETREATMENT After treating in the fall and again in spring, indicated on chart, apply at rate of 2 $\frac{1}{2}$ to 3 $\frac{1}{2}$ qts. per 1000 sq. ft. to maintain control of crabgrass and Poa Annua.

WHERE POA ANNUA IS A MAJOR PROBLEM SPRING TREATMENT Apply 3 $\frac{1}{2}$ to 5 $\frac{1}{2}$ qts. per 1000 sq. ft. Repeat treatment every 4 to 6 weeks between August 1st and October 1st. To maintain control of Poa Annua, annual treatments of 2 $\frac{1}{2}$ to 3 $\frac{1}{2}$ qts. of Turf Cal Flowable.

PRECAUTIONS Water Turf Cal Flowable in well water after application. This will get material into contact with soil and prevent any temporary scorching. Do not overlap areas when spraying. Keep any areas of turf away from lawn so that it sprouts lawn will not be harmed. Do not exceed rate of application recommended. Inadequate watering may cause some temporary scorching. Do not apply to Dichondra or St. Augustine lawns. Avoid contact with flowers and shrubs and applications in their immediate root zone. Do not apply Turf Cal Flowable on newly seeded areas until grass has become well established and has received a second cutting. Do not use a high phosphate fertilizer while program is in force. To help desirable grasses form, a high nitrogen fertilizer may be applied in spring and fall as needed. Apply lime as needed spring and fall but not at the same time as Turf Cal Flowable. Leave an interval of at least several months between applications of lime and Turf Cal Flowable.

SPECIAL NOTES: Kentucky, Merion Bluegrass, Zoysias, Bermudas and Ryegrass are more tolerant of arsenic than fescues and Bent grass.

TURF RENOVATION: Vertical cutting and over seeding should be encouraged, preferably before or even during the top of the turf. Plugging or vegetative improvement may also be desired.

STORAGE AND DISPOSAL

PROHIBITIONS: Do not use or store in or around the home. Do not contaminate water, food or feed by storage or disposal. Fertilizer dumping is prohibited. Do not reuse empty container.

PESTICIDE DISPOSAL Pesticide spray material should not be used except according to label instructions. Do not use or store in or around the home. Do not use or store in or around the home. Do not use or store in or around the home. Do not use or store in or around the home.

CONTAINER Triple rinse or equivalent, and offer for recycling or reconditioning, or dispose of in a sanitary landfill, or by other approved state and local procedures.

NOTICE: The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

Manufacturers

WOOLFOLK CHEMICAL WORKS, INC.

Fort Valley, Ga.