

1/4 769-501 PM 23

SUREGARD BRAND  
CAPTAN-MOLY SEED PROTECTANT  
WITH MOLYBDENUM  
15% NON-OIL LUBRICANTS ADDED

CE  
Per all Center  
OCT 13 1987  
769-501

174

ACTIVE INGREDIENTS:	
Captan -----	25.57%
Related Derivatives -----	0.58%
INERT INGREDIENTS: -----	73.85%
	100.00%

EPA Reg. No. 769-501                      EPA Est. 769-GA-1  
 \*\*Added ingredient Molybdenum 4.75%  
 Reg. USA Pat. Nos. 2,553,770; 2,553,771  
 Not licensed for use or export outside the United States,  
 its possessions, Canada or Mexico.

DANGER  
 KEEP OUT OF REACH OF CHILDREN

STATEMENT OF PRACTICAL TREATMENT  
 IF IN EYES: Flush with plenty of water. Call a physician.  
 IF SWALLOWED: Drink promptly a large quantity of milk, egg  
 whites, gelatin solutions, or if these are not available,  
 drink large quantities of water. Avoid alcohol. IF ON  
 SKIN: Wash with plenty of soap and water. IF INHALED: Move  
 victim to fresh air. If not breathing, give artificial  
 respiration, preferably mouth-to-mouth. Get medical  
 attention.

PRECAUTIONARY STATEMENTS  
 HAZARDS TO HUMANS AND DOMESTIC ANIMALS  
 DANGER! Causes irreversible eye damage. Harmful if  
 swallowed or inhaled. May cause allergic skin reactions.  
 Do not get in eyes. Wear goggles or face shield when  
 handling. Avoid contact with skin and clothing. Remove and  
 separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS  
 This pesticide is toxic to fish. Do not apply directly to  
 water or wetlands (swamps, bogs, marshes and potholes).  
 Drift and runoff from treated areas may be hazardous to  
 aquatic organisms in neighboring areas. Do not contaminate  
 water by cleaning of equipment or disposal of wastes. IF  
 the event that the seeds are spilled and cannot be salvaged,  
 completely cover them with soil.

ENDANGERED SPECIES RESTRICTIONS  
 The following restrictions apply to use of this product  
 after February 1, 1988.  
 Before using this pesticide on soybeans in the counties  
 listed below, you must obtain the PESTICIDE USE BULLETIN FOR  
 PROTECTION OF ENDANGERED SPECIES for the county in which the  
 product is to be used. The bulletin is available from your  
 County Extension Agent, State Fish and Game Office, or your

BEST AVAILABLE COPY

2/4  
pesticide dealer. Use of this product in a manner inconsistent with the PESTICIDE USE BULLETIN FOR PROTECTION OF ENDANGERED SPECIES is a violation of federal laws.

ALABAMA (Lauderdale, Limestone and Madison); ARIZONA (Graham, Maricopa, Mohave, Pima, Pinal and Santa Cruz); ARKANSAS (Benton and Polk); CALIFORNIA (Imperial, Modoc and Riverside); MISSISSIPPI (Claiborne, Copiah and Hinds); MISSOURI (Barry, Benton, Camden, Christian, Dallas, Greene, Hickory, Jasper, Lawrence, Miller, Newton, Osage, Polk, St. Clair, Stone and Webster); NEVADA (Clark); NEW MEXICO (Chaves, DeBaca and Eddy); OHIO (Pickaway); OKLAHOMA (Delaware, McCurtain and Pushmataha); OREGON (Lake); TENNESSEE (Lawrence and Wayne); TEXAS (Bastrop, Burleson, Comal, Harris, Hays, Jeff Davis, Pecos and Reeves); UTAH (Utah and Washington).

#### PRODUCT INFORMATION

Captan-Moly is especially formulated for planter box treatment of soybean seed to control seed and soil-borne diseases, including damping-off, seed decay organisms and seedling blights and to provide essential Molybdenum for plant growth where soils have been found to be deficient in available Molybdenum. Molybdenum helps the nodule bacteria and plants make more efficient use of nitrogen. It is prepared for use directly in the planter box, thereby eliminating seed handling to have seed treated.

A non-oil lubricant has also been added to help prevent cracking and splitting of the seed so they will flow easily and evenly from the planter box.

This material provides adequate coverage of seed and any excess flows with the seed into the furrow gives additional protection against soil-borne diseases.

DO NOT USE TREATED SEED FOR FOOD, FEED OR OIL PURPOSES.

#### RECOMMENDATIONS

Use as a planter box treatment at the rate of 4 oz. per bushel of seed. Plant soybeans at the rate of at least one bushel per acre, or the amount of Molybdenum applied to each acre will not be adequate.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

**BEST AVAILABLE COPY**

USE PRECAUTIONS: Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse. Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron. A dust mask must be worn when mixing and loading this product.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

Mix thoroughly with seed in planter box as follows: Fill the box one-half full with seed then treat with one-half the amount of Captan-Moly and mix well by stirring with a stick or paddle until color is imparted uniform to all seed; add the remaining seed to box and cover with balance of Captan-Moly. Mix thoroughly as above.

PLANTER BOX RATES ARE AS FOLLOWS

SIZE OF PLANTER BOX	NO. 1 OZ. SCOOPS PER BOX
1/2 bushel	2 heaping scoops
1 bushel	4 heaping scoops

Plant at the rate of at least one bushel soybeans per acre.

For further specific information, consult the State Agricultural Experiment Station or the State Agricultural Extension Service.

STORAGE AND DISPOSAL

STORAGE: Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks. PESTICIDE

DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous. Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. CONTAINER DISPOSAL: Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by


BEST AVAILABLE COPY

4/4  
incineration or, if allowed by state and local authorities,  
by burning. If burned, stay out of smoke.

NOTICE: DISCLAIMER OF WARRANTY, MERCHANTABILITY AND  
LIABILITY - If for any reason this product is not used,  
applied, stored, transported or disposed of in accordance  
with the instructions set forth on this label, then and  
thereafter SureCo, Inc. terminates, ends and excludes all  
warranties, expressed or implied, including the warranty of  
merchantability and fitness.

NET WEIGHT: 10 LBS.

MANUFACTURED FOR  
SURECO, INC.  
FORT VALLEY, GA 31030



**BEST AVAILABLE COPY**