Maria laga

EPA REGISTRATION NO. DATE OF ISSUANCE US ENVIRONMENTAL PROTECTION AGENCY SEP 3.0 1986 769-516 TERM OF ISSUANCE OFFICE OF PESTICIDES PROGRAMS REGISTRATION DIVISION (TS-767) WASHINGTON, DC 20460 NAME OF PESTICIDE PRODUCT NOTICE OF PESTICIDE: REGISTRATION REREGISTRATION Security brane incremane (Under the Federal Insecticide, Fungicide, Dispersible willing and Rodenticide Act, as amended) NAME AND ADDRESS OF REGISTRANT (Include ZIP code) Security nawn & Garden Products Co. P. U. DOX 938 Fort Vailey, 55 31,36 ┙ NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number. On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act. A copy of the labeling accepted in connection with this Registration/Reregistration is returned berewith. Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered based on your response to the logistration Standord outdance Fachage, aPA has reregistered the product listed above. This registration is manted under the authority of section 3(q) of FIFRA, as amended. The labeling referred to above, submitted in connection with registration under FIFAA, is acceptable subject to the comments listed below. Five copies of the limished labeling must be succitted. Incorporate the required language from Packotice (3-1, far worker) talety. I. Non-the remodeling payrical mazard statement: Sulfur dust suspensed in air lightles cosily. Kee, aday from hear, starks, or flame. be not smore while applying this product. Henry II. Jacoby Product Manager (21) Fungacide-Herbiciae branch Legistration Division (15-7070) ATTACHMENT IS APPLICABLE SIGNATURE OF APPROVING OFFICIAL

PREVIOUS EDITION MAY BE USED WITH SUPPLY IS EXHAUSTED.

88773: Jacoby: J-3: KENCO: 9/18/86: 9/30/86:bjp: vo

EPA Form 8570-6 (Rev. 5-76)

١,

SECURITY BRAND MICROFINE* DISPERSIBLE SULPHUR

ACCRATED
with COMMENTS
in EPA Letter Dated:
REREGISTERED

KEEP OUT OF REACH OF CHILDREN CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS: Harmful if swallowed. Avoid breathing dust. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin and clothing. Do not contaminate feed and foodstuffs. Do not use for medical purposes or as a feed supplement.

FIRST AID: IF IN EYES, immediately flush with plenty of water for at least 15 minutes. Call a physician if irritation persists. FOR SKIN CONTACT, wash with soap and water. IF INHALED and overcome by vapor or dust, remove immediately from exposure and call a physician. Administer aritificial respiration if breathing is irregular or has stopped. IF SWALLOWED, drink 1 to 2 glasses of water and induce vomiting by sticking finger in back of throat. Do not induce vomiting or give anything by mouth to an unconscious person. Call a physician.

ENVIRONMENTAL HAZARDS: Do not apply directly to water. Do not contaminate water by cleaning or equipment or disposal of waste. Apply this product only as specified on this label.

SE PRECAUTIONS: CAUTION - Some fruits and other plants are susceptible to injury from Alphur under certain climatic conditions. The user is advised not to use sulphur on any crop unless local use has proven that sulphur is safe in that locality. During periods of high temperatures, sulphur may burn foliage and fruit. Do not make sulphur applications at excessively high temperatures. Do not use sulphur with oil or within 3 weeks after oil application. When using Lead Arsenate with Security Microfine Dispersible Sulphur, always add the same amount of hydrated lime or other safener as would be used when applying Lead Arsenate without sulphur. If the crop to be treated will be used for commercial canning or freezing purposes, consult food processor before applying sulphur.

REENTRY INTERVAL: Do not reenter treated fields within 24 hours.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

GENERAL INFORMATION: When more than one dosage range is given, use the higher rate when conditions for diseases are most favorable and use the lower rate when conditions for diseases are least favorable.

APPLES: To control POWDERY MILDEW and SCAB, use 8 to 24 lbs. of Security Microfine Dispersibl Sulphur per acre in sufficient water for good coverage. Apply in prebloom, petal fall and cover periods as necessary. NOTE: MacIntosh, Golden Delicious, Jonathan and certain other varieties may be injured by sulphur.

BLACKBERRIES AND OTHER BRAMBLES: To control POWDERY MILDEW, apply 5 to 12 lbs. of Security Microfine Dispersible Sulphur per acre in sufficient water for good coverage...Begin when new shoots are 6 inches long and before blossoms open. Repeat at 10-day intervals or as recessary.

BLUEBERRIES, HUCKLEBERRIES: To control POWDERY MILDEW, apply 4 to 8 lbs. per 100 gallons of water. Begin when disease is expected to appear. Repeat at 7 to 10 day intervals or as necessary.

CHERRIES: To control POWDERY MILDEW, LEAF SPOT and BROWN ROT, apply 15 to 18 pounds per acre to TART CHERRIES and 20 to 24 pounds per acre to SWEET CHERRIES. Use sufficient vater for good coverage and apply as necessary from bloom through cover periods or as rected by your state spray guide.

CITRUS (ALL): To control RUST MITE, apply 5 to 7 lbs. per 100 gallons of water. Apply as necessary for control. Under heavy mite pressure or during cool weather, use the higher rate.

GRAPES: To control POWDERY MILDEW, apply 1.6 to 4 lbs. of Security Microfine Dispersible Sulphur per acre in sufficient water for good coverage. To control RUST, apply 40 lbs. per acre. Begin when new shoots are 6 to 10 inches long. Repeat before blooms open and continue at 10 to 14 day intervals as necessary. NOTE: Concord and other labrusca type grapes may be injured.

MANGOES: To control POWDERY MILDEW, apply 4-3/4 lbs. per 100 gallons of water. Begin before bloom. Repeat in bloom, after fruit set, and 3 weeks later.

NECTARINES, PEACHES: For control of BROWN ROT, SCAB and POWDERY MILDEW, apply 8 to 24 lbs. per acre of Security Microfine Dispersible Sulphur in sufficient water for good coverage. Apply from pink through cover sprays. A coarser, less adherent sulphur such as Security Nutonex or Security Wettable Sulphur is preferred in the preharvest sprays.

PEARS: For control of POWDERY MILDEW and SCAB, apply 8 to 24 lbs. per acre of Security Microfine Dispersible Sulphur in sufficient water for good coverage. Apply from prebloom through cover period as necessary.

STRAWBERRIES: To control POWDERY MILDEW, apply 4 to 8 lbs. per acre of Security Microfine Dispersible Sulphur in sufficient water for good coverage. Begin at first sign of disease and repeat as necessary. NOTE: Sulphur may injure strawberries under certain climatic conditions.

BEANS, PEAS: To control POWDERY MILDEW and RUST, apply 6 to 8 lbs. per acre in sufficient water for thorough coverage. Begin applications at first sign of infection and repeat in 7 to 14 days as necessary. Keep new growth protected.

POTATOES: To control POWDERY MILDEW, apply 2.4 to 4.8 lbs. of Security Microfine Dispersible Sulphur per acre in sufficient water for good coverage Begin before disease appears and repeat as necessary.

ROSES: To control BLACK SPOT, BROWN CANKER, LEAF SPOTS, POWDERY MILDEW, and RUSTS, apply 3.8 to 10 lbs. per 100 gallons of water and spray for thorough coverage. Begin when growth starts in spring. Repeat at 7 day intervals and after rains.

STORAGE AND DISPOSAL

Pesticide, spray mixture or rinse water that cannot be used according to label instructions must be disposed of according to applicable federal, state or local procedures. Completely empty bag by shaking and tapping sides and bottom to loosen clinging particles, Empty. residue into application equipment. Then dispose of bags in a sanitary landfill or by.... incineration if allowed by state and local authorities.

NOTICE

The seller makes no implied warranty of merchantability nor any other warranties which extend beyond the description on this label.

WOOLFOLK CHEMICAL WORKS, INC.
MANUFACTURERS
FORT VALLEY, GEORGIA