



# Sevin Bait

## Controls Cutworms, Sowbugs, on Vegetables and Mole Crickets, Armyworms on Lawns

**ACTIVE INGREDIENT**  
Carbaryl (1-naphthyl methyl carbamate) 95.0%

**INERT INGREDIENTS** 5.0%

EPA Reg. No. 729-517 EPA Est. 269-0A-1 Fort Valley, GA 31030

Sevin is the registered trademark of or one of the trademarks of the Carbaryl Insecticide.

### KEEP OUT OF REACH OF CHILDREN CAUTION

#### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED** Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

**IF INHALED** Remove victim to fresh air. **IF ON SKIN** Wash with soap and water. Remove any contaminated clothing. **IF IN EYES** Flush eyes with water.

**NOTE TO PHYSICIAN** Carbaryl is a moderate inhibitor of cholinesterase. Inhibit Atropine, 1 mg. intravenously. Do not use if FAM plates or cholinesterase inhibiting drugs.

#### PRECAUTIONARY STATEMENTS

##### HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**HARMFUL IF INHALED OR SWALLOWED** Avoid breathing dust. Do not take internally. Avoid contact with skin and eyes. Wear long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating. Wash thoroughly after handling. Keep bait away from domestic animals.

**NET WEIGHT**  
5 lbs

#### ENVIRONMENTAL HAZARDS

Avoid the application of this bait to rivers and ponds. Do not apply when weather conditions favor drift into areas that may contaminate water by spraying, dusting, or disposal of waste.

#### DIRECTIONS FOR USE

In accordance with Federal Law, use this product only in manner consistent with its labeling.

#### VEGETABLES

SEVIN Bait will control cutworms, armyworms, mole crickets, darkling ground beetles, grasshoppers, mole crickets and sowbugs when applied in a broadcast application to the listed vegetables. Repeat treatments are necessary. Follow the time interval between last application and harvest indicated by the following table. A full crop of vegetables may be used the day of harvest.

**Cucumbers (0), Melons (0), Peanuts (0), Squash (0)** Apply 20 pounds per acre or 1 lb. per 1000 square feet.

**Peas (0)** Apply 20 pounds per acre or 1 lb. per 1000 square feet.

**Beans (0), Carrots (0), Corn (0), Eggplants (0), Okra (0), Peppers (0), Potatoes (0), Tomatoes (0), Sweet Corn (0), Asparagus (1), Strawberries (1), Broccoli (3), Brussels Sprouts (3), Cabbage (3), Cauliflower (3), Lettuce (Head 3, Leaf 14), Garden Beets (Roots 3, Tops 14), Horseradish (3), Parsnips (3), Radishes (3), Rutabagas (3), Turnips (Roots 3, Tops 14), Collards (14), Endive (Escarole) (14), Kale (14), Parsley (14), Spinach (14) and Swiss Chard (14)** Apply 40 pounds per acre or 1 lb. per 1000 square feet.

#### LAWN TURF & ORNAMENTALS

To control cutworms, armyworms, crickets, darkling ground beetles, grasshoppers, mole crickets and sowbugs, apply 20 to 40 pounds per acre or 1 lb. per 1000 square feet.

**Lawn and Turf** Treat entire area evenly with a spray or dust or spreader or similar applicator to obtain uniform coverage. Paying close attention to edges of sidewalks and along paths and patios. Foundations. For best results, lawns should be cut and watered thoroughly before treatment.

**Ornamental Plants** Apply evenly to shrubs and plants and to trees at a rate of 1 pound per 1000 square feet. Trees and shrubs to be treated should be dormant.

#### DISPOSAL

Do not reuse container. Wrap in plastic and place in airtight container.

#### NOTICE

Sevin Bait is sold with a limited warranty. For details, see the full label. Warranties which extend beyond the label are not available.

#### Manufacturers

**WOOLFOLK CHEMICAL WORKS, INC.**  
Fort Valley, Georgia 31030