

SECURITY BRAND
EZE (R)
ORNAMENTAL & FRUIT
WEED PREVENTER

Controls Weeds and Grasses in Roses, Ornamentals and 16 Fruit and Nut Crops
CONTAINS CASORON (R)

ACTIVE INGREDIENT:

Dichlobenil* ----- 2.0%

INERT INGREDIENTS: ----- 98.0%

*2,6-dichlorobenzonitrile, U. S. Patent 3,027,248. TOTAL 100.0%

EPA Reg. No. 769-536

EPA Est. 769-GA-1

CASORON is a Reg. TM of Duphar, B.V., Amsterdam, Holland.

KEEP OUT OF REACH OF CHILDREN

CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting by drinking water and touching back of throat with finger. IF
INHALED: Remove to fresh air. IF IN EYES: Rinse with water for 15 minutes. IF ON SKIN:
Wash with soap and water.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION. HARMFUL IF SWALLOWED. Do not breathe dust. Do not allow contact with eyes or
skin. Do not allow livestock to graze in treated areas.

ENVIRONMENTAL HAZARDS

Do not apply directly to any body of water. Do not contaminate water by cleaning of equipment
or disposal of waste.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its
labeling.


DO NOT USE IN VEGETABLE GARDENS

IMPORTANT: PRECAUTIONS FOR USE. Do not apply this product to plants under unusual stress
resulting from drought, frost, insects, nematodes, etc. as injury may result. Do not use
on sloping areas as run-off may result in injury to lawn or other plants in adjacent areas.
Do not use on plants not listed on this label. Do not apply to transplant or cutting beds
or to newly set transplants and do not seed or transplant into treated areas. Do not use
on plants, any part of which is used for human or animal consumption except as listed on this
label. Do not use on extremely light soils. Once treatment is made, do not disturb the
soil in treated area as long as weed control is being maintained.

WEEDS CONTROLLED

When applied as directed, this product will give selective control of annual grassy and
broadleaved weeds and certain perennial weeds including:

Alligator Weed	Foxtail	Purslane
Annual Bluegrass (poa annua)	Gisekia	Ragweed
Bluegrass	Groundsel	Red Deadnettle
Bull Thistle	Henbit	Redroot Pigweed
Camphorweed	Horsetail	Rosarypea
Carpetweed	Jerusalem oak Goosefoot	Russian Thistle
Chickweed	Lotweed	Shepherdspurse
Citron Melon	Lambsquarter	Smartweed
Clovers	Laypop	Spanish Needles
Coffeeweed	Milkweed Vine	Spurge
Crabgrass	Miners Lettuce	Spurweed
Cudweed	Mugwort	Teaweed
Dandelion	Natalgrass	Texas Panicum
Dogfennel	Nutgrass	Wild Barley
Evening Primrose	Old Witchgrass	Wild Mustard
Fiddleneck	Peppergrass	Wild Radish
Florida Betony	Pineappleweed	Yellow Woodsorrel
Florida Purslane (pusley)	Plantain	



WOODY ORNAMENTALS

Apply this product only to established woody plants - those which have been planted for at least four months. Security EZE Ornamental & Fruit Weed Preventer may be used for weed control on the following ornamental plants:

Amur Corktree	Dogwood	Ivy (Hedera Helix)	Pecan
Arboe Vitae	Eastern Red Cedar	Juniper	Photinia
Ash	Elm	Kinnikinnick	Pine
Azalea (1)	Elwood Cypress	Laurel	Pittosproim
Barberry	English Walnut	Leucothoe	Poplar
Beauty Bush	Euonymous	Lilac	Privet
Birch	Filbert	Linden	Pyracantha
Boxelder	Flowering Almond	Locust	Rhododendron
Boxwood	Flowering Quince	Magnolia	Rockrose
Camellia	Forsythia	Maple	Rose
Caragana	Gardenia	Mock Orange	Russian Olive
Cleyera Japonica	Goldenraintree	Mountain Ash	Spiraea
Cotoneaster	Hackberry	Nandina	Sqaw-carpet
Cottonwood	Heather	Oak	Weigela
Crabapple	Holly (2)	Osmanthus (Holly-Olive)	Willow
Dentzia	Honeysuckle		Yew (Toxus)
		Pachistima	

(1) Krume, Mollis, Hardy Hybrid types and hardy native species only.

(2) Except I. Crenata, I. Rotunda, I. Vomitoria

BEFORE APPLICATION TO ORNAMENTALS: Remove existing weeds. Thoroughly work up and smooth the area to be treated. Carefully measure area to be treated to determine the number of square feet. If the area is large, it should be divided into smaller sections and each section treated individually.


APPLICATION AND RATE: Apply SECURITY EZE Ornamental & Fruit Weed Preventer uniformly to the soil at the rate of 3/4 to 1 ounce (2 to 2-2/3 tablespoons) for each 10 square feet to be treated. This is equivalent to 22 to 33 ounces per 300 square feet. Use the HIGHER RATE to control perennial and hard-to-control weeds. For circular areas around individual shrubs or trees, apply the amount indicated in the following table:

DIAMETER OF CIRCLE	RATE OF PRODUCT
2 feet	2 - 3 level teaspoons
3 feet	4 - 6 level teaspoons
4 feet	2-1/2 - 3-2/3 tablespoons
5 feet	4 - 6 level tablespoons
6 feet	5-2/3 - 8-1/2 tablespoons

Each ounce of SECURITY EZE Ornamental & Fruit Weed Preventer equals 8 level teaspoons. 3 teaspoons = 1 tablespoon.

Rates higher than those indicated may cause injury. More accurate and uniform application can be obtained by applying 1/2 of the required amount of material over the entire area and repeating the application with the remaining material.

As a guide for the proper application rate (lower rate) and uniformity of distribution of the granules, consult the diagram below.



AFTER APPLICATION: Unless the soil is wet, sprinkle lightly to moisten the soil 1/4 to 1/2 inch deep. Do not flood, avoid puddling and washing. Avoid disturbing soil in the treated area as long as weed control is being maintained.



FRUIT AND NUT CROPS

To control the weeds listed above, apply this product to the crops listed in EARLY SPRING before weeds germinate or after cultivation has removed the growing weeds. Distribute granules uniformly over the soil surface. This product works best in cool weather. For SOUTHERN AREAS, best results are obtained by mixing the granules with the upper inch of soil immediately after application. In other areas, shallow incorporation or sprinkling with water is suggested when applications are made during periods of high temperatures.

ALMOND, APPLE, BLUEBERRY, CHERRY, GRAPE, MANGO, NECTARINE, PEACH, PEAR, PRUNE, PLUM, FILBERT, PECAN and ENGLISH WALNUT: This product may be used on bearing, non-bearing and nursery stock. Do not apply within 4 weeks after transplanting (6 months on pecan transplants). Do not apply within one month of harvest of nuts or stone fruit. Apply 4-1/2 to 7 lbs. per 1000 square feet. For hard-to-control and perennial weeds, use the higher rate. For treating circular areas around individual fruit trees, follow the chart and rates provided in the ORNAMENTALS section.

BLACKBERRY and RASPBERRY: Use only on established plantings. Do not apply during new shoot emergence. Use 4-1/2 lbs. per 1000 square feet.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not store with propagative materials such as seeds, bulbs, tubers, nursery stock, etc. or with feed or food products.

Keep container tightly closed while in storage. Do not reuse empty container. Wrap container and place in trash collection.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

NET WEIGHT: 2-1/2 POUNDS

WOOLFOLK CHEMICAL WORKS, INC.

MANUFACTURERS

FORT VALLEY, GEORGIA