


SECURITY BRAND
PURGE 2
LAWN WEED KILLER

FOR ST. AUGUSTINE GRASS, CENTIPEDE GRASS, AND DORMANT BERMUDA GRASS



ACTIVE INGREDIENTS:

Atrazine: 3-chloro-4-ethylamino-6-isopropylamino-s-triazine	22.4%
Related Compounds	1.2%
INERT INGREDIENTS:	76.4%
	TOTAL 100.0%

EPA Reg. No. 769-

EPA Est. 769-GA-1

This product contains 2 lbs. Atrazine per gallon.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Induce vomiting and seek medical attention immediately. IF IN EYES OR ON SKIN: Flush eyes with water; wash skin with soap and water.

PRECAUTIONARY STATEMENTS

HAZARDOUS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful or fatal if swallowed. Do not get in eyes, on skin or on clothing. Avoid inhalation of vapors or spray mist. Do not contaminate feed or food.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

SECURITY Purge 2 Weed Killer controls spurweed, Florida betony, annual bluegrass and many other problem weeds in home lawns, ornamental and recreational turf areas.

GENERAL INFORMATION

SECURITY Purge 2 is a flowable weed killer that should be mixed with water and applied uniformly to the lawn. SECURITY Purge 2 may be applied with any pump-up or compressed air type sprayer or with a hose-on type sprayer. Shake well before using.

SECURITY Purge 2 will control both emerged weeds and weeds from seed. Rain or watering within 2 to 3 days of application may decrease the effectiveness on emerged weeds. However, for control of weeds from seeds, rainfall or watering is necessary within 7 to 10 days after treatment.

CAUTION: Do not apply within the active root zone of desirable ornamental plants such as camellias, azaleas, boxwoods, etc. Do not plant any crop (flower or vegetable gardens) to treated areas for 18 months or injury may result. Do not apply to grass that has been subjected to stress conditions such as drought, excessive moisture, winter injury, insects, diseases, nutrient imbalance or chemical injury. Do not apply when temperatures are expected to exceed 90° F. Do not apply in combination with liquid fertilizer or other pesticides unless experience indicates that they are compatible. Do not apply to plant species not listed on the label.

WEEDS CONTROLLED OR SUPPRESSED

Annual bluegrass (*Poa annua*), chickweed (common and mouseear), crabgrass, dandelion, cudweed, dichondra, Florida betony, henbit, knotweed, lespedeza, moneywort, nasturtium, narrowleaf vetch, parsley-piert, sandspur, smutgrass, spurge, spurweed, wintergrass, woodchick, various annual clovers, and fescue grass.

TIMING OF APPLICATION

The timing of applications to achieve control may vary quite a bit with different weed species. The following application times are suggested for certain problem weeds.

Spurweeds: The best control of spurweed can be obtained by applying SECURITY Purge 2 when spurweed has emerged (December and January).

Florida Betony: This weed emerges in the fall, so an application of SECURITY Purge 2 in mid to late October followed with a second application in mid to late February would give the best control.

Dichondra, Moneywort: The best control for these weeds can be obtained by applying SECURITY Purge 2 in early April followed by a second application in July. Do not apply to growing Bermuda grass.

Applications for spurweed or Florida betony generally will give control or suppression of the other weeds listed. However, as a general rule SECURITY Purge 2 will give the best control when applied to young tender weeds or just prior to weed emergence.

ST. AUGUSTINE AND CENTIPEDE GRASSES

SECURITY Purge 2 may be applied to established St. Augustine and centipede grasses during both the dormant and the growing season. Best results are usually obtained in the early spring or dormant periods when weeds are small or have not emerged. Follow rates below.

DORMANT BERMUDA GRASS

SECURITY Purge 2 may be applied to Bermuda grass during the dormant season only. Applications to Bermuda grass during the growing season will cause injury. Follow rates below.

RATE OF APPLICATION

Determine the total area to be sprayed and base rate of application on the chart below. Best results can be obtained by dividing total area into smaller areas for treatment. Measure area to be treated accurately and follow rates exactly as overtreatment can cause injury. Avoid overlapping spray pattern while treating. Shake spray mixture during use to prevent settling.

<u>AREA TO BE TREATED</u>	<u>AMOUNT OF PURGE 2</u>	<u>SUGGESTED AMOUNT OF WATER</u>
500 sq. ft.	3 tablespoons	1 gallon
1,000 sq. ft.	6 tablespoons (3 oz.)	2 gallons
3,000 sq. ft.	8 ounces	6 gallons
5,450 sq. ft.	1 pint	10 gallons
10,900 sq. ft. (1/4 acre)	2 pints	20 gallons

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Do not reuse empty container. Wrap empty container and put in trash collection.

NOTICE: The seller makes no implied warranties which extend beyond the description on this label.

NET CONTENTS: 1 PINT

WOOLFOLK CHEMICAL WORKS, INC.

MANUFACTURERS

FORT VALLEY, GEORGIA