


SUREGARD BRAND
CAPTAN 50-WP
AGRICULTURAL FUNGICIDE



ACTIVE INGREDIENTS:

Captan -----
Related Derivatives -----

INERT INGREDIENTS: -----

EPA Reg. No. 769-540

EPA Est. 769-GA-1

KEEP OUT OF REACH OF CHILDREN

DANGER - PELIGRO

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este
producto hasta que la etiqueta le haya sido explicada
ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Flush eyes immediately with fresh water for 15
minutes. Call a physician. IF SWALLOWED: Drink promptly a
large quantity of milk, egg whites, gelatin solutions, or if
these are not available, drink large quantities of water.
Avoid alcohol. IF ON SKIN: Wash with plenty of soap and
water. IF INHALED: Remove victim to fresh air. If not
breathing, give artificial respiration, preferably mouth-to-
mouth. Get medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER! Causes irreversible eye damage. Harmful if
swallowed or inhaled. May cause allergic skin reactions.
Do not get in eyes. Wear goggles or face shield when
handling. Avoid contact with skin and clothing. Avoid
inhalation of dust or spray mist. Remove and separately
launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to
water or wetlands (swamps, bogs, marshes and potholes).
Drift and runoff from treated areas may be hazardous to
aquatic organisms in neighboring areas. Do not contaminate
water by cleaning of equipment or disposal of wastes.

PHYSICAL OR CHEMICAL HAZARDS

Protect from excessive heat. Store in a cool, dry place.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a
manner inconsistent with its labeling. READ ENTIRE LABEL.
USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND
DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL
REGULATIONS.

REENTRY STATEMENT

BEST AVAILABLE COPY

Keep all unprotected persons out of operating areas or vicinity where there may be danger of drift. Do not allow persons to enter treated areas within 4 days following application unless a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet and chemically resistant gloves are worn. Conspicuously post reentry information at site of application. Certain states may require more restrictive reentry intervals; consult your State Department of Agriculture for further information. Written or oral warnings regarding use of protective clothing and accidental exposure must be given to workers who are expected to be in treated areas or in areas about to be treated.

USE PRECAUTIONS: Wear goggles or face shield when handling. Wash thoroughly with soap and water after handling. Remove contaminated clothing and separately launder clothing before reuse. Mixers/loaders when mixing and loading this product must wear a long sleeved shirt and long pants or a coverall that covers all parts of the body except the head, hands and feet. Wear chemical resistant gloves, goggles or a face shield and a chemically resistant apron. Fieldworkers and harvesters must wear chemical resistant gloves. Leather or fabric gloves are not acceptable. A dust mask must be worn when mixing and loading this product. During application only, if from a tractor with a completely enclosed cab, a long sleeved shirt and long legged pants must be worn. Chemically resistant gloves must be available in the cab and worn while exiting.

AFTER WORK: Take off all clothes and shoes. Shower using soap and water. Do not wear contaminated clothing. Wash protective clothing and equipment with soap and water after each use. Personal clothing worn during use must be laundered separately from household articles. Clothing and protective equipment heavily contaminated or drenched with captan must be destroyed according to state and local regulations.

NOTE: For concentrate sprayer application, the same quantity (per acre) of Captan 50-WP as would normally be applied when spraying dilute mixture should be applied to each acre in sufficient water for coverage. When more than one dosage rate is given, use the higher rate when conditions are favorable for disease development and use the lower rate when disease conditions are less favorable.

Do not apply this product through any type of irrigation system.

COMPATIBILITY: Captan 50-WP may be combined with commonly used insecticides, adjuvants and fungicides with the exception of strongly alkaline materials such as hydrated lime which reduces the fungicidal activity. Do not use

BEST AVAILABLE COPY

this product in combination with oil sprays or with solvent formulations of parathion products. Do not use this product on foliage closely following oil sprays.

APPLES: Primary Scab Infection - Apply 8 lbs. Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment. Applications should be made at 5-7 day intervals as needed to maintain control in prebloom, bloom, petal-fall and first cover spray. Secondary Apple Scab, Brooks Fruit Spot, Flyspecking, Bitter Rot, Black Rot, Botryosphaeria, Black Pox - Apply 4 to 8 lbs. Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment. Apply at 10 to 14 day intervals in second and later cover sprays up to start of harvest. CAUTION: Captan 50-WP should not be used in combination with or closely following or in alternation with wettable sulfur products on sulfur-sensitive varieties of apples such as Red Delicious, Staygreen, Baldwin, King, etc. as severe injury and defoliation may occur. Foliage injury may at times occur to Red Delicious, Winesap and other sensitive varieties of apples in early season sprays.

CANTALOUPE, CUCUMBERS: Anthracnose - Apply 4 lbs. Captan 50-WP per acre in sufficient water for thorough coverage. Begin applications at first signs of infection and repeat at 5 to 7 day intervals or as needed to maintain control.

CHERRIES: Cherry Leaf Spot, Brown Rot, Botrytis Rot - 4 lbs. per acre in 20 to 200 gallons of water using ground equipment. Spray at prebloom, bloom, petal-fall, shuck, cover and preharvest sprays. Application at 3 to 4 day intervals may be necessary during bloom to control Blossom Blight. Repeat applications at 7 to 10 day intervals as needed to maintain control up to start of harvest.

GRAPE: Downy Mildew, Black Rot - 4 lbs. per acre in 10 to 200 gallons of water using ground equipment. Make first application just before bloom and repeat immediately after bloom and at 10 to 14 day intervals for 1 to 3 additional applications depending on disease pressure and susceptibility of the grape variety. Dead Arm Disease (California) - 2 lbs. per 100 gallons spray, 3 to 6 lbs. per acre in 10 to 200 gallons of water using ground equipment. Apply first spray when green tissue begins to show but... before shoots are 1 inch long and repeat application when shoots are 6 to 6 inches long. (Northeast) - 2 to 4 lbs. per acre in 10 to 200 gallons water using ground equipment. Make first application when shoots are 1/2 to 1 inch long and repeat application when shoots are 4 to 6 inches long.

NECTARINES: Brown Rot, Scab - Apply 5 lbs. Captan 50-WP per acre in 20 to 250 gallons of water using ground equipment. Apply in full pink, bloom, petal-fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control Blossom Blight. Repeat

4/6
applications at 7 to 14 day intervals as necessary to maintain control. Continue applications through harvest if conditions favor Brown Rot. Coryneum Blight (Shot-Hole) - Apply 5 lbs. Captan 50-WP per acre in 20 to 250 gallons water using ground equipment. Apply in pink bud, full bloom, petal-fall and cover sprays as necessary and as a post harvest spray (but before leaves drop).

PEACHES: Brown Rot, Scab - 8 lbs. per acre in 20 to 400 gallons water using ground equipment. Apply at full pink, bloom, petal-fall, shuck, cover and preharvest sprays. Applications at 3 to 4 day intervals may be necessary during bloom to control Blossom Blight. Repeat applications at 7 to 14 day intervals as needed to maintain control. Continue applications through harvest if conditions favor Brown Rot. Coryneum Blight (Shot-Hole) - Apply 8 lbs. Captan 50-WP per acre in 20 to 400 gallons of water using ground equipment. Apply at pink bud, full bloom and petal fall stages and in cover sprays as necessary and as a post harvest spray (but before leaf drop).

PEARS: Pear Scab (except Pacific Northwest) - Primary Infection - 8 lbs. per acre in 20 to 200 gallons water using ground equipment. Apply in pre-blossom, petal-fall and cover sprays. Repeat cover spray applications at 7 to 14 day intervals under severe scab infection periods. Russetting may be produced on Bose pears. Do not use on D'Anjou pears.

PRUNES, PLUMS: Brown Rot - 6 lbs. per acre in 20 to 300 gallons water using ground equipment. Apply at green bud, popcorn, bloom and petal-fall stages. Repeat in cover sprays as conditions warrant. Prune Russet Scab (Lacy Scab in California) - 8 lbs. per acre in 20 to 300 gallons water using ground equipment. Apply at full bloom.

STRAWBERRIES: Botrytis Rot - 6 lbs. per acre in sufficient water for thorough coverage. Begin applications when new growth starts in the spring and before fruit starts to form. Repeat applications weekly. Under conditions favorable to fruit rot, continue applications through harvest period treating immediately after each picking.

TOMATOES: Gray Leaf Spot, Anthracnose, Septoria Leaf Spot - 6 lbs. per acre in sufficient water for thorough coverage. Severe Infection - 6 to 8 lbs. per acre in sufficient water for thorough coverage. Apply at first sign of disease. Repeat at 5 to 7 day intervals or as necessary to keep disease under control.

ORNAMENTAL USES

BEST AVAILABLE COPY

5/10
AZALEA, CARNATION AND CHRYSANTHEMUM CUTTINGS for Damping-Off: Dip cutting in a suspension of 4 lbs. per 100 gallons spray before bedding.

ORNAMENTALS: Roses - Black Spot; Carnations - Rust and Leaf Spot; Chrysanthemums - Flower Blight: 2 lbs. per 100 gallons of spray. Apply at first sign of disease; cover upper and lower leaf surfaces thoroughly. Repeat every 7 to 10 days.

TURF, LAWN: Brown Patch, Leaf Spot, Damping-Off, Root Rot - 2 lbs. per 100 gallons spray. Apply 10 gallons to each 1000 square feet or apply 8 lbs. per acre in water for coverage. Apply prior to infection. Repeat at 7 to 10 day intervals as necessary. Do not graze treated areas or feed clippings to livestock.

SOIL AND GREENHOUSE BENCH TREATMENT: Preplanting for damping-off and root rot diseases on seedlings or transplants on roses (and other shrubs and trees), many flowers and lawn seedbeds - Apply as a dust at 4 to 5 oz. per 1000 square feet or spray at 2 lbs. to 100 gallons spray using 15 gallons to each 1000 square feet. Cultivate into up to 3 to 4 inches of soil before planting. USE PRECAUTIONS: Only the applicator is permitted to be in the greenhouse during application of Captan. Open vents to greenhouse during application and for at least 1 hour following application. Workers planting in Captan-treated soil must wear chemically resistant gloves.

Do not apply under conditions involving possible drift to food, forage or other plantings that might be damaged or the crops thereof rendered unfit for sale, use or consumption.

STORAGE AND DISPOSAL

STORAGE: Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Pesticide wastes are acutely hazardous.

Improper disposal of excess pesticide, spray mixture or rinsate is a violation of federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Representative at the nearest EPA Regional Office for guidance. CONTAINER

DISPOSAL: Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

BEST AVAILABLE COPY

6/6
NOTICE: The seller makes no implied warranty of
merchantability nor any other warranties which extend beyond
the description on this label.

NET WEIGHT: 4 LBS.

MANUFACTURED FOR
SURECO, INC.
FORT VALLEY, GA 31030

