



COMPANY
NUMBER 769-

PRODUCT
SERIAL NO. 541

LABEL
APPROVAL
DATE 09/18/86

petal fall, when first leaves unfold and every 21 days for 3 additional sprays. Russeting may be produced on Rose pears. Do not use on O Anjou pears. **RASPBERRIES** (spur blight, anthracnose). Apply when canes are 8 to 10 in. long and 2 weeks later. Fruit rot. Apply 3 to 5 days before harvest, mid harvest 8 to 10 days later. **STRAWBERRIES** (botrytis rot). Apply when new growth starts in spring. Repeat applications weekly to harvest. For severe infection, continue through harvest period, treating immediately after each picking. **SEED TREATMENT** (seed rot, damping off) - Beans, cabbage, corn, melons, peas, squash - Use 1/2 teaspoon per pound of dry seed. Beets, chard, spinach, grass - use 1 teaspoon per pound of dry seed. Mix thoroughly in paper bag or glass jar. Do not use treated seed for feed or food or oil purposes. **SOIL AND GREENHOUSE BENCH TREATMENT:** Preplanting for damping off and root rot diseases on seedling or transplants of roses and other shrubs and trees, many flowers and vegetables such as beans, peas and cucurbits and lawn seedbeds - Apply as a dust at 4 to 5 oz. per 1000 square feet or spray 4 teaspoons per gallon of water using 15 gallons to each 1000 square feet. Cultivate into upper 3 to 4 inches of soil before planting. Only the applicator is permitted to be in the greenhouse during application of Captan to the soil. Open vents to greenhouse during application and for at least 1 hour after application. Workers planting in Captan treated soil in greenhouses must wear gloves impermeable to Captan.

ORNAMENTALS - AZALEAS, CARNATIONS, CHRYSANTHEMUMS (damping off). Dip cuttings in a suspension of 2 1/2 tablespoons per gallons of water before bedding. **CARNATIONS** (rust, alternaria leaf spot), **CHRYSANTHEMUMS** (botrytis flower blight). Apply 4 teaspoons per gallon of water at first sign of disease and repeat at 7 to 10 day intervals. Use the shorter interval under conditions favorable to infection. **GLADIOLUS**, **TUBEROUS BEGONIAS** (rot, damping off). Mix 4 teaspoons per gallon of water and dip bulbs for 30 minutes. Stir, drain and plant. **ROSES** (black spot). Apply 4 teaspoons per gallon of water at first sign of disease and repeat at 7 to 10 day intervals. Use the shorter interval under conditions favorable to infection.

LAWNS, TURF: To control brown patch, damping off, leaf spot and root rot, mix 4 teaspoons per gallon of water and apply to 100 square feet of turf. Repeat at 7 to 10 day intervals as needed. For melting out, use 5 to 8 1/2 teaspoons per gallon and apply to 100 square feet beginning when grass resumes growth in the spring (March - April). Repeat at 7 to 10 day intervals as long as cool, wet weather persists. Do not graze treated areas or feed clippings to livestock.

VEGETABLES: Unless otherwise specified the following vegetable diseases can be controlled using a standard rate of 4 teaspoons per gallon of water. Spray for thorough coverage. **CARROTS** (septoria leaf spot, cercospora blight). Apply at first sign of disease and repeat at 6 to 10 day intervals. **CUCURBITS** (cantaloupes, cucumbers, summer and winter squash, pumpkins, melons) (anthracnose, downy mildew). Apply at 5 to 7 day intervals beginning at first sign of disease and repeat at 7 to 10 day intervals as necessary to maintain control. **LETTUCE, SPINACH** (downy mildew). Apply at first sign of disease and repeat at 7 to 10 days intervals as necessary to maintain control. **POTATOES** (early and late blight). Use 8 teaspoons per gallon of water. **TOMATOES** (early and late blight, gray leaf spot, septoria leaf spot). Use 4 teaspoons per gallon of water. For severe infection on tomatoes, use 8 teaspoons per gallon of water. Begin when disease first appears and repeat at 5 to 7 day intervals as necessary to keep diseases under control.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal of this product. Store this product in original container and keep in a locked storage area. Do not reuse container. Securely wrap original container in several layers of newspaper and discard in trash.

NOTICE: Buyer assumes all responsibility for safety and use not in accordance with directions.

SECURITY

Captan

Garden Spray

For Plant Disease Control

ACTIVE INGREDIENTS:	BY WT.
*Captan	47.3%
Related Derivatives	2.7%
INERT INGREDIENTS:	50.0%
	TOTAL 100.0%

*N-Trichloromethylthio-4-cyclohexeno-1,2-dicarboximide.

EPA REG. NO. 769-541

EPA EST. 769-GA-1

KEEP OUT OF REACH OF CHILDREN
DANGER!

(See Additional Precautionary Statements
on Side Panel)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Give the victim water or milk to drink and telephone for medical advice. DO NOT make the victim vomit unless directed to do so by medical personnel. If medical advice cannot be obtained, then take the victim to the nearest medical emergency treatment center or hospital.

IF INHALED: Remove victim to fresh air and administer artificial respiration if indicated.

IF ON SKIN: Wash area immediately with soap and water. Seek medical attention if irritation persists.

IF IN EYES: Flush eyes for 15 minutes with water and seek medical attention.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER: Causes irreversible eye damage. Harmful if swallowed or inhaled. May cause allergic skin reactions. Do not get in eyes. Wear goggles or face shield when handling. Avoid contact with skin and clothing. Remove and separately launder clothing before reuse.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water or wetlands (swamps, bogs, marshes, and potholes). Drift and runoff from treated areas may be hazardous to aquatic

organisms in neighboring areas. Do not dispose of wastes. In the event that you have a spill, completely cover them with soil.

ENDANGER

The use of this product in a manner that modifies their habitat is prohibited by the general location of their habitats. Alabama (Lauderdale, Limestone and Blackwater River); Alabama (Lauderdale and Wayne); Pecos gambusia; New and Pecos); Freshwater Mussels; Texas (Virginia (Smyth, Scott, Lee, W. Pulpish (Texas (Reeves and Jeff Davis). Prior to the use of this product in the area not located in, or immediately adjacent to your regional U.S. Fish and Wildlife outdoor use of Captan is prohibited.

PHYSICA

Protect from excessive heat. Store in

Do not

It is a violation of Federal law to use this product in a manner that modifies their habitat. **PROTECTIVE CLOTHING:** Wear non-permeable to Captan when using pants and long-sleeved shirt and apply thoroughly with soap and water before must be laundered separately from treated crop with crops other than the enter treated areas within 4 days for. Conspicuously post reentry information. microfine wettable powder specially diseases of fruits, vegetables and ornamentals for the control of certain seed rots and

DIRTY

Captan Garden Spray is a wettable powder suspension for spray application. completely dispersed. This product contains insecticides, adjuvants and fungicides. **FRUITS:** To control the diseases listed on the side panel, apply Captan Garden Spray per gallon of water.

APPLES (scab, fruit spot, sooty blotch, rot). Apply at weekly intervals beginning 10 to 14 day intervals thereafter until harvest. **PEACHES** (brown rot, botrytis rot). Apply at pre-bloom, bloom, and post-bloom stages. **BLACKBERRIES** (black rot). Apply first application just before bloom. Make 1 to 3 additional applications. **DEAD ARM** (current season), apply when shoots are 4 to 6 inches long. **SCAB**, Apply in pink bud, full bloom, and every 12 to 14 days until harvest.

ACCEPTED

SEP 18 1986

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 769-541