

pm04

769-612

7/9/98

llk



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

Stacey A. Schmidt
SureCo
7501 Harps Mill Road
Raleigh, N.C. 27615-5422

OFFICE OF
PREVENTION, PESTICIDES AND
TOXIC SUBSTANCES

JUL 9 1998

Subject: SureCo Garden & Kennel Dust 10%
EPA Registration No. 769-612
Amendment dated June 9, 1998
Corrections to Reinstatement of Ornamentals Section dated
May 15, 1998.

Dear Ms. Schmidt:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

1. On page 3 under the heading PREHARVEST USE LIMITATION place a comma between the words squash and cantaloupe.
2. On page 3 under the same heading as above change the sentence "Allow 14 days between harvest and harvest on leafy vegetables.....and spinach" to "Allow 14 days between application and harvest on leafy vegetables.....spinach".

Submit two copies of your final printed labeling before you release the product for shipment. A stamped copy of the labeling is enclosed for your records.

Sincerely,

George Tompkins, Ph.D., Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505C)



Recycled/Recyclable
Printed with Soy/Canola Ink on paper that
contains at least 50% recycled fiber

218 ACCEPTED
with COMMENTS
In EPA Letter Dated:

JUL 9 1998
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
769-612

SureCo GARDEN & KENNEL DUST 10%

CONTROLS INSECTS ON
VEGETABLES - PETS - LAWNS - AND ORNAMENTALS

ACTIVE INGREDIENT:	
Carbaryl (1-naphthyl N-methylcarbamate)	10.0%
INERT INGREDIENTS:	90.0%
Total:	100.0%

EPA Reg. No. 769-612

EPA Est. No. 769-GA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

See Back Panel for Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.


IF INHALED: Remove victim to fresh air.

IF IN EYES: Flush with water.

NOTE TO PHYSICIAN: Carbaryl is a moderate, reversible, cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase-inhibiting drugs.

Manufactured by:
SureCo, Inc.
Fort Valley, GA 31030

Net Contents: _____



PRECAUTIONARY STATEMENTS
Hazards to Humans & Domestic Animals
CAUTION

May be harmful if swallowed. Avoid breathing dust. Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water by cleaning of equipment or disposal of waste. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

4/18

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls, Waterproof gloves, and Shoes plus socks.

CAUTION: Some phytotoxicity may occur on tender foliage in the presence of rain or high humidity of several days' duration following application.

PREHARVEST USE LIMITATION: Allow 2 days between application and harvest for sweet corn. Allow 3 days between application and harvest for fresh beans, broccoli, Brussels sprouts, cabbage, cauliflower, cowpeas, kohlrabi, cucumbers, squash cantaloupe, pumpkins, melons, tomato, eggplant, pepper, and okra. Allow 7 days between application and harvest on root and tuber crops including radishes, garden beets, rutabagas, turnips, carrots, and potato. Allow 14 days between harvest and harvest on leafy vegetables including collards, garden beet tops, rutabagas, turnip tops, kale, lettuce (head and leaf), mustard greens, and spinach. Allow 21 days between application and harvest for dried beans or peas. Allow 7 days between application and harvest on small fruits and berries including blackberries, boysenberries, dewberries, loganberries, raspberries, blueberries, grapes, and strawberries. Allow 7 days between application to poultry and game birds or premise treatment and slaughter of birds.

APPLICATION: Begin applications when insect or damage is first observed. Do not repeat application more than once a week.

VEGETABLES

BEANS: To control Mexican bean beetle, bean leaf beetle, cucumber beetles, leafhoppers, flea beetles, Japanese beetles and corn earworm, apply at the rate of 10 pounds per acre or one ounce per 100 feet of row. Where Mexican bean beetle is the only problem, a rate of 10 pounds per acre is sufficient. To control lygus bug, stink bug and Lima bean pod borer, apply 20 pounds per acre.

CABBAGE, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, KOHLRABI: To control flea beetles and harlequin bugs, apply at the rate of 10 pounds per acre. To control corn earworm, armyworm, imported cabbageworm, diamond-back moth larvae, apply 10 to 20 pounds per acre or 1 to 2 ounces to 100 feet of row. Will not control cabbage loopers.

COLLARDS, GARDEN BEETS, KALE, LETTUCE, MUSTARD GREENS, RADISHES, RUTABAGAS, SPINACH, TURNIPS: To control flea beetles, harlequin bug and leafhoppers, apply at the rate of 10 pounds per acre. To control armyworms, imported cabbageworm, diamond-back moth larvae, corn earworm, tarnished plant bug and stink bugs, apply 10 to 20 pounds per acre or 1 to 2 ounces to 100 feet of row. For best control of corn earworm after lettuce heads begin to form, use 20 pounds per acre in a 7 day schedule. Will not control cabbage looper.


CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control pickleworm, melonworms, cucumber beetles, squash bug, flea beetles, apply at the rate of 10 pounds per acre or 1 ounce to 100 feet or row.

CORN: To control corn earworm, European corn borer, fall armyworm, sap beetles, and Japanese beetles, apply at the rate of 15 to 20 pounds per acre or 2 ounces to 100 feet or row. For budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply 7 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending upon severity of infestation.

CARROTS: To control flea beetles, leafhoppers, spittlebugs, lygus and stink bugs, apply at the rate of 10 to 20 pounds per acre or 1 to 2 ounces to 100 feet of row.

OKRA: To control corn earworm and stink bugs, apply at the rate of 10 to 20 pounds per acre or 1 to 2 ounces to 100 feet of row.

TOMATO, EGGPLANT, PEPPER: To control tomato fruitworm, fall armyworm, flea beetles and leafhoppers, apply at the rate of 10 pounds per acre or 1 ounce to 100 feet of row. For control of tomato hornworm, European corn borer and stink bugs, apply at the rate of 10 to 20 pounds per acre.



cel

POTATO: To control Colorado potato beetle, apply at the rate of 10 pounds per acre. For control of leafhoppers and flea beetles, apply at 10 pounds and for lygus bugs, apply at 20 pounds per acre.

SMALL FRUIT

BLACKBERRIES, BOYSENBERRIES, DEWBERRIES, LOGANBERRIES, RASPBERRIES: To control Japanese beetles, leafrollers, leafhoppers and raspberry aphid, apply at the rate of 20 pounds per acre.

BLUEBERRIES: To control blueberry maggot, cherry and cranberry fruitworms and Japanese beetles, apply at the rate of 15 to 20 pounds per acre. Apply 3 weeks before harvest and repeat 10 days later if necessary.

GRAPES: To control grape leaf folder, grape leafhopper and grape leaf skeletonizer, apply at the rate of 10 to 20 pounds per acre. Apply just before first brood leaf folder larvae emerge from rolls and as needed for leafhoppers. To control grape berry moth, Japanese beetle, June beetle and red-banded leafroller, apply at the rate of 20 pounds per acre.

STRAWBERRIES: To control meadow spittlebug, strawberry leafroller and strawberry weevil apply at the rate of 10 to 20 pounds per acre.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses. Do not allow children or pets to contact treated area until dust has settled.

RESIDENTIAL USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use,
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

DIRECTIONS FOR RESIDENTIAL USE

Wearing household latex or rubber gloves, use a dust applicator such as a shaker, can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow

dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week. Apply when insects or their damage appear. Foliage should be dry before dust is applied. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

ORNAMENTALS

To control bagworm, blister beetles, boxelder bug, boxwood leafminer, flea beetles, Japanese beetle, lacebugs, leafhoppers, leaf rollers, periodical cicada, plant bugs, psyllids, rose slug, rose aphid, scale insects in the crawler stage, tent caterpillars, and exposed thrips, apply thoroughly to the infested plants at first sign of damage and repeat at 7 day intervals or as necessary for good insect control. To obtain scale control apply in spring and early summer when scale crawlers are present.

LAWNS

To control fleas, ticks, leafhoppers, fall armyworm, and sod webworm, apply at the rate of 2 pounds per 1000 square feet. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

EARWIGS

This material is effective in controlling earwigs and when properly used will reduce the number gaining entrance to homes and buildings. Apply heavily at the rate of 1 pound to 150 linear feet completely around the foundation wall of buildings. The material should be applied heavily in a band 3 to 4 inches wide using a shaker-type can. Earwigs breed over wide areas around yards and fields and are driven by cold weather to seek shelter in warm buildings. As they move along the foundation wall seeking entrance, they come in contact with the treated barrier and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation wall. Treat also around refuse piles, lumber, mulch and other areas in the yard offering areas of concealment to earwigs. Repeat application in 10 days or when earwigs become troublesome. If the dust barrier is washed away by rain, it should be replaced with a new application if earwigs are still a problem.

POULTRY INSECT CONTROL

CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS AND PIGEONS:

LITTER TREATMENT - Apply 1/2 pound per 40 square feet to control Northern fowl mite, chicken mite, lice, fleas and bed bugs. Use a scoop and scatter by hand or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

HAND APPLICATIONS TO BIRDS: Use 1/2 pound to treat 100 birds by means of a shaker can, squeeze bottle or hand duster to control Northern fowl mite, chicken mite, lice and fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and fleas is a supplement to litter treatment for control of these pests. For the control of sticktight fleas, it is especially important to treat necks and head.

8/18

DUST-BATH BOXES: Use 2 1/2 pounds per box for each 100 birds to control Northern fowl mite, chicken mite, lice and fleas. This is a supplement to litter treatment for chicken mite and flea-control. On ranges, two natural wallows may be substituted for each dust-bath box.

PET INSECT CONTROL

Use this product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Dust bedding and doghouse using a shaker can or other dust applicator.

DOGS AND CATS: For ticks and fleas, dust liberally over the entire animal, avoiding the eyes. Rub thoroughly into the skin. Comb out dead ticks a few hours after treatment. Repeat at weekly intervals if needed. Do not use on pregnant dogs.

KENNELS

DOG AND CAT QUARTERS: For ticks and fleas, dust liberally in and around sleeping quarters, kennels and dog houses by means of a shaker can, hand or powder dust. Force dust into cracks and crevices. Repeat as necessary.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Store pesticide in original container and place in a locked storage area.

Disposal: Do not reuse container. Wrap and discard in the trash. Wastes resulting from this product may be disposed of on site or at an approved waste disposal facility.

This container is sold by weight, NOT by volume. You can be assured of the proper weight even though some settling of contents normally occurs during transportation and handling.

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product.
DISCLAIMER: SureCo warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. SureCo makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use, handling or application of this product.

per EPA letter 5/15/98