

JAN - 4 1994

SureCo, Inc.
c/o Mr. H. R. McLane
H. R. McLane, Incorporated
7210 Red Road, Suite 206
Miami, Florida 33143-5321

Subject: Revised Labeling--Watering-in Requirement
Dursban 2E Insecticide
EPA Registration Number 769-641
Your Application Dated December 9, 1993

Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable provided that you make the labeling changes indicated below before you release the product for shipment bearing the amended labeling:

1. Add the following sentence to the Hazards to Humans and Domestic Animals section: "Keep children and pets off treated areas until dry."

2. Revise the Environmental Hazards section to:

This pesticide is toxic to birds, fish, and aquatic invertebrates. Birds feeding in treated areas may be killed. Clean up spilled product to reduce exposure to wildlife. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply or allow to drift to blooming crops or weeds if bees are visiting the treatment area.

294

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

DURSBAN® 2E INSECTICIDE

CONTROLS NUMEROUS PESTS, AS LISTED, OF HOUSEHOLDS, TURF AND ORNAMENTAL PLANTS ALSO RECOMMENDED FOR MOSQUITO CONTROL AND AREA CONTROL OF TICKS AND CHIGGERS. USE AS A MOUND TREATMENT TO KILL IMPORTED FIRE ANTS SEE SUPPLEMENTAL LABELING FOR ADDITIONAL USES

To be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts and Other Trained Personnel Responsible for Insect Control Programs

ACTIVE INGREDIENT:
Chlorpyrifos (0,0-diethyl 0-(3,5-
trichloro-2-pyridyl)phosphorothioate) 22.5%

INERT INGREDIENTS 77.5%

Total 100.0%

Contains 2 pounds Chlorpyrifos per gallon
DURSBAN Reg. Trademark, Dow/EI

KEEP OUT OF REACH OF CHILDREN WARNING

STATEMENTS OF PRACTICAL TREATMENT

IF SWALLOWED Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention. **IF INHALED** Remove victim to fresh air. Get medical attention. **IF IN EYES** Flush eyes with plenty of water. Get medical attention if irritation persists. **IF ON SKIN** In case of contact, remove contaminated clothing and wash affected area with soap and water. **NOTE TO DURSBAN** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine only by reaction to the anticholinergic syndrome.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING May be fatal if swallowed. Excessive absorption through skin may be fatal. Do not get in eyes, skin or clothing. Avoid breathing of vapors and spray mist. Wear eye protector. Wear protective clothing and chemically resistant gloves when handling. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Keep away from food, feedstuffs and water supplies.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic areas. Cover or incorporate soils. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply or allow to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use, pour, spit or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: Avoid storage at high temperatures. Protect from moisture. Avoid contamination with water, acids or alkalis. Do not store near heat or open flame. Keep container closed. Store in original container only. Separate pesticides to avoid cross-contamination of other pesticides, fertilizer, food and feed. Keep container in locked storage areas. If spilled, absorb with dry material such as sand or sawdust and sweep up for disposal. **Disposal:** Do not contaminate water, food or feed by storage and disposal. Pesticide wastes are toxic. Proper disposal of excess pesticide, spray mixture, or waste is a violation of Federal law. If these wastes, or no, be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste Response Team at the nearest EPA Regional Office for guidance. **Container:** Large Al. Metal: Treat here for equipment. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and Local authorities. **Plastic:** Treat here for equipment. Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or if allowed by State and Local authorities, by burning. If burned, stay out of smoke. (709-641 P&M12-09-83)

EPA Reg. No. 709-641 NET CONTENTS 1 GALLON SURECO, INC. Fort Valley, GA 31030 EPA Est. No. 709-GA-1

PESTS INDOORS

General Information: Use DURSBAN 2E insecticide to control the pests indicated in the areas listed below by applications as a 0.25% or 0.5% Chlorpyrifos spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute DURSBAN 2E with water or with a suitable base as shown in DILUTION RATES paragraph. Repeat treatment as necessary. Use only decontaminated equipment for oil based sprays to be applied in food handling establishments.

Do not mix the spray into the air or allow the spray to contact food or food containers. Areas treated may include dark corners of rooms and closets, along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass CRACK & CREVICE treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or which occur at points between different elements of construction or between equipment and floors. Equipment capable of delivering a stream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back), Durban 2E may be applied after 2 days from the last treatment but emergency use should be limited to once per month.

APPLICATIONS OF THIS PRODUCT IN FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A SPOT AND/OR CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

APPLICATION WITHIN RESIDENTIAL BUILDINGS

including Homes and Apartment Buildings and Within Non-Food Areas of Industrial, Institutional and Commercial Buildings including Hospitals, Stores, Manufacturing Plants and Warehouses. For Spot Treatment Only. Apply as a coarse spray or with a paint brush to localized areas where **Ants, Cockroaches, Clover mites, Confused Flour Beetle, Crickets, Earwig, Firebrats, Indian Meal Moth, Mediterranean Flour Weevil, Rice & Grain Weevils, Millipedes, Red Flour Beetle, Saw Toothed Grain Beetle, Scorpion, and Spiders** are found or normally occur including dark corners of rooms and closets, cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets, around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For **Clothes moths** use 0.5% spray only. Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches.

For the Control of **Carpenter Beets**, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting under carpeting, rugs, furniture in closets and on shelving and wherever else these insects are seen or suspected.

DILUTION RATES

To make a 0.25% spray solution:
FOR WATER SPRAY - mix 1 1/3 oz. in 1 gallon of WATER
FOR AN OIL SPRAY - mix 1 fl. oz. in 1 gallon of DEO-BASE OIL.
To make a 0.5% spray solution:
FOR WATER SPRAY - mix 2 2/3 oz. in 1 gallon of WATER
FOR AN OIL SPRAY - mix 2 fl. oz. in 1 gallon of DEO-BASE OIL.
For larger amounts multiply the base ounce (amount) in 1 gallon by number of gallons required for the "job". (Decorated bees, **Kanawha** **Blue**, **White** **Chigger** and/or **Flashing** **Chigger** is covered elsewhere. (DOVP) or pyrethrins may be added to the spray mixture containing Durban 2E at 1:1000. For best results use dichlorvos at a concentration of 0.25% in both water and oil-based sprays. Consult labeling for the scheduler or pyrethrin product used to determine the amount to use. Do not use a spray concentration. Also read and observe all use and handling precautions for these products. **DO NOT** use Durban 2E in combination with dichlorvos (DDVP) or pyrethrins for treatment of food areas of food handling establishments.

APPLICATION WITHIN FOOD HANDLING ESTABLISHMENTS

PLACES OTHER THAN PRIVATE RESIDENCES IN WHICH FOOD IS HELD, PROCESSED, PREPARED OR SERVED INCLUDING BUT NOT LIMITED TO RESTAURANTS, GROCERY STORES, BAKERIES, BOTTLING PLANTS, CANNERIES AND GRAIN MILLS.

a Durban 2E according to the DILUTION RATES. Use as a **TREATMENT** to control **Ants, Cockroaches, Clover mites, Confused Flour Beetle, Crickets, Earwig, Firebrats, Indian Meal Moth, Mediterranean Flour Weevil, Rice & Grain Weevils, Millipedes, Red Flour Beetle, Saw Toothed Grain Beetle, Scorpion, and Spiders** in and around both food and non-food areas by application as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen, suspected or having or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff. **Brown Dog Ticks:** Thoroughly apply 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices, along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product.

NOTE: (Pests Indoors) Use of **base spray only** for cockroaches, water housing, junction or switch boxes. Prior to application de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switch in the "ON" position. **DO NOT** use water-based sprays in these places. Do not use in poultry houses and other animal buildings. Do not use in serving areas where food is exposed. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

PESTS OUT-OF-DOORS

General Information: Use Durban 2E for the control of certain pests of lawns and ornamental plants. For area control of ticks and chiggers and for mosquito control. Consult the following directions for dosage and application details to control the pests listed.

PESTS OF LAWNS AND OTHER ORNAMENTAL AND RECREATIONAL TURF GRASS AREAS

Use Durban 2E to control the pests listed on the following table by application at the recommended dosage and in accordance with the directions given below. Dilute Durban 2E in enough water to obtain complete and uniform coverage of pest infested areas and apply as a coarse, low pressure spray using suitable application equipment. For soil insects thoroughly water after treatment to wash the insecticide into the soil, except as noted. Washing-in of the insecticide is not required for control of surface insects. For best results the lawn should be mowed at the time of treatment. Spray when pests first appear. Repeat as needed.

SURFACE INSECTS

Use 1 1/2 fl. oz. or 11,000 sq. ft. to control **Ants, Armyworms, Brown Dog Ticks, Centipedes, Chiggers, Chinchbugs, Clover Mites, Crickets, C. weevils, Earwig, Flea, Fire Ant (mound), Fleas, Grubs, Grasshoppers, Ground Squirrels, Jay Weevils, Leafhoppers, Lucerne Moths, Millipedes, Mosquitoes, Nymphs, Spiders, Spittlebugs, Sod Webworms, Spotted Spiders, Spotted and Ticks**. Spray when pests first appear; repeat as needed. For **Sod Webworms**, delay watering or mowing the treated area for 12 to 24 hours after treatment.

SOIL INSECTS

For **Lebanese Hardwood Weevils** use 3 fl. oz. or 11,000 sq. ft. Spray suspected problem areas in mid April and again in mid May or as recommended by your local Ag. Extension Service Specialist for your area. For **European Crane Fly Larvae** use 2 fl. oz. or 11,000 sq. ft.

For **White Grubs, Black Anthrenus weevils, European chigger, Southern meadow chigger & meadow chigger (larvae)** use 3 to 6 fl. oz. or 11,000 sq. ft. Spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by local Ag. Extension personnel. Immediately after spraying irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil. For **Lebanese Hardwood** use 1 1/2 to 6 fl. oz. or 11,000 sq. ft. Spray early in season when adults first appear. Repeat as needed.

PREMISE TREATMENT FOR CRAWLING INSECTS

To help prevent infestation of buildings by **ants, chiggers, scorpions, millipedes and spiders (wolfspiders)**, treat a band of soil 6 to 10 feet wide around and adjacent to the building into the building foundation to a height of 2 to 3 feet, where pests are active and may find entrance. Use 10 fluid ounces of Durban 2E per 100 gallons of water and apply as a coarse spray to thoroughly and uniformly wet the band area.

AREA CONTROL OF TICKS AND CHIGGERS

Use Durban 2E to control ticks and chiggers infesting non-crop land areas such as roadsides, forests and trails, picnic and camping sites, parts recreational areas and ground coverings where these pests are present.

Create a nuisance or area (equipment to 1 fluid ounce) backpack sprayer or of undershoot, grassy areas, etc. to allow use of treated areas when weather conditions allow, streams, ponds or they be killed if these water.

PEST: Use Durban 2E to treat mosquitoes found to be a nuisance according to the directions in a manner to apply a wetting spray to the areas. Attempt to prevent mosquito nuisance. Treat with ATTENTION. Keep children of lawn and garden fish, carnations, poinsettias, etc.

Use 0.5 fl. oz. (1 fl. oz.) in 3 fl. oz. water. Use 2.0 fl. oz. (4 fl. oz.) in 3 fl. oz. water. Use 1 fl. oz. (2 fl. oz.) in 3 fl. oz. water.

For larger amounts multiply required for the "job". Mow ornamental plant pests depending on measure.

INDOOR: Use 2 fl. oz. in 4 gal. water. 1 gallon of the mixture or surrounding area of 3 x 4 ft. areas. F. early or late in sprays may disturb area.

Apply Durban 2E to mow and areas shown in the table as necessary or No. 2 band of area treated. Apply the and power ground sprayer. dosage will also give faster or no vegetative cover. Un there are no fish or ornamentals of feed chigger. NOTICE: Read complete Fish & Game Department.

CONDITIONS: For areas of no vegetation 1.6 fl. oz. of Durban 2E of Durban 2E will treat 1/2 acre. For areas of heavy vegetation to 0.05 lb. A.I. Chlorpyrifos area.

CONDITIONS: For areas of no vegetation 1.6 fl. oz. Durban 2E per 180 Acres of area. For areas of heavy vegetation to 0.05 lb. A.I. Chlorpyrifos area.

See SUPPLIES OF Chloranil and M. (DORMAN)

BEST AVAILABLE COPY

USE 1 TO 2 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF BEETLES
 Beetles (such as, Cottonwood Leaf, Elm Leaf, Willow leaf, (1) - For Cottonwood Leaf Beetles use to control and adults infesting cottonwoods. Apply when insects are present or when foliage damage is ant.

USE 2 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS
 Borers (Clearwing moth (such as, Ash, Dogwood, 1 cedar Peachtree, Lusc, Oak Rhododendron) Metallic wood (such as, Bronze Birch, Flatheaded Appletree Twined Chestnut, Longhorned Beetles (such as Locust, Red Oak) Leaf miners, Needleminers (such as Jeffrey Pine, Lodgepole Pine, Spruce) Scale Crawlers such as (Cottonycushion, Cottony Maple, Eucalyptus scale, Fleckers scale, Florida Wax scale, Golden Oak Hemispherical, Lecanium Magnolia, Oak Kermes, Oystershell, Pine Needle, San Jose Tea White Birch)
 (1) - For borers apply to trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your Ag. Ext. Service for proper time to spray. Apply uniformly as a course spray.
 (2) For Cranberry Gardier larvae infesting Douglas Fir seedlings apply 1 quart per acre. Direct spray at the lower crown and stems using 50 gallons of water per acre. Irrigate immediately after application for soil penetration of 12 inches. Treat after egg laying during the summer.

USE 1 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS
 Borers (such as, Cottonwood, Peachtree(1), (1) For Peachtree borers, apply 3 sprays the 1st of May, then June 1st and the 3rd on July 1st) to flowering trees and shrubs of the genus Prunus as a trunk spray before newly hatched larvae enter the trees. Apply as a course, low pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.

6

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
WARNING: May be fatal if swallowed. Excessive absorption through skin may be fatal. Do not get in eye, sun or clothing. Avoid breathing of vapors and spray mist. Wear eye protection. Wear protective clothing and chemically resistant gloves when handling. Wash thoroughly after handling. Remove contaminated clothing and wash before reuse. Keep away from food, feedstuffs and water supplies. **NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.
ENVIRONMENTAL HAZARDS
 This pesticide is toxic to birds and aquatic life, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply or allow to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage: - Do not contaminate water, food or feed by storage and disposal. Store product in its original labeled container in a cool, dry, locked place out of reach of children. **Pesticide Disposal:** - Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or residue is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representatives at the nearest EPA Regional Office for guidance. **Container Disposal:** - a) Metal - Triple rinse (or equivalent). Then offer for recycling or reconditioning or disposal of in a sanitary landfill or by other approved State and local procedures. b) Plastic - Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill or incineration, or if allowed by State and local authorities, by burning. If burned stay out of smoke (2).

2

DORMANT SPRAY (TREE AND FOREST PESTS)

Use as a dormant or delayed dormant spray at the rates indicated to control the listed insect. Spray the entire tree to runoff using suitable ground spray equipment. Use the higher dosage for severe infestations. Use as recommended by State Ag. Ext. Service. **RESTRICTIONS:** - Make only one application during the dormant season. Do not allow meat or dairy animals to graze in treated areas.

USE 1/2 TO 1 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF FOLLOWING PESTS

Aphid (such as, Mealy Plum, Raw Apple, Woolly Apple), Borer (such as, Peach Twig), Cutworms (such as, climbing), Leafrollers (such as, Pandanus), Pear Psylla Adults, Plant Bugs, Scales (such as San Jose) Tank mix with 1 or 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water (7)
 (76-841SP PM412-08-83)



h6h

ORNAMENTALS (OUTDOOR AND NURSERY STOCK)

USE 1/2 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS

Apply when insects first appear or their damage is evident. Repeat as needed for control.
 Aphids (galls), Ants, Aphids (Apple, Cynanthemum, Cottonwood, Elm Leaf, Peach, Rose, Spirea, Woolly), Armyworms (Fall, Yellowlegged), Bagworms, Birch Leaf Miner, Birchwood Leaf Miner, Borer (Bugs), Cankworms, Citrus Mealy Bug, Cockroaches (American, Brownbanded, German, Oriental, Smoky Brown), Caterpillars (Fall Webworm, Grasshoppers, Green Fruitworms, Hornworms, Jack pine Budworms, Juniper Webworms, Katydid, Lace Bugs, Leafhoppers, Leafrollers, Maple Leaf-cullers, Moths (Clover, Red Spider, Southern, Red), Oriental Caterpillars, Orange Tortix, Periodical Cicada, Planthoppers, Poplar Tentmaker, Pyralis, Pines Caterpillars, Rose Chalcids, Sawflies (Pine Pin Oak Redheaded), Sawbugs, Spittlebugs, Spring Elm Caterpillars, Springtails, Spruce Budworm, Tent Caterpillars (Eastern, Forest, Western), Thornbug, Walnut Caterpillar, Western Spruce Budworm, Whiteflies, Yellow-necked Caterpillars.

For Bagworms treat when insects are in crawler stage. For effective control of leafrollers sp. should be applied before leaves are tightly rolled. For effective control of Maple Leaf Cullers on maple trees apply spray to larvae as ceciae are being formed. Do not treat sugar maple trees intended for maple syrup production. For effective control of Spider Moths when large numbers of eggs are present, apply a second spray 3 to 5 days in the south or 7 to 10 days in the north after initial treatment to control newly hatched nymphs.

USE 1 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS

Armyworms (Beet) Beetles (Fuller Rose, Native Elm Bark), Cutworms, Leafhoppers, Mealybugs, Mahogany Webworms, Memosa Webworms, Moths (such as, Brown-tail, Cypress tip, Douglas fir tussock, European Pine Shoot, Gypsy(2), Holly Bud, Nantucket Pine Tip, Pandora, Pitch Pine Tip, Subtropical Pine Tip, Tussock) Oakworms (such as California, Orangestemmed, Redhumped) Redhumped Caterpillar, Thrips (Exposed Stages), Weevils (such as, Blackvine (3), Yellow Poplar Pine Reproduction).

(1) - To reduce leg and branch feeding by bark beetles, applications should be made in the spring or early summer. (2) - To kill migrating and invading Gypsy moth larvae treat trunks and foliage. (3) - Blackvine Weevils are night feeders. Late afternoon spraying will give control in some areas.

3

USE 1 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS

Beetles (1) (such as, Cottonwood, Elm Leaf, Fuller Rose, Native Elm Bark, (2), Willow Leaf) (1) For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exist from nearby infested trees. For remedial treatment apply the spray to the main trunk of infested trees or logs when damage occurs but before adult beetles begin to emerge. For plantation trees apply to individual trees using a suitable hand or powered operated spray equipment.

(2) - To prevent Native Elm Bark Beetles from overwintering in uninfested trees, apply Dursban 4 E in water to the bottom 9 feet of the trunk. Wet the trunk thoroughly but do not spray to runoff. Care should be taken to apply the spray right to the base of the trunk. Application can be made with either a backpack mistblower or a hydraulic pressure sprayer from spring through to early fall. To reduce the leg and branch feeding on uninfested trees deemed to be of high value, apply a water spray to the tree crown. Application should be made in the spring or early summer using a sprayer that will give thorough coverage to the tree crown.

USE 1 PINT IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS

Beetles (such as, Ambrosia, Ambrosia Black Turp, Elm, (2), Formosa, Leaf, (1), Leaf European Elm Bark, Flea, Fuller Rose, Japanese, Juniper, Native Elm Bark, Southern Pine, Willow Leaf) (1).
 (1) For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees. For remedial treatment apply the spray to the main trunk of infested trees when damage occurs but before adult beetles begin to emerge.

USE 2 PINTS IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS

Weevils (such as, Northern Pine, Pines, Pitch spruce, Long(1) (1) For Southern Pine Seedlings treat immediately after transplanting. Treat each seedling with enough spray to thoroughly wet the foliage and stems to the point of runoff but do not use more than .5 gal. mix of spray per acre.

7

SUPPLEMENTAL LABELING
DURSBAN® 2E INSECTICIDE

See labeling affixed to DURSBAN 2 E product for instructions in using for INDOOR AND OTHER OUTDOOR PESTS

CONTROLS NUMEROUS PESTS AS LISTED ON ORNAMENTAL, OUTDOOR & NURSERY PLANTS, TREES AND FOREST PESTS OF NURSERY PLANTATIONS AND FIELDED TREES AND AS A DORMANT SPRAY FOR TREES AND FOREST PESTS

To be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, Mosquito Abatement Districts and Other Trained Personnel Responsible for Insect Control Programs

ACTIVE INGREDIENT	Chlorpyrifos (0.6-dialyl)	23.5%
	(3,5-Dichloro-2-pyridyl) phosphorothioate	76.5%
INERT INGREDIENTS		76.5%
		Total 100.0%

Contains 2 pounds of chlorpyrifos per gallon U.S. Patent No. 3,244,588 DURSBAN®

Reg. Trademark DowElanco
KEEP OUT OF REACH OF CHILDREN
WARNING

SEE DIRECTIONS FOR USE AND CAUTIONS ON SIDE PANELS
 EPA Registration No. 768-841 EPA Est. No. 768 GA 1

SURECO, INC.
 Fort Valley, GA 31030

BEST AVAILABLE COPY