

NOV 30 1994

SureCo, Inc.
c/o Mr. H. R. McLane
H. R. McLane, Incorporated
7210 Red Road, Suite 206
Miami, Florida 33143-5321

Subject: Revised Labeling to Remove from
Scope of Worker Protection Standard
Dursban .5% G Insecticide
EPA Registration Number 769-662
Your Application Dated October 27, 1994
(Previously Submitted August 23, 1994)

Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended labeling:

The Environmental Hazards section must be revised to:

This pesticide is toxic to birds, fish, and aquatic invertebrates. Birds feeding in treated areas may be killed. Clean up spilled product to reduce exposure to wildlife. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwaters.

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DTE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure



For Effective Long Lasting Control of Chinch Bugs and Sod Webworms in Ornamental Turf.

USE FOR MOUND TREATMENT CONTROL OF IMPORTED FIRE ANTS IN LAWNS AND RECREATIONAL AREAS

25 LBS TREATS UP TO 5,000 SQ.FT.

Controls Ants, Armyworms (Such as Beet, Fall, Yellowstriped), Brown Dog Ticks, Centipedes, Chiggers (1) Chinch Bugs, Clover Mites, Cutworms, Crickets, Earwigs, Fairy Skipper, Fleas, Gnats, Grasshoppers, Greenbugs, Leafhoppers, Lucerne Moth, Millipedes, Mites (Such as Clover, Formula Grass, Winter Grain), Mosquitoes (2), Sod Webworms (Lawn Moth)(3), Sow Bugs, Springtails, Ticks (1), Turfgrass Weevil (Hyperodes) (4), European Crane Fly (5), White Grubs (Such as Black Turfgrass Ateenia, European Chafer, Southern Chafer, Japanese Beetle, June Beetle larvae) (5), Bluegrass Billbugs(6) Fire Ants (7).

The numbers in parenthesis refer to SPECIFIC DIRECTIONS on side panel.

USE FOR CRAWLING INSECT CONTROL OUTSIDE OF HOMES, APARTMENTS, CONDOMINIUMS, WAREHOUSES, HOSPITALS, SCHOOLS, FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL BUILDINGS

ACTIVE INGREDIENT:

Chlorpyrifos: [O,O.-diethyl O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] - 0.50%

INERT INGREDIENTS: 99.50%

DURSBAN® is a Registered Trademark of The Dow Chemical Company

Total 100.00%

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or convulsing person. Get medical attention.

IF IN EYES Flush eyes with plenty of water.

IF ON SKIN In case of contact, remove contaminated clothing and wash affected area with soap and water.

NOTE TO PHYSICIAN Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote only by injection.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION May be harmful if swallowed. Avoid contact with eyes, skin and clothing. Avoid breathing dust. Wash thoroughly after handling and before eating or smoking. Wash contaminated clothing before reuse. Keep away from food, feedstuffs, and water supplies. When watering in of the insecticide is required children and pets should be kept off treated areas until material is washed into the soil and grass is dry.

ENVIRONMENTAL HAZARDS

This product is toxic to fish, birds, and other wildlife. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not contaminate water when disposing of equipment washwaters. (Use only as directed.)

SEE SIDE/PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENT

EPA REG. NO. 769-662

EPA EST. NO. 769 GA-1

NET WEIGHT (1 lb.; 5 lb.; 25 lb.; 50 lb.) POUND(S)

SURECO, INC.

Fort Valley GA 31030

NOV 30 1994

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 769-662

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not use for any purpose not listed on the label. Do not use for any purpose not listed on the label. Do not use for any purpose not listed on the label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal. **STORAGE:** Store product in its original labeled container in a cool, dry, locked place out of reach of children. **PESTICIDE DISPOSAL:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke. **HOUSEHOLD:** Do not re-use empty container. Wrap container and put in trash.

SPECIFIC DIRECTIONS

The following numbered paragraphs correspond to numbers in parenthesis (1) on the front panel following the insect names. Insects not numbered may be controlled by following the GENERAL DIRECTIONS. Repeat treatments as needed for control.

(1) CHIGGERS, TICKS, & FLEAS

Use 1 lb / 200 sq. ft. (5 lb./1,000 sq. ft.) for area control of these insects, infesting non-crop land areas such as roadsides, footpaths, and trails, picnic and camping sites, parks and other recreational areas where these pests are present and create a nuisance or a possible public health problem. DO NOT allow public use of treated area during applications. For Fleas (out-of-doors; only) treat sleeping quarters and other areas frequented by pets such as beneath porches and along fencing.

(2) MOSQUITOES

Mosquitoes coming to rest on areas treated for control of turf pests will be killed for varying periods of time after treatment depending on exposure of treated areas to weathering conditions.

(3) SOD WEBWORMS

For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment. Treat in the spring or when insects first appear troublesome. Additional treatments at 30 day intervals may be needed for new infestations. Apply at rate of 8 lb per 1,000 Sq Ft.

(4) TURF GRASS WEEVIL

Use 25 lbs per 2,500 sq ft. Treat suspected problem areas in mid-April and again in mid-May or as recommended by your local Agricultural Extension Service Specialist.

(5) WHITE GRUBS; CRANE FLY LARVAE

To control White Grubs (Black Turfgrass Ateenia, European Chafer, Southern Mole Cricket, Japanese Beetle and June Beetle Larvae) use 25 lb to treat 2,500 sq ft. After applications apply enough water to wash the insecticide into the turf. For best results remove thatch first & treat when grubs are young and actively feeding near the soil surface, usually during late July and August. Consult your local Agricultural Extension Service Specialist for best time to treat. European Crane Fly larvae will be controlled using 25 lb per 3,600 Sq Ft. or will be controlled simultaneously with the same application being applied for control of "White Grubs" at (25 lb/2,500 Sq Ft.). Consult your Agricultural Agent for proper time to apply. For best results, first remove thatch. Water the granules into the soil.

(6) BLUEGRASS BILLBUGS

Use 25 lb / 2,500 sq ft. for Billbug grubs. Remove thatch before treatment. Wash granules into soil.

(7) IMPORTED FIRE ANT CONTROL

MOUND TREATMENT: For treatment of Fire Ant mounds occurring in lawns, turf and premise areas, treat individual ant mounds measuring from 3 to 12 inches in diameter by applying 8 tablespoonfuls (1 1/2 oz.) of granules evenly in a circle around the outer edges (1 to 2 feet from the center) of the mound. Finish the treatment then by applying 8 tablespoonfuls (1 oz.) sprinkled evenly over the center of the mound. Best results will be obtained if mound is not disturbed after treatment thus allowing ants to come and go and to be exposed in the insecticide. Repeat treatments as needed for additional control but not more than once every 7 days. (Amend to remove from WPS - Sod Farms deleted - 1 subm'd 01-14-94)

GENERAL DIRECTIONS FOR USE IN LAWNS, RECREATIONAL AREAS AND ORNAMENTAL TURF:

Apply DURSBAN .5% and recreational turf areas. Apply to lawn and recreational turf areas.

For control of Ants, Ticks, Centipedes, Skipper, Fleas, Gnats, Millipedes, Mites (Lawn Moth), Sow Bugs, necessary to water.

Refer to SPECIFIC DIRECTIONS for control of the proper Spreader Set of water for control.

Area to be Treated
2,500 Sq Ft
5,000 Sq Ft
10,000 Sq Ft

USE FOR CRIMINAL HOMES, APARTMENTS, CONDOMINIUMS, WAREHOUSES, HOSPITALS, SCHOOLS, FOOD HANDLING ESTABLISHMENTS AND COMMERCIAL BUILDINGS

To protect against insects around the outside of homes, use 1 pound to treat. Apply when insects are in spring or early fall. Repeat more than once every 7 days. **SPREADER SETTING:** due to age and use, 1.5% into your spreader. Area of 500 Sq. Ft. (5' x 100') in the test, open the spreader at 1/2 inch, close (shut) it.

SPREADER
Scott Lawn Cyclone B1

Read these entire DIRECTIONS on the label and Directions For Use, it makes no other application or use consequences, special product.