

X-CEL VEG KILL

ACTIVE INGREDIENT:

Sodium Cacodylate	5.80%
Dimethylarsinic Acid	1.00%
OTHER INGREDIENTS:	<u>93.20%</u>
TOTAL	100.00%

Total arsenic content as elemental, all in water soluble form 3.25%



Contains the equivalent of 0.65 lbs. Dimethylarsinic Acid per gallon

KEEP OUT OF REACH OF CHILDREN

CAUTION

See side panel for Precautionary Statements and First Aid

NET CONTENTS: 32 FL. OZ. [1 Qt.]



PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed. Wash thoroughly with soap and water after handling. Avoid contact with eyes, skin, or clothing. Keep children and pets off treated areas until this material has been washed into the soil.

FIRST AID

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF ON SKIN: Wash with plenty of water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing give artificial respiration, preferably mouth to mouth. Get medical attention.

IF IN EYES: Flush with plenty of water. Call a physician if irritation persists.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwater, or rinsate.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. **READ ENTIRE LABEL BEFORE USING THIS PRODUCT. USE STRICTLY IN ACCORDANCE WITH LABEL PRECAUTIONARY STATEMENTS AND DIRECTIONS.**

When X-Cel Veg Kill is applied it kills most weeds and grasses on contact. X-Cel Veg Kill may be used around bushes trees or shrubs. Excellent for use around patios, sidewalks, driveways, fences, decks, retaining walls, foundations, and curb sides.

SPOT TREATMENT:

To kill unwanted vegetation along side walks, patios, driveways, fences, retaining walls, around shrubs, trees, and bushes. Use a hand operated sprayer only. Mix at rate of 32 oz. per 3 gallons of water. Covers 800 sq. ft. For tall weeds or grasses, cover foliage thoroughly and repeat in 5 days if necessary. To kill the leaves on Poison Ivy, Brambles, Briars, and Stump Sprouts, wet all foliage thoroughly and repeat if new growth develops. For most rapid kill, apply during warm, sunny weather. Do not apply during windy conditions that will carry spray to desired vegetation. Product will cause brown or yellowing dead spots in lawn.

TRIM AND EDGING TREATMENT:

For use along driveways, sidewalks, fence lines, patios, and parking lots. Apply anytime during the growing season. Apply using a hand operated sprayer only. Mix at rate of 32 oz. per 3 gallons of water. Thoroughly wet foliage of unwanted vegetation. Maximum control will be achieved during warm sunny weather and when unwanted vegetation is actively growing. Wilting will occur in 1 to 3 days and complete browning in about 5 to 7 days. Avoid drift to desirable

foliage. Heavy lawn grasses and difficult to control perennial weeds such as Bermudagrass and Dallisgrass will require more than one application to give complete control of top growth.

LAWN RENOVATION

Mow grass to about 1 inch height. Mix at rate of 32 oz. per 3 gallons of water. Apply X-Cel Veg Kill at any time during the growing season, preferably on a bright sunny day. Cover foliage thoroughly. X-Cel Veg Kill works only through foliage, not the root system. Average turf should be brown and dead within 5 days. Certain weeds and grasses may require a second application. When all vegetation is dead, treated area should be raked. Water area heavily prior to seeding. Seeding, sodding, or sprigging may be done 5 days after application. Keep area moist. On sunny days water if possible at mid-day until plants are established.

USE PRECAUTIONS

Care should be taken when treating weeds in flower beds or around shrubs and ornamental plantings. Do not allow spray to come in contact with foliage, green bark, grafted unions, or damaged bark when spraying around desirable plants. **Do not use for spot treatment of weeds in lawns.** Do not water within 24 hours after application. Do not apply if rainfall is expected within 24 hours after application.

STORAGE & DISPOSAL

Storage: Store product in original container in a cool, dry locked place out of reach of children.

Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash.

Container Disposal: Do not reuse container. Rinse thoroughly before discarding in trash.

SureCo, Inc. will not accept liability for damage or injury resulting from misuse.

Distributed by SureCo, Inc., an indirect subsidiary of Ringer Corp.,
9555 James Avenue South, Suite 200, Bloomington, MN 55431
(612) 703-3300

For information on this pesticide product (including health concerns, medical emergencies or pesticide incidents) call the National Pesticide Telecommunications Network at 1-800-858-7378.

If you are not completely satisfied with this product, or for consumer information, call (612) 703-3300 to arrange for refund of the purchase price or replacement of the product. Proof of purchase is required.

EPA Reg. No 769-664

EPA Est. No 192-CA-1 769-GA-1, 44616-MO-1

Circled letter corresponds to first letter of lot number on side of container

