

769-665

04/11/2000

1/10



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

Richard W. Fresh
Consep, Inc., A Subsidiary of Verdant Brands, Inc.
213 SW Columbia Street
Bend, Oregon 97702-1013

APR 11 2000

Subject: SMCP Sevin 10% Dust
EPA Registration No. 769-665
Amendment dated January 14, 2000 in Response to EPA
Letter dated 23 November 1999 in response to GDE.

Dear Mr. Fresh:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended labeling:

1. Correct the PHI for peanuts listed under "Forage Crops" to 14 days.

Submit two copies of your final printed labeling before you release the product for shipment. A copy of the labeling stamped "Accepted With Comments" is enclosed for your records.

Sincerely,

George Tompkins
George Tompkins, Ph.D., Entomologist
Insecticide-Rodenticide Branch
Registration Division (7505C)

SMCP SEVIN® brand carbaryl insecticide 10% DUST

Active Ingredient:	
Carbaryl (1-Naphthyl N-methylcarbamate	10.0%
Inert Ingredients:	90.0%
Total	100.0%

Sevin® is a registered trademark of Rhone-Poulenc for carbaryl insecticide

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

STATEMENT OF PRACTICAL TREATMENT

This product is an N-methyl carbamate and is a cholinesterase inhibitor.

IF SWALLOWED: Drink 1 or 2 glasses of water and induce vomiting by touching the back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person. Get medical attention.

IF INHALED: Remove victim to fresh air.

IF IN EYES: Flush eyes with plenty of water.

Note To Physician: Carbaryl is a moderate, reversible cholinesterase inhibitor. Atropine is antidotal. Do not use 2-PAM, opiates or cholinesterase inhibiting drugs.

EPA Reg. No. 769-665

EPA Est. No. 769-GA-1

Manufactured By:
SureCo, Inc.
Fort Valley, GA 31030

**ACCEPTED
with COMMENTS
In EPA Letter Dated:
APR 11 2000**

**Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under EPA Reg. No.
769-665**

**PRECAUTIONARY STATEMENTS
HAZARDS TO HUMANS AND DOMESTIC ANIMALS**

Caution: May be harmful if swallowed. Avoid breathing vapors (dust). Avoid contact with eyes, skin or clothing.

PERSONAL PROTECTIVE EQUIPMENT

Applicators and other handlers must wear: Long-sleeved shirt and long pants, waterproof gloves and shoes plus socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:
*Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
*Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to aquatic and estuarine invertebrates. For terrestrial uses, do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark, except for rice uses. Discharge from rice fields may kill aquatic and estuarine invertebrates. Do not contaminate water by cleaning of equipment or disposal of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl labels.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulations.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

Coveralls, Waterproof gloves, and Shoes plus socks.

Note To User: Avoid applying this material during very hot periods of the day.

VEGETABLES

APPLICATION: Begin applications when insect or damage is first observed and repeat on a regular 5 to 7 day schedule or as necessary for good insect control.

BEANS: To control Mexican Bean beetle, bean leaf beetle, cucumber beetles, leafhoppers, flea beetles, Japanese beetles, and corn earworm, apply at the rate of 10 pounds per acre. To control lygus bug, stink bug and lima bean pod borer, apply 20 pounds per acre.

CABBAGE, COLLARDS, BROCCOLI, BRUSSELS SPROUTS, CAULIFLOWER, LETTUCE, MUSTARD GREENS, TURNIPS, KOHLRABI AND SPINACH: To control flea beetle and harlequin bug, apply at the rate of 10 pounds per acre. To control corn earworm, armyworm, cabbageworm and small cabbage loopers, apply 10 to 20 pounds per acre.

CUCUMBERS, SQUASH, CANTALOUPE, PUMPKINS, MELONS: To control pickleworm, melonworm, cucumber beetles, squash bug, flea beetles, apply at the rate of 10 pounds per acre.

CORN: To control corn earworm, European corn borer, fall armyworm, sap beetle and Japanese beetles apply at the rate of 15 to 20 pounds per acre. For budworm damage in the whorl, apply to entire plant and repeat as necessary. For insects attacking silks and ears, apply at 2 to 3 day intervals starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending upon the severity of infestation.

CARROTS: To control flea beetles, leafhoppers, spittlebugs, lygus and stink bugs, apply at the rate of 10 to 20 pounds per acre.

LETTUCE: To control flea beetles, harlequin bug, and leafhoppers, apply at the rate of 10 pounds per acre. To control corn earworm, armyworms, cabbage caterpillars, spittlebugs, lygus and stink bugs, apply at the rate of 10 to 20 pounds per acre.

OKRA: To control corn earworm and stink bugs, apply at the rate of 10 to 20 pounds per acre.

TOMATO, EGGPLANT, PEPPERS: To control tomato fruitworm, fall armyworm, flea beetles, and leafhoppers, apply at the rate of 10 pounds per acre. For control of tomato hornworm, European corn borer and stink bugs, apply at the rate of 10 to 20 pounds per acre.

POTATO: To control Colorado potato beetle, leafhoppers and flea beetles, apply at the rate of 10 pounds and for lygus bugs, apply at 20 pounds per acre.

PREHARVEST INTERVAL FOR VEGETABLES: Beans = 3 days (succulent), 21 days (dried). Cabbage, broccoli, Brussels sprouts, cauliflower, kohlrabi = 3 days. Collards, mustard greens, lettuce (head & leaf), spinach, turnip (tops) = 14 days. Cucumbers, squash, cantaloupes, pumpkins, melons, okra, tomato, eggplant, peppers = 3 days. Turnips (roots), carrots, potato = 7 days. Corn (sweet) = 2 days (ears).

FORAGE CROPS

APPLICATION: Apply when insects or signs of damage appear and repeat 14 days later if necessary.

ALFALFA, CLOVERS, COWPEAS, PEANUTS AND SOYBEANS: To control Mexican bean beetle, blister beetles, thrips, leafhoppers, three-cornered alfalfa hopper, cucumber beetles, bean leaf beetle, Japanese beetle, alfalfa caterpillar and velvet bean caterpillar, apply at rate of 10 pounds per acre. To control armyworms, webworms, corn earworm, stink bugs and grasshoppers, apply at the rate of 10 to 15 pounds per acre. For alfalfa and clovers do not apply within 7 days of harvest or grazing; for cowpeas and soybeans do not apply within 14 days of grazing or harvest for forage or within 21 days of harvest of dried beans or peas, seed or hay. Peanuts have a 21 days preharvest interval. — 14

RICE, FORAGE GRASS, PASTURE: To control thrips, armyworms, stink bugs and grasshoppers, apply at the rate of 10 to 15 pounds per acre. For optimum thrip control in grasses grown for seed, direct dust to boot. For forage grass and pasture, do not apply within 14 days of harvest or grazing. Allow 14 days between last application and harvest on rice. Do not use on rice fields in which crayfish and/or catfish is included in the cultural practice.

CORN (Forage and fodder): To control earworm, European corn borer, fall armyworm, sap beetles and Japanese beetles, apply at the rate of 10 to 20 pounds per acre. Do not apply within 48 days of harvest of grain and fodder or within 14 days of harvest or grazing of forage or silage.

SORGHUM (Grain and forage): To control armyworms, webworms, corn earworms, stink bugs and grasshoppers, apply at the rate of 10 to 20 pounds per acre. Do not apply within 21 days of harvest for grain or fodder or within 14 days of harvest or grazing of forage or silage.

TOBACCO

To control hornworms, budworms, and flea beetles, apply at the rate of 15 to 25 pounds per acre. Begin applications when worms are small and repeat as necessary for control. Avoid excessive application. For budworm control with a hand duster, direct the dust to the bud and upper leaves of each plant, only 4 to 6 pounds per acre is necessary. Do not apply with cloth bags or apply heavily because plant injury may result.

POULTRY

CHICKENS, TURKEYS, DUCKS, GEESE, GAME BIRDS, AND PIGEONS: (LITTER TREATMENT): Apply 1/2 pound per 40 square feet to control Northern fowl mite, chicken mite, lice, fleas and bed bugs. Use a scoop and scatter by hand, or apply by hand duster. Avoid contamination of nests, eggs, feeding and watering troughs.

HAND APPLICATION TO BIRDS: Use 1/2 pound to treat 100 birds by means of a shaker can, squeeze bottle or hand duster to control Northern fowl mite, chicken mite, lice and fleas. Direct dust to vent and fluff areas. Repeat in 4 weeks if necessary. Hand application to birds for chicken mite and fleas is a supplement to litter treatment for control of these pests. For the control of sticktight fleas, it is especially important to treat necks and heads.

DUST-BATH BOXES: Use 2 1/2 pounds per box for each 100 birds to control Northern fowl mite, chicken mite, lice and fleas. This is a supplement to litter treatment for chicken mite and flea control. On ranges, one natural wallow may be substituted for each dust-bath box.

NOTE: Allow 7 days between application to poultry and game birds or premise treatment and slaughter of birds.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forest, nurseries, or greenhouses.

Do not allow children or pets to contact treated area until dust has settled.

RESIDENTIAL USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product.
- Change clothing as soon as possible after use,
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.

LAWNS

To control leafhoppers, fall armyworm, and sod webworm, apply at the rate of 2 pounds per 1000 square feet. For best results, apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 or 3 weeks if necessary.

EARWIGS

This material is effective in controlling earwigs and when properly used will reduce the number gaining entrance to homes and buildings. Apply heavily in a band 3 to 4 inches wide using a shaker-type can. Earwigs breed over wide areas around yards and fields and are driven by cold weather to seek shelter in warm buildings. As they move along the foundation wall seeking entrance, they come in contact with the treated barrier and are killed. Be sure to treat in front of all steps and other entrance ways as well as along the foundation wall. Treat also around refuse piles, lumber, mulch and other areas in the yard offering areas of concealment to earwigs. Repeat application in 10 days or when earwigs become troublesome. If the dust barrier is washed away by rain, it should be replaced with a new application if earwigs are still a problem.

PEST INSECT CONTROL (TICKS & FLEAS)

DOGS: Use this product only outside or in a well-ventilated area. Wearing household latex or rubber gloves, dust entire animal, avoiding pet's eyes, nose, mouth and genital areas. Rub or brush pet's hair to work dust down to the skin. Comb out dead ticks a few hours after treatment.

This product should not be applied by children. Do not use on pregnant dogs. Do not use on kittens or puppies under 12 weeks of age. Repeat at weekly intervals if needed.

DOG QUARTERS: Dust bedding and doghouse using a shaker can or other dust applicator. Force dust into cracks and crevices. Repeat as necessary.

HOME AND GARDEN USE

DIRECTIONS FOR USE: SMCP Sevin 10% Dust is to be used without dilution. Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin. Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering. For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week. Apply when insects or their damage appear. Foliage should be dry before dust is applied. Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

CROPS	INSECTS CONTROLLED	LAST APPLICATION ALLOWED BEFORE HARVEST
Asparagus (on Spears), Asparagus (On Ferns & Brush)	Asparagus Beetle, Do not repeat application within 3 days. Asparagus Beetle, Apache Cicada Post-Harvest application	1 Day
Beans	Lygus Bugs, Stink Bugs, Corn Earworm, Leafhoppers, Flea Beetles, Tarnished Plant Bug, Bean Leaf Roller, Mexican Bean Beetle, Bean Leaf Beetle, Western Bean Cutworm, Japanese Beetle, Velvetbean Caterpillar, Armyworms. Do not feed forage to dairy animals or animals being finished for slaughter.	3 days (succulent) 21 days (dried)
Cucumbers, Squash, Cantaloupes,	Striped Cucumber Beetle, Spotted Cucumber Beetle, Melonworm,	3 days

Watermelons	Pickleworm, Squash Bug, Leafhoppers, Flea Beetles	3 days
Potatoes, Tomatoes, Eggplants, Peppers	Flea Beetles, Armyworms (Fall), Tomato Fruit Worm, Corn Earworm, Tomato Hornworms, Leafhoppers, European Corn Borer, Stink Bugs, Colorado Potato Beetle, Lace Bug, Lygus Bugs	7 days (potato) 3 days (tomato, eggplant, pepper)
Lettuce	Flea Beetles, Stink Bugs, Corn Earworm, Leafhoppers, Armyworms, Harlequin Bug, Spittlebugs, Lygus Bugs	14 days (head & leaf)
Carrots	Flea Beetles, Leafhoppers, Six-spotted Leafhopper, (aster yellow vector), Armyworms, Spittlebugs, Lygus & Stink Bugs	7 days
Cabbage, Broccoli, Brussels Sprouts, Cauliflower, Kohlrabi	Flea Beetles, Harlequin Bug, Corn Earworm, Armyworms	3 Days
Okra	Corn Earworm, Stink Bugs	3 days
Strawberries	Spittlebugs, Strawberry Leaf Roller, Strawberry Weevil	7 days
Grapes	Grape Leafhoppers, Grape Leaf Folder, Grape Berry Moth, European Fruit Lecanium, Grape Leaf Skeletonizer, Japanese Beetle, June Beetles, Red-banded Leaf Roller	7 days

STORAGE AND DISPOSAL

STORAGE: Store pesticide in original container and place in a locked storage area. Opened pesticide containers should be securely resealed prior to storage. Protect from penetration of moisture, excessive heat or cold. Avoid cross-contamination with other pesticides or fertilizers. Use caution when storing pesticides in stacks.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Wastes resulting from the use of this product may be disposed of on site or at approved waste disposal facility.

CONTAINER DISPOSAL: Completely empty bag into application equipment, then dispose of empty bag in a sanitary landfill or by incineration or if allowed by state and local authorities, by burning. If burned, stay out of smoke.

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product. **DISCLAIMER:** SureCo warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. SureCo makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use, handling or application of this product.