

DEC 03 1996

SureCo, Inc.
c/o Mr. H. R. McLane
H. R. McLane, Inc.
7210 Red Road, Suite 206
Miami, Florida 33143-5321

Subject: Revised Labeling--Ant Species
Dursban 1E Insecticide
EPA Registration Number 769-666
Your Application Dated August 16, 1996

Dear Mr. McLane:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, is acceptable provided that you make the labeling change indicated below before you release the product for shipment bearing the amended labeling:

Restore the Non-Agricultural Use Requirements box. It may appear anywhere in the Directions for Use section after the Agricultural Use Requirements box. The box must contain the following text:

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Keep children and pets off treated areas until spray has dried.

274

(You may also add "(for all species of Ants, including Argentine, Carpenter, Ghost, and Pharaoh)" after "Ants" in the list of pests near the beginning of the section APPLICATION WITHIN FOOD HANDLING ESTABLISHMENTS.)

Submit five (5) copies of your final printed labeling before you release the product for shipment.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

DHE

Dennis H. Edwards, Jr.
Product Manager (19)
Insecticide-Rodenticide Branch
Registration Division (7505C)

Enclosure

DURSBAN® 1E INSECTICIDE

To be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, and Other Trained Personnel Responsible for Insect Control Programs.

CONTROLS NUMEROUS PESTS, AS LISTED, OF HOUSEHOLDS, TURF AND ORNAMENTAL PLANTS. FOR AREA CONTROL OF TICKS AND CHIGGERS. USE AS A MOUND TREATMENT TO KILL FIRE ANTS, ALSO CONTROLS CARPENTER ANTS

ACTIVE INGREDIENT:

Chlorpyrifos (0,0-diethyl,0-(3,5,6-

trichloro-2-pyridyl)(phosphorothioate)

12.9%

INERT INGREDIENTS:

87.1%

100.0%

Contains 1 pound Chlorpyrifos per gallon
DURSBAN Reg. Trademark DowGlance

KEEP OUT OF REACH OF CHILDREN WARNING / AVISO

Si usted no entiende la etiqueta, busque a alguien para que se le explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)

STATEMENTS OF PRACTICAL TREATMENT

THIS PRODUCT IS AN ORGANOPHOSPHATE. IF SWALLOWED, Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. DO NOT induce vomiting or give anything by mouth to an unconscious or comatose person. Get medical attention. IF INHALED: Remove victim to fresh air. Get medical attention. IF IN EYES: Flush eyes with plenty of water. Get medical attention if irritation persists. IF ON SKIN: In case of contact, remove contaminated clothing and wash affected area with soap and water. NOTE TO PHYSICIAN: Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine, only by injection is the preferable antidote.

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. Excessive absorption through skin may be fatal. Do not get in eyes, on skin or clothing. Avoid breathing of vapors and spray mist. Keep away from food, feedstuffs and water supplies.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category (F or G) on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear coveralls over short-sleeved shirt and short pants; Chemical Resistant gloves such as barrier laminata or Viton 14 mils. Chemical resistant footwear plus socks. Protective eyewear. Chemical resistant apron when cleaning equipment. A Dust/Mist filtering respirator (MSHA/NIOSH) approval number prefix TC-21C). Follow manufacturers instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should; Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet. Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing. Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds and wildlife, and extremely toxic to fish and aquatic organisms. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply or allow to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL AND CHEMICAL HAZARDS

Do not use, pour, spill or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Do not apply this product through any type of irrigation system.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) notification to workers, and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours. PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: coveralls over short sleeved shirt and short pants; Chemical Resistant gloves such as barrier laminata or Viton, 14 mils.; Chemical resistant footwear plus socks; Protective eyewear; Chemical resistant headgear for overhead exposures.

E.P.A. Reg. No. 769-666 NET CONTENTS 1 GALLON E.P.A. Est. No. 769-GA-1

SURECO, INC. - Fort Valley, GA 31030

STORAGE AND DISPOSAL

Storage: Avoid storage at high temperatures. Protect from moisture. Avoid contamination with water, acids, or alkalis. Do not store near heat or open flame. Keep container closed. Store in original container only. Separate pesticides to avoid cross-contamination of other pesticides, fertilizer, food and feed. Keep container in locked storage areas. If spilled, absorb with dry material such as sand or sawdust and sweep up for disposal. **Pesticide Disposal:** Do not contaminate water, food or feed by storage and disposal. Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal:** Metal - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by other procedures approved by State and Local authorities. Plastic - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or by incineration, or, if allowed by State and Local authorities, by burning. If burned stay out of smoke.

PESTS INDOORS

General Information: Use DURSBAN 1E insecticide to control the pests indicated in the areas listed below by applications as a 0.25% or 0.5% Chlorpyrifos spray. Use the 0.25% spray to control light infestations and the 0.5% spray to quickly reduce heavy infestations or as directed. To prepare the spray, dilute DURSBAN 1E with water or with a suitable base oil as shown in DILUTION RATES paragraph. Repeat treatment as necessary. Use only deodorized kerosene for oil based sprays to be applied in food handling establishments.

Do not introduce the spray into the air or allow the spray to contact food or food contacting surfaces. Areas treated may include dark corners of rooms, and closets; along and behind baseboards, beneath and behind stoves, refrigeration units and similar food preparation areas and equipment, floor drains and around plumbing and other utility installations. For ants apply to ant trails and wherever these pests may find entrance. Spot treatment may encompass CRACK & CREVICE treatment by applying small amounts of material directly into openings leading to voids and hollow spaces in walls, equipment legs and bases or which occur at joints between different elements of construction or between equipment and floors. Equipment capable of delivering a pinstream of insecticide should be used. Repeat treatment as needed but not more than once every 7 days in restaurants and similar food service establishments or more often than once every 14 days in other types of food handling establishments. In case of emergency (call back), Durban 1E may be applied after 2 days from the last treatment but emergency use should be limited to once per month.

APPLICATIONS OF THIS PRODUCT IN FOOD HANDLING ESTABLISHMENTS OTHER THAN AS A SPOT AND/OR CRACK AND CREVICE TREATMENT ARE NOT PERMITTED.

APPLICATION WITHIN RESIDENTIAL BUILDINGS

Including Homes and Apartment Buildings and Within Nonfood Areas of Industrial, Institutional and Commercial Buildings including Hospitals, Stores, Manufacturing Plants and Warehouses: For Spot Treatment Only: Apply as a course spray or with a paint brush to localized area where ants for all species of ants, including Argentine, Carpenter, Ghost and Pharaoh; apply spray to ant trails and wherever they are seen; Cockroaches, Clover mites, Confused Flour Beetles, Crickets, Earwigs, Firebrats, Indian Meal Moth, Mediterranean Flour Moth, Rice & Granary Weevils, Millipedes, Red Flour Beetle, Saw Toothed Grain Beetle, Silverfish, and Spiders are found or normally occur including dark corners of rooms and closets; cracks and crevices in walls, along and behind baseboards, beneath and behind sinks, stoves, refrigerators and cabinets; around plumbing and other utility installations. For ants apply to ant trails, also around doors and windows and wherever else these pests may find entrance. For Clover mites use 0.5% spray only. Note: A period of 4 to 7 days is normally required for maximum effect on cockroaches. For the Control of Carpet Beetles, thoroughly apply a 0.5% spray as a spot treatment along baseboards and edges of carpeting, under carpeting, rugs, furniture, in closets and on shelving, and wherever else these insects are seen or suspected.

DILUTION RATES

To make a 0.25% spray solution:

FOR WATER SPRAY - mix 2 2/3 oz. in 1 gallon of WATER.

FOR AN OIL SPRAY - mix 2 fl. oz. in 1 gallon of DEO-BASE OIL.*

To make a 0.5% spray solution:

FOR WATER SPRAY - mix 5 1/3 oz. in 1 gallon of WATER.

FOR AN OIL SPRAY - mix 4 fl. oz. in 1 gallon of DEO-BASE OIL.*

DILUTION RATES (Cont.)

For larger area multiply the fluid ounces (amount) in 1 gallon by number of gallons required. *The "Job", "Deodorized base Kerosene.

NOTE: Where quick knockdown and/or flushing action is desired, dichlorvos (DDVP) or pyrethrin may be added to the spray mixture containing Durban 1E insecticide. For best results, use dichlorvos at a concentration of 0.25% in both water and oil-base sprays. Consult labeling for the dichlorvos or pyrethrin product used to determine the amount needed to give the desired spray concentration. Also read and observe all use and handling precautions for these products. DO NOT use Durban 1E in combination with dichlorvos (DDVP) or pyrethrin for treatment of food areas of food handling establishments.

APPLICATION WITHIN FOOD HANDLING ESTABLISHMENTS

PLACES OTHER THAN PRIVATE RESIDENCES IN WHICH FOOD IS HELD, PROCESSED, PREPARED OR SERVED INCLUDING, BUT NOT LIMITED TO; RESTAURANTS, GROCERY STORES, BAKERIES, BOTTLING PLANTS, CANNERIES, AND GRAIN MILLS: Mix Durban 1E according to the DILUTION RATES. USE AS A SPOT TREATMENT to control Ants, Cockroaches, Clover mites, Confused Flour Beetles, Crickets, Earwigs, Firebrats, Indian Meal Moth, Mediterranean Flour Moth, Rice & Granary Weevils, Millipedes, Red Flour Beetle, Saw Toothed Grain Beetle, Silverfish, and Spiders in and around both food and nonfood areas by application as a coarse, low pressure (20 psi or less) spray to localized areas where the above pests have been seen, suspected of hiding or entering. Apply only enough spray to thoroughly cover the surfaces treated using special care to avoid unnecessary runoff.

Brown Dog Ticks: Thoroughly apply 0.5% spray as a spot treatment to infested areas such as pet beds and resting quarters, nearby cracks and crevices; along and behind baseboards, window and door frames and localized areas of floor and floor covering where these pests may be present. Old bedding of pets should be removed and replaced with clean, fresh bedding after treatment. Do not treat pets with this product.

NOTE: (Pests Indoors): Use oil base spray only for conduits, motor housing, junction or switch boxes. Prior to application de-energize electrical equipment. After application, allow time for oil to evaporate before starting motors or placing switch in the "ON" position. DO NOT use water-base sprays in these places. Do not use in poultry houses and other animal buildings. Do not use in serving areas while food is exposed. Thoroughly wash dishes and food handling utensils with soap and water if they become contaminated by application of this product. Do not allow children or pets to contact treated surfaces until spray has dried. Remove pets and cover fish bowls (tanks) before spraying.

PESTS OUT-OF-DOORS

General Information: Use Durban 1E for the control of certain pests of lawns and ornamental plants, for area control of ticks and chiggers. Consult the following directions for dosage and application details to control the pests listed.

PESTS OF LAWNS AND OTHER ORNAMENTAL AND RECREATIONAL TURF GRASS AREAS

Use Durban 1E to control the pests listed on the following table by application at the recommended dosage and in accordance with the directions given below. Dilute Durban 1E in enough water to obtain complete and uniform coverage of pest infested areas and apply as a coarse, low pressure spray using suitable application equipment. Thoroughly water immediately after treatment to wash the insecticide into the turf, except as noted. For best results, the lawn should be moist at the time of treatment. Spray when pests first appear. Repeat as needed.

SURFACE INSECTS

Use 3 fl. oz./1,000 sq. ft. to control Ants, for all species of ants, including Argentine, Carpenter, Ghost, and Pharaoh; ants apply spray to ant trails and wherever these pests are seen; Armyworms, Brown Dog Ticks, Centipedes, Chiggers, Chinchbugs, Clover Mites, Crickets, Cutworms, Earwigs, Fire Ants (mounds), Fleas, Gnats, Grasshoppers, Greenbugs, June beetles, Leafhoppers, Lucerne Moths, Millipedes, Mites (clover, formula, winter grain), Sod webworms, Sowbugs, Springtails and Ticks. Spray when pests first appear, retreat as needed. For Sod Webworms, delay watering or mowing the treated area for 12 to 24 hours after treatment.

SOIL INSECTS

For Turfgrass Hyperodes Weevils use 6 fl. oz./1,000 sq. ft. Spray suspected problem areas in mid April and again in mid May or as recommended by your local Ag. Extension Service Specialist for your area. For European Crane Fly Larvae use 4 fl. oz./1,000 sq. ft. For White Grubs, (Black Turfgrass Athanas, European chaffer, Southern masked chaffer & Japanese beetle larvae) use 6 to 12 fl. oz./1,000 sq. ft. Spray when grubs are young and actively feeding near the soil surface, usually during late July and August or as recommended by local Ag Extension personnel. Immediately after spraying, irrigate the treated area with 1/2 to 1 inch of water to wash the insecticide into the underlying soil.

For Bl...
adults first

PRE

To help prevent
building, elec
and may find
and apply as

Use Durban
rodenticides, for
around dwellin
a disease vec

fluid ounces
sprayer or ot
grass areas,
obtain throu
use of treatm
weather cond
streams, pond
be killed if the

Use Durban
evergreens for
water accordi
equipment in
apply a wetting
areas. Attemp
excessive run
needed. ATTE
Keep out of h
azales, cam

Use 1 fl. oz./2
Fall/Winter
flies, Yellow

Use 2 fl. oz./
Thrips/Exo

Use 4 fl. oz./8
Lecanium, Pin

For larger amo
required for th
ornamental pl
depending on

Mix 4 fl. oz. in
1 gallon of
surrounding
degrees F. ea
sprays may di
(DCC-AMD, PM

27
S

(Outd

10

494

USE 2 TO 4 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF BEETLES.

Leaf Beetles (such as; Cottonwood Leaf, Elm Leaf, Willow leaf, For Cottonwood Leaf Beetles, use to control larvae and adults infesting cottonwoods. Apply when insects are present or when foliage damage is imminent.

USE 4 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Borers (Clearwing moth (such as; Ash, Dogwood, Lesser Peachtree, Lilac, Oak, Rhododendron), Metallic wood (such as; Bronze Birch, Flatheaded Appletree, Twolined Chestnut, Longhorned Beetles, (such as; Locust, Red Oak), Leaf miners, Needleminer (such as; Jeffrey Pine, Lodgepole Pine, Spruce), Scale Crawlers (Cottony Cushion, Cottony Maple, Euonymus scale, Fitchers scale, Florida Wax scale, Golden Oak, Hemispherical, Lecanium, Magnolia, Oak Kermes, Oystershell, Pine Needle, San Jose, Tea, White Birch)

(1) - For borer apply to the trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your Ag. Ext. Service for time to spray. Apply uniformly as a coarse low-pressure spray.

USE 12 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Borers (such as; Cottonwood, Peachtree)(1); (1) For Peachtree borers, apply 3 sprays (the 1st of May, then June 1st and the 3rd on July 1st) to flowering trees and shrubs of the genus Prunus as a trunk spray before the newly-hatched larvae enter the trees. Apply as a coarse, low pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.

USE 2 PINTS IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Beetles (such as; Ambrosia, Anobidae, Black Turpentine, Cottonwood Leaf, Elm Leaf, European Elm Bark, Flea, Fuller Rose, Japanese, June, Native Elm Bark, Southern Pine, Willow Leaf)(1);

(1) - For preventive treatment apply the spray to the main trunk of trees in the early spring or when threat of attack exists from nearby infested trees.

For remedial treatment apply the spray to the main trunk of infested trees when damage occurs but before beetles, begin to emerge.

4

TREE AND FOREST PESTS (NURSERIES, PLANTATIONS, AND FELLED TREES)

USE 1 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Adelgids (such as; Cooley and Eastern Spruce Gail, Pine Bark galls), Aphids (Apple, Chrysanthemum, Cottonwood, Elm Leaf, Peach, Rose, Spirea, Woolly), Bagworms, Boxelder Bugs, Cankerworms, Caterpillar Sphinx, Citrus mealybugs, Elm Spanworm, Fall Webworm, Greenstriped Maple worm, Jackpine Budworms, Juniper Webworms, Katydid, Lace Bugs, Leafhoppers, Leafrollers(2), Maple Leafcutters(3), Mites (Clover, Red Spider, Southern, Red), Oak Skeletonizers, Periodical Cicada, Poplar Tentmaker, Psyllids, Puss Caterpillars, Sawflies (Pine, Pin Oak,), Spring Elm Caterpillars, Spruce Budworm, Tent Caterpillars (Eastern, Forest, Western), Walnut Caterpillar, Western Spruce Budworm, Yellownecked Caterpillars, (1) - For Bagworms, treat when insects are in crawler stage. (2) - For effective control of Leafrollers spray should be applied before leaves are tightly rolled. (3) - For Maple Leafcutter on maple trees apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. (4) - For effective control of Spider Mites, when large numbers of eggs are present, apply a second spray after 3 to 5 days in the south or 7 to 10 days in the north after initial treatment to control newly hatched nymphs.

USE 2 FL. OZ. IN 3 GALLONS OF WATER FOR THE FOLLOWING INSECTS.

Beetles (Fuller Rose, Native Elm Bark), Leafhoppers, Mahogany webworms, Mealybugs, Mimosa Webworms, Moths (such as; Brownail, Cypress tip, Douglas fir tussock, European Pine Shoot, Gypsy(2), Holly Bud, Nantucket Pine Tip, Pandora, Pitch Pine Tip, Subtropical Pine Tip, Tussock), Oakworms (such as; California, Orangestriped, Redhumped), Redhumped Caterpillar, Thrips (Exposed Stages), Weevils (such as; Blackvine (3), Yellow Poplar, Pine Reproduction).

(1) - To reduce twig and branch feeding by bark beetles, applications should be made in the spring or early summer. (2) - To kill migrating and invading Gypsy Moth Larvae treat trunk and foliage. (3) - Blackvine weevils are night feeders. Late afternoon spraying will give control in some areas.

5

USE 2 TO 4 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF BEETLES.

Beetles (such as; Cottonwood Leaf, Elm Leaf, Willow leaf, (1) - For Cottonwood Leaf Beetles, use to control larvae and adults infesting cottonwoods. Apply when insects are present or when foliage damage is imminent.

USE 4 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Borers (Clearwing moth (such as; Ash, Dogwood, Lesser Peachtree, Lilac, Oak, Rhododendron), Metallic wood (such as; Bronze Birch, Flatheaded Appletree, Twolined Chestnut, Longhorned Beetles, (such as; Locust, Red Oak), Leaf miners, Needleminer (such as; Jeffrey Pine, Lodgepole Pine, Spruce), Scale Crawlers such as (Cottonyushion, Cottony Maple, Euonymus scale, Fitchers scale, Florida Wax scale, Golden Oak, Hemispherical, Lecanium, Magnolia, Oak Kermes, Oystershell, Pine Needle, San Jose, Tea, White Birch)

(1) - For borers apply to trunks and lower limbs of trees and shrubs when the adults begin to emerge. Consult your Ag. Ext. Service for proper time to spray. Apply uniformly as a coarse spray.

(2) For Cranberry Girdler larvae infesting Douglas Fir seedlings apply 1 quart per acre. Direct spray at the lower crown and stems using 50 gallons of water per acre. Irrigate immediately after application for soil penetration of 1-2 inches. Treat after egg laying during the summer.

USE 12 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Borers (such as; Cottonwood, Peachtree)(1); (1) For Peachtree borers, apply 3 sprays (the 1st of May, June 1st and the 3rd on July 1st) to flowering trees and shrubs of the genus Prunus as a trunk spray before newly hatched larvae enter the trees. Apply as a coarse, low pressure spray. Thoroughly wet all bark areas from ground level to scaffold limbs.

6

ORNAMENTALS (OUTDOOR AND NURSERY STOCK)

USE 1 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Apply when insects first appear or their damage is imminent. Repeat as needed for control. Adelgids(galls), Ants, Aphids (Apple, Chrysanthemum, Cottonwood, Elm Leaf, Peach, Rose, Spirea, Woolly), Armyworms (Fall, Yellowstriped), Bagworms, Birch Leaf Miner, Boxwood Leaf Miner, Boxelder Bugs, Cankerworms, Citrus Mealy Bugs, Cockroaches (American, Brownbanded, German, Oriental, Smokey Brown), Catalpa Sphinx, Elm Spanworm, Fall Webworm, Grasshoppers, Green Fruitworms, Hornworms, Jackpine Budworms, Juniper Webworms, Katydid, Lace Bugs, Leafhoppers, Leafrollers, Maple Leafcutters, Mites (Clover, Red Spider, Southern, Red), Oleander Caterpillars, Orange Tortix, Periodical Cicada, Plants Bugs, Poplar Tentmaker, Psyllids, Puss Caterpillars, Rose Chafers, Sawflies (Pine, Pin Oak, Redheaded), Sowbugs, Spittlebugs, Spring Elm Caterpillars, Springtails, Spruce Budworm, Tent Caterpillars (Eastern, Forest, Western), Thornbug, Walnut Caterpillar, Western Spruce Budworm, Whiteflies, Yellownecked Caterpillars,

For Bagworms, treat when insects are in crawler stage; For effective control of leafrollers spray should be applied before leaves are tightly rolled; For effective control of Maple Leaf Cutters on Maple trees apply spray to larvae as cases are being formed. Do not treat sugar maple trees intended for maple syrup production. For effective control of Spider Mites, when large numbers of eggs are present, apply a second spray 3 to 5 days in the south or 7 to 10 days in the north after initial treatment to control newly hatched nymphs.

USE 2 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF THE FOLLOWING INSECTS.

Amygdal (Bead), Beetles, (Fuller Rose, Native Elm Bark), Cutworms, Leafhoppers, Mealybugs, Mahogany Webworms, Mimosa Webworms, Moths (such as; Brownail, Cypress tip, Douglas fir tussock, European Pine Shoot, Gypsy(2), Holly Bud, Nantucket Pine Tip, Pandora, Pitch Pine Tip, Subtropical Pine Tip, Tussock), Oakworms (such as; California, Orangestriped, Redhumped), Redhumped Caterpillar, Thrips (Exposed Stages), Weevils (such as; Blackvine (3), Yellow Poplar, Pine Reproduction).

(1) - To reduce twig and branch feeding by bark beetles, applications should be made in the spring or early summer. (2) - To kill migrating and invading Gypsy moth larvae treat trunks and foliage. (3) - Blackvine Weevils are night feeders. Late afternoon spraying will give control in some areas.

3

Refer to Label Affixed to Container for Instructions Concerning the Worker Protection Standard Requirements PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

WARNING: May be fatal if swallowed. May be fatal if absorbed through skin or clothing. Avoid breathing of vapors. Keep away from food, feedstuffs and water supplies. **NOTE TO PHYSICIAN:** Chlorpyrifos is a cholinesterase inhibitor. Treat symptomatically. Atropine is an antidote.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish, birds, and other wildlife. Do not apply directly to water. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. Cover or incorporate spills. Do not contaminate water when disposing of equipment washwaters.

CHEMICAL & PHYSICAL HAZARDS

Do not use, pour, spill or store near open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Storage - Do not contaminate water, food or feed by storage and disposal. Store product in its original labeled container in a cool, dry, locked place out of reach of children. **Pesticide Disposal** - Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If these wastes cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. **Container Disposal** - a) Metal - Triple rinse (or equivalent). Then offer for recycling or reconditioning or dispose of in a sanitary landfill or by other approved State and Local procedures. b) Plastic - Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by State and local authorities, by burning. If burned stay out of smoke.

2

DORMANT SPRAY (TREE AND FOREST PESTS)

Use as a dormant or delayed dormant spray at the rates indicated to control the listed insect. Spray the entire tree to runoff using suitable ground spray equipment. Use the higher dosage for severe infestations. Use oil as recommended by State Ag. Ext. Service **RESTRICTIONS** - Make only one application during the dormant season. Do not allow meat or dairy animals to graze in treated areas.

USE 1 to 2 FL. OZ. IN 3 GALLONS OF WATER FOR CONTROL OF FOLLOWING PESTS.

Aphid (such as; Mealy Plum, Roxy Apple, Woolly Apple), Borer (such as; Peach Twig), Cutworms, (such as; climbing), Leafrollers (such as; Pandemis), Pear Psylla Adults, Plants Bugs, Scale (such as; San Jose)

Tank mix with 1 or 2 gallons of a petroleum spray oil recommended for dormant use in 100 gallons of water(7), (888AMOSP.PM4 / 01-03-9508-06-06

ACCEPTED with COMMENTS in EPA Letter Dated

DEC 03 1996

Under the Federal Insecticide, Fungicide, and Rodenticide Act no amended, for the pesticide registered under EPA Reg. No. 769-166

SUPPLEMENTAL LABELING DURSBAN® 1E INSECTICIDE

See labeling affixed to DURSBAN 1-E INSECTICIDE product for instructions in using for INDOOR AND OTHER OUTDOOR PESTS.

CONTROLS NUMEROUS PESTS, AS LISTED, ON ORNAMENTAL OUTDOOR & NURSERY PLANTS, TREES AND FOREST PESTS OF NURSERIES, PLANTATIONS AND FELLED TREES, AND AS A DORMANT SPRAY FOR TREES AND FOREST PESTS

To be Applied Only By or Under the Supervision of Pest Control Operators, Public Health Organizations, and Other Trained Personnel Responsible for Insect Control Programs.

ACTIVE INGREDIENT: Chlorpyrifos [0,0-diethyl, O-(3,5,6-trichloro-2-pyridyl) phosphorothioate] 12.9%
INERT INGREDIENTS: 87.1%
Total 100.0%

Contains 4 pounds of chlorpyrifos per gallon U.S. Patent No. 3,244,586 DURSBAN®

Reg. Trademark Dow Chemical Co.
KEEP OUT OF REACH OF CHILDREN WARNING/AVISO

SEE DIRECTIONS FOR USE AND CAUTIONS ON INSIDE PANELS.
E.P.A. Registration No. 799-606 E.P.A. Est. No. 799-GA-1

SURECO, INC.
Fort Valley GA 31030