

520371

MAR 18 1997

Ms. Stacy Schmidt
SureCo, Inc.
10012 N. Dale Mabry Hwy., Suite 221
Tampa, FL 33618

Dear Ms. Schmidt:

Subject: Amendment- Comply with Agency letter of November 22, 1996
SMCP Diazinon AG500
EPA Registration Number 769-689
Your submission dated February 17, 1997

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling:
 - a) The heading RESTRICTED USE PESTICIDE has the same type size requirements as the signal word -see 40 CFR 156.10 (h)(1)(iv). Also, the heading should appear on a separate line, with the "Due to Avian and Aquatic Toxicity" on the next line, rather than just "Due to Aquatic Toxicity".
 - b) On the container label and in the booklet, in the ingredients statement, revise ACTIVE INGREDIENTS to ACTIVE INGREDIENT, since there is only one.
 - c) On the container label and in the booklet, under the heading PERSONAL PROTECTIVE CLOTHING the usual chemical resistance category for 48% EC diazinon products is "F", not "G" as shown on your label. Also, revise "such s Barrier Laminate" to "such as Barrier Laminate".

- d) On the container label and in the booklet, in the Environmental Hazards revise "If waterfowl, i.e., ducks or geese, can be" to "On lawns, if waterfowl (ducks or geese) can be". Also, revise "water "lawn/areas" to just "water.lawn".
- e) In the booklet, in the Agricultural Use Requirements block, delete "of 24 hours" after "during the restricted entry interval (REI)" because there are different REIs for some groupings. Therefore, under each crop group heading the appropriate REI will have to be given (see below).
- f) In the booklet, directly under the heading **Fruit and Nut Crops** add "REI - 24 hours".
- g) In the booklet, add "to wait" between "days" and "between" in the PHI note under the PHI table for Fruits and Nuts.
- h) In the booklet, in the directions for the crop Caneberries, delete "Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur." from the Timing/Special Directions for control of Leafhoppers, etc. and delete "Do not apply within 7 days of harvest or illegal residues may occur." from the Timing/Special Directions for control of Dryberry Mites, then under all the directions for Caneberries add a Notes section as follows:

Notes: (1) Do not apply more than 2 qts. of product per acre per application and no more than 5 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest.

- i) For the crop Cranberries, in the directions for control of Blackheaded Fireworms, replace "Apply when pest appears." with "Apply to larval stage. Pheromone trap captures and sweep monitoring may be used to optimize treatment timing.". Retain the rest of the text as is, except that you may chose to delete the PHI sentence and have it appear in the revised NOTES (see item "j", below).
- j) Also for the crop Cranberries, for Cranberry Fruitworms the rate should be 2-3 qts. and in the directions replace the current text with "Cranberry growth stage and/or berry inspection for eggs may be used to optimize treatment timing; consult your local extension agent or pest management adviser. Repeat application every 14 days, as needed. A maximum of 4 applications per season is permitted for the 3 qt. rate. A maximum of 6 applications is permitted for the 2 qt. rate.". These revised directions do not include the PHI sentence since it will appear in the Notes. Add a NOTES section as follows:

NOTES: (1) Do not apply more than 12 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 7 days between the last application and harvest. (3) This product may be applied to cranberries through sprinkler irrigation. Refer to the section CHEMIGATION for specific chemigation directions for use. (4) Do not use water from irrigated or flooded cranberry beds for drinking purposes or to irrigate crops other than those appearing on EPA approved diazinon labels.

- k) In the booklet, for the crop Strawberries delete "Do not apply within 5 days of harvest or illegal residues may occur." from the Timing/Special Directions for control of Aphids, etc. and for control of Cyclamen Mites. Also, for Cyclamen Mites delete "Application may be made to plants after harvest." since this use is not supported by the residue data submitted by CIBA for this crop. Then, add a NOTES section as follows:

Notes: (1) Do not apply more than 1 qt. of product per acre per application and no more than 4 qts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 5 days between the last application and harvest.

- l) In the booklet, in the directions for the crop Walnuts delete "Do not make more than 3 applications per season. Do not apply within 45 days of harvest or after husks open, or illegal residues may occur." from the Timing/Special Directions and then add a Notes section as follows:

Notes: (1) Do not apply more than 3 qts. of product per acre per application. (2) Do not apply more than 9 qts. of product per acre per season. (3) To avoid illegal residues, allow a minimum of 45 days between the last application and harvest and do not apply after husks open.

- m) In the booklet, directly under the heading **Vegetable Crops** add "REI - 24 hours".
- n) In the booklet, add "to wait" between "days" and "between" in the PHI note under the PHI table for Vegetables.
- o) In the booklet, for all Vegetable Crops entries that include the pest Mole Crickets, you may replace "Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil." with "One to two days before planting, broadcast and immediately incorporate into the top 1-2 inches of soil."
- p) In the booklet, for all Vegetable Crops entries that include the pest Root Maggots, you may replace "Apply to plant beds

or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil." with "Broadcast just before planting and immediately incorporate into the top 3-4 inches of soil."

- q) In the booklet, for the crop Sweet Potatoes, delete the pest Flea Beetles since soil incorporation is not appropriate as a control for this foliar pest.
- r) In the booklet, directly under the heading **Field Crops** add "REI - 24 hours".
- s) In the booklet, add "to wait" between "days" and "between" in the PHI note under the PHI table for Field Crops.
- t) Revise the Note under the directions for the crop Sugar Beets to read "Sugar beet tops may be fed to beef and dairy animals."
- u) In the booklet, directly under the heading **Insect Control on Ornamentals** add "REI - 12 hours".
- v) In the booklet, in the directions paragraph for Insect Control on Ornamentals, add "Apply when pests first appear." before "Try to spray...". And add "Repeat as necessary." after "... and penetrate dense foliage."
- w) In the booklet, in the Precautions under the heading **Lawn Pest Control**, revise "if waterfowl, i.e., ducks or geese, can be" to "if waterfowl (ducks or geese) can be" and revise "1/4 inch water" to "1/4 inch of water".
- x) In the booklet, in the **Lawn Pest Control** table revise the Remarks for control of Frit Flies to read "For frit flies, before treatment mow grass and water well, but delay application until grass is dry."

3. Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records. The Agency notes "AllPro" in place of "SMCP" in the product name. Do you intend to change the master name of record for this product?

In compliance with the Diazinon Registration Standard

you must submit revised Confidential Statement(s) of Formula which must include upper certified limits for the impurities of toxicological concern as identified in the Diazinon Registration Standard (TEPP, O,S-TEPP, S,S-TEPP, and oxo-diazinon), and include signed and dated certification statements relative to FIFRA Sec 12 (a)(1)(C) and the method of arriving at the certified limits for the impurities (i.e., reliance on Ciba-Geigy's analysis, or another producer's analysis if another source of diazinon is used) and addressing data requirements under Guidelines 61-3 and 61-2b. If you have more than one diazinon supplier or Basic and Alternate Formulas, separate CSF forms should be submitted for each supplier for each Basic and Alternate Formula since the levels of toxicologically significant impurities are different for each diazinon technical and the certification of analyses are only appropriate to each supplier's own material (the Agency notes that one of your previously indicated suppliers is apparently no longer supplying this active ingredient). The text of these certification statements is as follows:

I hereby certify that, for the purposes of FIFRA Sec.12(a)(1)(C), the description of the composition of [your product's name], EPA Reg. No. [insert correct number for your product], refers to the composition set forth on this Statement of Formula and supporting materials. This description includes the representation that (1) no ingredient will be present in the product in an amount greater than the upper certified limit or in an amount less than the lower certified limit specified for that ingredient in this Statement of Formula; and (2) if the Agency requires that the source of supply of an ingredient be specified, that all quantities of such ingredient will be obtained from the source specified in the Statement of Formula.

I certify that the upper certified limits for the impurities listed on this Statement of Formula are based on [your supplier's name] analysis and certification of limits for this product based on [name of source product that your supplier has analyzed]. I further certify under Section 61-3 and 61-2(b) that there are no chemical reactions between diazinon and/or its impurities with the intentionally added inerts.

If you do not already have the impurities information required on the CSF(s), please immediately contact your supplier(s) directly.

Sincerely,

George T. LaRocca
 Product Manager 13
 Insecticide/Rodenticide Branch
 Registration Division (7505C)

RESTRICTED USE PESTICIDE. Due to Aquatic Toxicity.
For retail sale to, and use only by, Certified Applicators or persons under their direct supervision
and only for those uses covered by the Certified Applicator's certification.

AIIPro™

Diazinon AG500

For Control of Certain Insects on Fruits, Nuts, Vegetables, Field Crops, and Ornamentals.
THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.

ACTIVE INGREDIENTS:

Diazinon [0, 0-diethyl 0- (2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate].....	48.0%
INERT INGREDIENTS:	52.0%
TOTAL	100.0%

*contains xylene range aromatic solvent.

Contains 4 lbs. Diazinon per gallon

E.P.A. Reg. No. 769-689 E.P.A. Est. No. 769-GA-1

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

Si usted no entiende la etiqueta, busque a alguien para se la explique a usted en detalle.
(If you do not understand the label, find someone to explain it to you in detail.)
See Side Panel For Additional Precautionary Statements

STATEMENT OF PRACTICAL TREATMENT

This is an organophosphate insecticide.

IF SWALLOWED: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. Do not give anything by mouth if person is unconscious or convulsing.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash promptly with soap and water. Rinse thoroughly. Get medical attention.

IF IN EYES: Rinse eyes with plenty of water. Call a physician if irritation persists.

NOTE TO PHYSICIAN: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains a xylene range aromatic solvent which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

NET CONTENTS:
ONE GALLON

ACCEPTED
with COMMENTS
in EPA Letter Dated

MAR 18 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

769-689

2/17

RESTRICTED USE PESTICIDE
Due to Air and Aquatic Toxicity

For retail sale to and use by Certified Applicators or persons under their direct supervision and only for those uses covered by the Certified Applicator's Certification.

THIS PRODUCT MUST NOT BE USED ON GOLF COURSES AND SOD FARMS.

All Pro DIAZINON AG500

For control of certain insects on fruits, nuts, vegetables, field crops, lawns and ornamentals. All Pro DIAZINON AG500 contains 4 lbs. diazinon per gallon.

ACTIVE INGREDIENTS:	Diazinon [0,0-dioethyl 0-(2-isopropyl-6-methyl-4-pyrimidinyl) phosphorothioate]	48%
INERT INGREDIENTS:		52%
TOTAL		100%

contains xylene range aromatic solvent

KEEP OUT OF REACH OF CHILDREN CAUTION

SEE ADDITIONAL PRECAUTIONARY STATEMENTS AND DIRECTIONS FOR USE INSIDE BOOKLET

IMPORTANT: Read the entire Directions for Use and Conditions of Sale and Warranty before using this product.

EPA Reg. No. 769-689

EPA Est. No. 769-GA-1

SURECO
FORT VALLEY, GA 31030

STATEMENT OF PRACTICAL TREATMENT

This is an organophosphate insecticide

- If swallowed: Call a physician or Poison Control Center immediately. DO NOT INDUCE VOMITING. Vomiting may cause aspiration pneumonia. Do not give anything by mouth if person is unconscious or convulsing.
- If inhaled, remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.
- If on skin, wash promptly with soap and water. Rinse thoroughly. Get medical attention.
- If in eyes, rinse eyes with plenty of water. Call a physician if irritation persists.
- Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. This product contains a xylene range aromatic solvent which may present an aspiration hazard. Gastric lavage may be indicated if product was taken internally.

PRECAUTIONARY STATEMENTS Hazards To Humans and Domestic Animals CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Causes moderate eye injury. Avoid contact with skin, eyes, or clothing. Avoid breathing vapor or spray mist. Avoid contamination of food and feed. Food utensils such as tablespoons and measuring cups should not be used for food purposes after use in measuring pesticides. Keep out of reach of domestic animals. Do not use on humans, household pets or livestock. Do not contaminate ornamental fish ponds.

Personal Protective Equipment: Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants.
- Chemical-resistant gloves, such as Barrier Laminate or Viton.
- Shoes plus socks.

Follow manufacturer's instructions for cleaning, maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

Users should:	User Safety Recommendations
• Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.	
• Remove clothing immediately if pesticide gets inside. Then wash thoroughly and change into clean clothing.	
• Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.	

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product.

DISCLAIMER: SureCo warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions for Use, subject to the inherent risks referred to above. SureCo makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use or handling or application of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited.

Pesticide: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If pesticide, spray mixture or rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance.

Container: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate, or burn if allowed by state and local authorities. Stay out of smoke from burning container.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking in treated areas may be killed. Because of the susceptibility of many birds to diazinon, it is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking in treated areas may be killed.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Agricultural Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard. Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 24 hours.

Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as Barrier Laminate or Viton
- Shoes plus socks

Note: The California Department of Agriculture has set a 5-day reentry period for diazinon on grapes, peaches, and nectarines. Five days should elapse between the time of application and the time a worker may enter the field to engage in any activity requiring substantial contact with treated foliage. When a mixture of two or more organophosphate pesticides are applied in combination, the interval should be prolonged by adding to the largest applicable interval an additional 50% of that interval.

NON-AGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses. Do not permit children and pets to go in treated areas until sprays have dried.

FAILURE TO FOLLOW ALL PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

CHEMIGATION

APPLICATION THROUGH AN IRRIGATION SYSTEM (CHEMIGATION) - CRANBERRIES ONLY
All Pro Diazinon AG500 may be applied alone or in combination with other pesticides registered for application through sprinkler irrigation systems to cranberries. To insure compatibility, pour the products into a small container of water in the correct proportions. After thorough mixing, let stand for five minutes. If the combination remains mixed, or can be remixed readily, mixture is compatible.

Do not apply this product through any type of irrigation system to other crops. For chemigation, apply this product to cranberries only through sprinkler irrigation systems. Do not apply this product to cranberries through any other type of irrigation system.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water. If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect the irrigation system used for pesticide application to a public water system unless the pesticide label-prescribed safety devices for public water systems are in place. A person knowledgeable of the chemigation system and responsible for its operation, or under the supervision of the responsible person, shall shut the system down and make necessary adjustments if the need arises.

Chemigation System Connected to Public Water Systems

1. Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.
2. Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

Sprinkler Chemigation

1. The system must contain a functional check valve, a vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.
2. The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.
3. The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.
4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Operating Instructions

1. Determine the acreage to be treated by the sprinkler system.
2. Measure the appropriate amount of All Pro Diazinon AG500 needed to treat the acreage to be treated. Refer to the specific directions for use for cranberries for the application rate and the amount of water per acre.
3. Add the premeasured All Pro Diazinon AG500 to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if All Pro Diazinon AG500 is tank mixed with other pesticides registered for sprinkler chemigation application to cranberries.
4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.

Fruit and

All Pro Di...
the table i...
use for ea...
methods (...
applied to...
ground ec...
appears i...
AG500 in...
ground sp...
used per...
Timing/St...
Observe t...
applicatio...

Crop

Almonds*
Aoncos
Blackber...
Boysenbe...
Cherries
Cranberri...
Dewberri...
Grapes
Loganber...
Nectarine...
Peaches
Plums
Prunes
Raspberri...
Strawberri...
Walnuts*

*CA on

Cy

Almonds
Aoncos
Blackber...
Boysenbe...
Cherries
Cranberri...
Dewberri...
Grapes
Loganber...
Nectarine...
Peaches
Plums, P...
Raspberri...
Strawberri...
Walnuts

Fruit Cit

Crop
Almonds

Aoncos

Notes: (1...
product (...
applicatio...

Cranberri...
(CA, OR...
WA only)
(Blackbe...
Boysenbe...
Dewberri...
Loganber...
Raspberri...

Cherries

...warranty of fitness or merchantability or any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use or handling or application of this product.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. Open dumping is prohibited. Pesticide: Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of federal law. If pesticide, spray mixture or rinsate cannot be disposed of by use according to label instructions, contact your State Pesticide or Environmental Control Agency, or the Hazardous Waste representative at the nearest EPA Regional Office for guidance. Container: Triple rinse (or equivalent). Then puncture and dispose of in a sanitary landfill, or incinerate, or, burn if allowed by state and local authorities. Stay out of smoke from burning container. For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup and disposal of wastes.

ENVIRONMENTAL HAZARDS

This product is highly toxic to birds, fish and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates. Rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. If waterfowl, i.e., ducks or geese, can be expected in the treated area after treatment (except spot treatment), water lawn/area with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes and estuaries. Do not apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters. This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

PHYSICAL OR CHEMICAL HAZARDS

Do not use, pour, spill, or store near heat or open flame.

GENERAL INFORMATION

All Pro Diazinon AG500 is an emulsifiable solution which, when diluted with water according to the directions, is intended for the control of insect pests of agricultural crops and other plants. Do not use on food crops grown in greenhouses. Do not use in barns or dwellings. Do not use on livestock. In the directions for use which follow, best control is obtained when application is made at first sign of pests. Application should be repeated only as directed to maintain control. To avoid spray drift, do not apply under windy conditions. Avoid spray overlap, since crop or plant injury may result.

- 4. The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.
- 5. The irrigation line or water pump must include a functional pressure switch which will stop the water pump motor when the water pressure decreases to the point where pesticide distribution is adversely affected.
- 6. Systems must use a metering pump, such as a positive displacement injection pump (e.g., diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.
- 7. Do not apply when wind speed favors drift beyond the area intended for treatment.

Operating Instructions

- 1. Determine the acreage to be treated by the sprinkler system.
- 2. Measure the appropriate amount of All Pro Diazinon AG500 needed to treat the acreage to be treated. Refer to the specific directions for use for cranberries for the application rate and the amount of water per acre.
- 3. Add the premeasured All Pro Diazinon AG500 to the chemigation system supply tank and dilute with water if needed. Proper agitation is needed if All Pro Diazinon AG500 is tank mixed with other pesticides registered for sprinkler chemigation application to cranberries.
- 4. Start the irrigation system and bring the system up to full pressure, making certain that all sprinkler heads are functioning properly.
- 5. Engage the chemigation injection or venturi system to add the diazinon to the irrigation lines.
- 6. Continue to run irrigation system for a few minutes after supply tank is empty to allow material in the irrigation lines to reach the sprinkler heads furthest from the chemigation injection point.

Work Safety Rules

REPEATED EXPOSURES TO CHOLINESTERASE INHIBITORS SUCH AS ARE CONTAINED IN THIS PRODUCT MAY, WITHOUT WARNING, CAUSE PROLONGED SUSCEPTIBILITY TO VERY SMALL DOSES OF ANY CHOLINESTERASE INHIBITOR. When handling All Pro Diazinon AG500 do not rub eyes or mouth with hands. If you feel sick in any way, STOP work and get help right away. See Practical Treatment section of this label. When handling this product, wear the personal protective equipment (PPE) as listed under personal protective equipment (PPE) paragraph on this label.

Boysen
Dewberry
Loganbe
Rausper

9/17

Chem:

Notes:
per acre
of 21 a
Crop
Cranbe

Notes: 1
for acre
Grape:

Notes
acre 0

9/17

All Pro Diazinon AG500 may be applied using ground or aerial application equipment as follows in the table below. The specific amount of product to be used appears under separate directions for use for each crop. This table indicates the minimum amount of water that can be used for the various methods of application. The rate of All Pro Diazinon AG500 for aerial spray or concentrated spray applied to fruit trees by ground equipment is based on a dilute full cover spray applied with conventional ground equipment at 300-400 gallons of water per acre. The specific amount of product to be used appears under the separate directions for use for each crop. If the rate calls for 1 pt. of All Pro Diazinon AG500 in 100 gallons of water, this amounts to 3-4 pints of product per acre when applied at dilute ground spray volumes, of 300-400 gallons of water per acre. In no case should the amount of product used per acre exceed the maximum amount per acre per application that is specified in the Timing/Special Directions for individual crops.

Observe stated time intervals between last application and harvest, as well as intervals between applications and total amount of product that can be made per season:

Crop	Air	Minimum Gallons/Acre	Minimum Gallons/Acre	
			Ground	Dilute Concentrate
Almonds*	Yes	20	Yes 100	20
Apricots	Yes	20	Yes 100	20
Blackberries**	Yes	20	Yes 100	20
Boysenberries**	Yes	20	Yes 100	20
Cherries	Yes	20	Yes 100	20
Cranberries	Yes	20	Yes 15	—
Dewberries**	Yes	20	Yes 100	20
Grapes	Yes	20	Yes 100	20
Loganberries**	Yes	20	Yes 100	20
Nectarines	Yes	20	Yes 100	20
Peaches	Yes	20	Yes 100	20
Plums	Yes	20	Yes 100	20
Prunes	Yes	20	Yes 100	20
Raspberries**	Yes	20	Yes 100	20
Strawberries	Yes	20	Yes 100	20
Walnuts*	Yes	20	Yes 100	20

*CA only **CA, OR, or WA only.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Boysenberries	7
Cherries	21
Cranberries	7
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums, Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

*PHI = Pre-harvest interval, or the number of days between the last application and harvest, may vary based on use pattern selected - see specific directions for use for grapes.
 ** Dormant spray only.
 Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds in the orchard are in bloom.

Nectarines	Oriental Fruit Moths	1 pt. in 100 gals. of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight.
	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers, Peach Twig Borers	1 pt. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days if necessary.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Peaches	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 pt. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days if necessary.
	Oriental Fruit Moths	1 pt. in 100 gals. of water	Apply at 100% petal fall. Make 2 additional applications at 10-15 day intervals. Time the last application to peak adult flight.
	Peach Twig Borers	1 pt. in 100 gals. of water	Apply as insects occur in May and June. Allow 7 days between applications.
	Leafhoppers	1 pt. in 100 gals. of water	Make 3-4 applications beginning at petal fall and continuing through the growing season. Allow at least 7 days between applications.
	White Peach Scale	1 1/2-2 pts. in 100 gals. of water	Apply post-harvest timed to coincide with peak crawler and immature scale activity. Under conditions of heavy infestation 2 applications (7 days apart) may be needed. Do not apply before fruit is harvested.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Plums	Brown Mites (Clover Mites), European Red Mites	1 pt. in 100 gals. of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	1/2-1 pt. in 100 gals. of water	

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Prunes	Brown Mites (Clover Mites), European Red Mites	1 pt. in 100 gals. of water	Apply as needed every 7 days.
	Leafcurl Plum Aphids, Mealy Plum Aphids, Thistle Aphids	1/2-1 pt. in 100 gals. of water	

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) Allow a minimum of 21 days between the last application and harvest.

Fruit Crops

Crop	Pest	Rate	Timing/Special Directions
Almonds (CA only)	San Jose Scale, Parlatoria Scale	1-1 1/2 pts. + 2-3 gals. dormant oil or 1-1 1/2 gals. superior type oil in 100 gals. of water	Apply as a dormant spray. Do not apply more than 6 pts. of diazinon or 9 gals. of oil per acre.
Apricots	Aphids, Brown Mites (Clover Mites), Two-spotted Spider Mites, Olive Scale Crawlers, San Jose Scale Crawlers	1 pt. in 100 gals. of water	Apply as infestations occur. Repeat application after 14 days if necessary.
	Olive Scale Crawlers	1/2 pt. + 1 1/2 gals. of light medium horticultural oil per 100 gals. of water	Apply when scale crawlers are present.
	Apricot Mealybugs	1 pt. per 100 gals. of water	Apply as a cover spray from petal fall to June. Allow 14 days between applications.

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Strawberries	Aphids, Two-spotted Spider Mites	1 pt. in 100 gals. of water per acre	Apply as pests occur. Repeat application if necessary every 7 days, but do not make more than 3 applications per season. Do not apply within 5 days of harvest or illegal residues may occur.
	Cyclamen Mites	1 qt. in 100 gals. of water per acre	Direct spray to the plant crowns. Plants should be sufficiently agitated to assure thorough coverage of the foliage. Repeat application after 7 days if necessary. Do not make more than 3 foliar applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Application may be made to plants after harvest.
	Mole Crickets	1 qt. per acre	Broadcast in sufficient water to obtain even coverage when insects are present. Apply 1-2 days before transplanting and immediately incorporate into the top 1-2 inches of soil.
	Strawberry Leafrollers	3/4-1 pt. in 100 gals. of water	Apply 200 gals. of spray per acre when blossoms show color. Do not apply within 5 days of harvest or illegal residues may occur.
Walnuts (CA only)	Aphids, Spider Mites	1 qt. per acre	Apply as a foliar spray. Repeat application in 14 days if necessary. Do not make more than 3 applications per season. Do not apply within 45 days of harvest or after husks open or illegal residues may occur. Make ground applications in a minimum of 100 gals. of water per acre.
	Codling Moths, Scale Crawlers, Walnut Caterpillars	3 qts. per acre	

Vegetable Crops

For foliar or soil applications, All Pro Diazinon AG500 should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. All Pro Diazinon AG500 may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application		Aerial Application Minimum Gals./Acre
	Minimum Gals./Acre	OR Minimum Gals./1,000 Sq. Ft.	
Beans (Lima, Pole, and Snap) (succulent only)	10	2	5
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabas	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5

Crop	Pest	Rate	Timing/Special Directions
Caneberries (CA, OR and WA only) (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries)	Leafhoppers, Thrips, Aphids, Two-spotted Spider Mites, Raspberry Fruitworms, Raspberry Sawflies	1 qt. per acre	Apply as insects occur. Repeat application after 14 days if necessary. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). For Raspberry Fruitworms make one application when blossom buds separate. Make a second application immediately before blossoms open.
	Dryberry Mites	1-2 qts. per acre	Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed but still closed. Apply in a minimum of 100 gals. of water per acre. Do not apply within 7 days of harvest or illegal residues may occur.
	Raspberry Crown Borers (Raspberry Root Borers)	2 qts. per acre	Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the Spring before buds break.
Cherries	Eyespotted Bud Moths, Fruit Tree Leafrollers	1 pt. in 100 gals. of water	Apply in cover sprays as infestations occur beginning at petal fall. Repeat application after 7 days if necessary.
	Leafhoppers	1 pt. in 100 gals. of water	Make 3-4 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications.

Pre-Harv Crop
 Beans (Lima, Pole, and Snap)
 Beets, Red (Table)
 Broccoli
 Brussels
 Cabbage
 Cantalou
 Carrots
 Casabas
 Cauliflow
 Chinese
 Chinese Broccoli
 Chinese Cabbage
 Chinese Mustard
 Collards
 Crenshaws
 Cucumbr
 Endive
 Ginseng
 Honeyd
 Kale
 Lettuce
 Muskm
 Mustard
 Onions
 Parsn
 Peas
 Pepper
 Parsn
 Potato
 Radish
 Spinac
 Squash
 Squash
 Sweet
 Swiss
 Tomak
 Turnip
 Water
 *PHI
 Notes: immed using incorp applic area a
 Crop
 Bean, Pole, (succ)

Notes: (1) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (2) To avoid illegal residues, allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Canberries (CA, OR and WA only) (Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries)	Leafhoppers, Thrips, Aphids, Two-spotted Spider Mites, Raspberry Fruitworms, Raspberry Sawflies	1 qt. per acre	Apply as insects occur. Repeat application after 14 days if necessary. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Apply in a minimum of 100 gals. of water per acre (200 gals. per acre maximum). For Raspberry Fruitworms make one application when blossom buds separate. Make a second application immediately before blossoms open.
	Dryberry Mites	1-2 qts. per acre	Apply as a dormant spray in the early Spring when canes are still on the ground and buds are closed. Make a second application when buds are well developed but still closed. Apply in a minimum of 100 gals. of water per acre. Do not apply within 7 days of harvest or illegal residues may occur.
	Raspberry Crown Borers (Raspberry Root Borers)	2 qts. per acre	Apply a single application in a minimum of 100 gals. of water per acre as a drench to the crown and lower canes in the Spring before buds break.
Cherries	Eye-spotted Bud Moths, Fruit Tree Leafrollers	1 pt. in 100 gals. of water	Apply in cover sprays as infestations occur beginning at petal fall. Repeat application after 7 days if necessary.
	Leafhoppers	1 pt. in 100 gals. of water	Make 3-4 applications beginning at petal fall and continuing through the growing season as infestations occur. Allow 7 days between applications.
	Cherry Fruitflies	1/2-1 pt. in 100 gals. of water	Make 3-4 applications beginning when adult flies begin to emerge and repeating at 10 day intervals.
	Black Cherry Aphids	1 pt. in 100 gals. of water	Apply as a cover spray when aphids appear. Repeat application after 10 days if necessary.
	San Jose Scale Crawlers	1 pt. in 100 gals. of water	Apply during the pre-pink stage or when crawlers first appear.
	Cherry Rust Mites	1/2-1 pt. in 100 gals. of water	Apply to trees as a single supplemental spray after harvest.

Notes: (1) Do not make more than 3 applications per season. (2) Do not apply more than 4 pts. of product per acre per application and no more than 12 pts. of product per acre per season. (3) Allow a minimum of 21 days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Cranberries	Blackheaded Fireworms (<i>Rhopobota neevana</i>)	2 qts. in a minimum of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation.	Apply when pest appears. Repeat application after 14 days if necessary. A maximum of 6 applications per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur.
	Cranberry Fruitworms	3 qts. in a minimum of 15 gals. of water per acre for ground application and up to 400 gals. of water per acre for chemigation.	Apply when pest appears. Repeat application after 14 days if necessary. A maximum of 4 applications per season is permitted. Do not apply within 7 days of harvest or illegal residues may occur.

Note: This product may be applied to cranberries through sprinker irrigation. Refer to the Chemigation section for specific chemigation directions for use.

Crop	Pest	Rate	Timing/Special Directions
Grapes	Pacific Spider Mites, Leafhoppers, Grape Berry Moths, Grape Leafhoppers, Omnivorous Leafhoppers, Aphids, Spider Mites	1-2 pts. per acre in a minimum of 100 gals. of water	Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary.
	<i>Drosophila</i> spp.	8 fl. oz. per acre in a minimum of 100 gals. of water (300 gals. of water maximum)	Apply as pest occurs as a thorough cover spray. Repeat application after 7 days, if necessary.

Notes: (1) Do not apply more than 2 pts. of product per acre per application and no more than 10 pts. per acre per season. (2) To avoid illegal residues, allow 28 days between the last application and harvest.

Vegetable Crops

For foliar or soil applications, All Pro Diazinon AG500 should always be applied as a spray in sufficient water to assure thorough coverage of the foliage or soil. All Pro Diazinon AG500 may be applied using ground or aerial application equipment as specified in the following table. This table indicates the minimum amount of water that can be used for the various methods of application. The specific amount of product to be used appears under the separate directions for use for each crop.

Crop	Ground Application		Aerial Application Minimum Gals./Acre
	Minimum Gals./Acre	OR Minimum Gals./1,000 Sq. Ft.	
Beans (Lima, Pole, and Snap) (succulent only)	10	2	5
Beets, Red (Table)	10	2	5
Broccoli	10	2	5
Brussels Sprouts	10	2	5
Cabbage	10	2	5
Cantaloupes	5	1	5
Carrots	10	2	5
Casabes	5	1	5
Cauliflower	10	2	5
Chinese Broccoli	10	2	5
Chinese Cabbage	10	2	5
Chinese Mustard	10	2	5
Chinese Radish (CA, FL only)	10	2	5
Collards	10	2	5
Crenshaws	5	1	5
Cucumbers	5	1	5
Endive (Escarole)	10	2	5
Ginseng	10	2	5
Honeydew Melons	5	1	5
Kale	10	2	5
Lettuce, Head	10	2	5
Lettuce, Leaf	10	2	5
Muskmelons	5	1	5
Mustard	10	2	5
Onions (Bulb and Green)	10	2	5
Parsley	10	2	5
Parsnips	10	2	5
Peas (succulent only)	10	2	5
Peppers	10	2	5
Persian Melons	5	1	5
*Potatoes (Irish)	10	2	5
Radishes	10	2	5
Spinach	10	2	5
Squash, Summer	5	1	5
Squash, Winter	5	1	5
Sweet Corn	20	4	5
Sweet Potatoes	10	2	5
Swiss Chard	10	2	5
Tomatoes	10	2	5
Turnips	10	2	5
Watermelons	5	1	5

*Do not use on commercially grown potatoes which will be hand-harvested.

Observe stated time intervals between last application and harvest as well as intervals between and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI)* for Foliar Applications

Crop	PHI* (Days)
Beans (Lima, Pole, and Snap)	7
Beets, Red (Table)	14
Broccoli	7
Brussels Sprouts	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli	10
Chinese Cabbage	10
Chinese Mustard	10
Chinese Radish	10
Collards	10
Cranberries	3
Cucumbers	7
Endive (Escarole)	14
Ginseng	30
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Mustard	10
Onions (Bulb and Green)	14
Paranips	14
Peas	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Rapishes	14
Sprach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Swiss Chard	14
Tomatoes	1
Tumpe, Roots and Tops	14
Watermelons	3

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

Notes: (1) Soil Incorporation: Following application of this product to control vegetable soil insects, immediately incorporate the product into the soil to the recommended depth (see Timing/Special Directions) using a rotary hoe, cultivator, disk, harrow, or other suitable means. For control of surface cutworms, incorporate 2-3 inches, for subterranean cutworms incorporate 3-6 inches. (2) To protect bees: Foliar applications of this product should not be made to peas or beans if these crops or weeds in the treatment area are in bloom, or to corn during the pollen shed period.

Crop	Pest	Rate	Timing/Special Directions
Beans (Lima, Pole, and Snap) (succulent only)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Black Bean Aphids, Mexican Bean Beetles, Dipterous Leafminers	1-1½ pts. per acre	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal residues may occur.
	Cucumber Beetles	¾ pt. per acre	Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. Make cucumber beetle application in 100 gals. of water per acre. Notes: (1) Diazinon will not control organophosphate-resistant leafminers. (2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
Beans, Lima (succulent only)	Leafhoppers, Spider Mites	1 pt. per acre	Apply as pests occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Bean forage or bean hay may be fed to dairy and beef cattle and sheep. Wait 4 days following treatment if forage is to be cut for hay. Note: To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
Beets, Red (Table)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur.

Vegetable	Pests	Rate	Application	Vegetable	Pests	Rate	Application	Notes	
Brussels Sprouts	Root Maggots	2-3 qts. per acre	Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.	Endive (Escarole)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	Potatoes	
		4-8 oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.		Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.		
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)		Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.		Aphids, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.		
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		Ginseng	Leafhoppers, Aphids, Lygus Bugs, Flea Beetles, Jumping Plant Lice	1/2-1 pt. per acre		Spray when insects first appear. For heavy insect infestation, use 1 qt. per acre. Apply by ground equipment using 10 to 100 gals. of water per acre. Do not apply more than one application per growing season. Do not apply during flowering of three or four year old crops. Do not apply within 30 days of harvest. Do not graze treated areas or feed treated forage to livestock.
	Aphids, Diamondback Moths, Imported Cabbageworms	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur.			Kale	Cutworms		2-4 qts. per acre
Cabbage	Root Maggots	2-3 qts. per acre	Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.	Lettuce	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	Spinach	
		4-8 oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.		Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.		
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)		Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.		Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin Bugs, Corn Earworms	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.		
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		Mustard	Cutworms	2-4 qts. per acre		Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 21 days of harvest, or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.			Wireworms	3-4 qts. per acre		Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Carrots	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	Melons (Cantaloupes, Casabas, Crenshaws, Honeydews, Muskmelons, Persians, and Hybrids of these) and Watermelons	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	Squash	
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.		Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		Aphids, Dipterous Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.		
	Aphids	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest, or illegal residues may occur.		Onions (Bulb and Green)	Onion Maggots	2-4 qts. per acre		Broadcast just before planting and mix into the top 3-4 inches of soil. Note: Diazinon will not control organophosphate-resistant onion maggots.
Cauliflower	Root Maggots	2-3 qts. per acre	Apply to plant beds or fields just before sowing of seed or transplanting and immediately incorporate into the top 3-4 inches of soil.	Onion Thrips		1 pt. per acre	Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.		
		4-8 oz. in 50 gals. of water	In transplant water as a drench application when 200-300 gals. of water are used per acre. Apply 1/2-1 cup (4 to 8 oz.) per plant by hand or by tractor-mounted sprayer equipped with drop nozzles to direct spray to the base of the plant. Note: Transplant water treatments may result in stand reduction due to plant stress at time of transplanting.	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.			
	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)	Parsley	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)		
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.		Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.		
Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.	Parsnips	Aphids, Flea Beetles, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.			
Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.		Peas (succulent only)	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)		
Chinese Broccoli	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.		Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.	Sweet Pea	
				Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.		Wireworms
Chinese Mustard (Gai Choy)	Aphids, Diamondback Moths	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.					Aphids	1/2-1 pt. per acre

Note: Do

Radishes

Spinach

Squash

Squash

Sweet Ct

Notes: (1) the last ac (3) Corn m

Sweet Pea

Swiss Ch.

Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Aphids, Diamondback Moths, Imported Cabbageworms, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days, with a minimum of 7 days between applications. Do not apply within 5 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Broccoli	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Cabbage (Bok Choy and Napa)	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Mustard (Gai Choy)	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Chinese Radish (Japanese Radish or Daikon) (FL and CA only)	1 pt. per acre	Mix in a minimum of 100 gals. of water per acre and apply as a foliar spray as pests occur. Repeat application as necessary, every 14 days. Do not make more than 3 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Collards	Cutworms	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Diamondback Moths, Imported Cabbageworms, Harlequin Bugs, Corn Earworms	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 10 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Cucumbers	Cutworms	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Onion Thrips	1 pt. per acre	Apply as thrips appear. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.
Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Parsley	Cutworms	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Parsnips	Aphids, Flea Beetles, Dipterous Leafminers	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
	Cutworms	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
Peas (succulent only)	Cutworms	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Peppers	Aphids, Dipterous Leafminers	Apply as insects occur. Repeat application, as necessary, every 5 days. Do not make more than 3 applications per season. To avoid illegal residues, allow a minimum of 7 days between the last application and harvest. Vines and hay may be fed to dairy and beef cattle and sheep. To avoid illegal residue, wait 7 days following treatment if forage is to be cut for hay. Notes: Diazinon will not control organophosphate-resistant leafminers. (2) To protect bees, do not apply foliarly if the crop or weeds in the treatment area are in bloom.
	Cutworms	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 5 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Notes: (1) the last ac (3) Corn fr Sweet Po

Swiss Cr

Tomatoes

Turnips and Top

Vegetable Crops continued

Potatoes	Cutworms (Surface and Subterranean)	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Southern Potato Wireworms (Southeastern U.S.)	2-4 qts. per acre	For potatoes grown on mineral soils in the Southeast, apply before planting and immediately incorporate into the top 3-6 inches of soil depending on location of wireworms.
	Aphids, Flea Beetles, Colorado Potato Beetles, Dipterous leafminers	1/2-3/4 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 35 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
Banded Cucumber Beetle, Leafhoppers, Southern Armyworms	3/4-1 pt. per acre	Apply as insects occur. Repeat application as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 35 days of harvest or illegal residues may occur.	

Note: Do not use on commercially grown potatoes which will be hand-harvested.

Radishes	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 3 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Spinach	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.

Squash, Summer	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	1/2-1 1/2 pts. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 7 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Squash, Winter	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Serpentine Leafminers, Thrips, Striped Cucumber Beetles, Melonworms, Leafhoppers, Spider Mites	1/2-1 1/2 pts. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 3 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Sweet Corn	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	See Corn Maggots	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast before planting and immediately incorporate into the top 4-8 inches of soil.
	Corn Earworms	1-1 1/4 qts. per acre	Apply when silks first appear. Repeat application every 7 days if necessary. Do not make more than 5 applications per season.
	Corn Rootworm (Adults)	1/2-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Corn Leaf Aphids	1-2 pts. per acre	Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Grasshoppers, Spider Mites, Flea Beetles	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Sap Beetles	2-2 1/2 pts. per acre	Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Notes: (1) To protect bees, do not apply liberally to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be pickled 7 days after the last application.

Sweet Potatoes	Wireworms, Flea Beetles	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Swiss Chard	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)

Field Crops

Soil and foliar applications of All Pro Diazinon AG500 should always be applied as a spray in sufficient water to ensure thorough coverage. For foliar applications, observe stated time intervals between last application and harvest, as well as intervals between applications and total number of applications that can be made per season.

Pre-Harvest Intervals (PHI)* for Foliar Applications

Crop	PHI* (Days)
Hops	14
Sugar Beets	14

*PHI = Pre-Harvest Interval or the number of days between the last application and harvest.

Crop	Pest	Rate	Timing/Special Directions
Hops	Aphids, Spider Mites	1 qt. per acre	Apply as pests occur. Repeat application, as necessary, every 14 days. Do not make more than 4 applications per season. Do not apply within 14 days of harvest.
	Wireworms	3-4 qts. per acre	Broadcast just before planting. Incorporate into soil 4-6 inches. Do not apply more than 4 qts. of this product to the soil, per season.
Sugar Beets	Beet Leafhoppers, Dipterous Leafminers, Aphids	3/4-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest. Note: Diazinon will not control organophosphate-resistant leafminers.
	Grasshoppers	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest.

Note: Tops may be fed to beef and dairy animals.

Grassland Insects

Ditch Banks, Roadside, Wasteland, Noncrop Areas, Barrier Strips
Grasshoppers: Spray 1/4-1 pt. per acre in water (minimum of 1 gal. of water per acre) for aerial applications or a minimum of 5 gals. of water per acre for ground applications or in oil (a minimum of 1 gal. of oil per acre) when insects first appear, preferably in the nymphal stage. Thorough coverage of the foliage is essential.

Do not graze or use treated grass for forage or feed.

Insect Control on Ornamentals

To control certain insects on Arborvitae, Azalea, Birch, Boxwood, Camellia, Carnation, Chrysanthemum, Douglas Fir, Elm, Gladiolus, Hawthorn, Holly, Juniper, Lilac, Locust, Maple, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Rose, Spruce and Willow, apply the recommended rates indicated below. Try to spray underside of leaves and penetrate dense foliage.

Insects	Amount of All Pro Diazinon AG500 to Use	
	Rate Per 3 Gals. Water	Rate Per 100 Gals. Water
Aphids, Bagworms, Carnation Bud Mites, Carnation Shoot Mites, Clover Mites, Cyclamen Mites, Dipterous Leafminers, European Pine Shoot Moths, European Red Mites, Flea Beetles, Holly Bud Moths, Leafhoppers, Obocure Root Weevil, Omnivorous Leafhoppers, Privet Mites, Scale Crawlers (Cottony-cushion, Lecanium, Pine Needle, San Jose, Soft), Thrips, Two-spotted Spider Mites, Juniper Webworms, Whiteflies	1/2 fl. oz.*	1 pt.**
Apple-and-Thorn Skeletonizers, Cottonweaver Webworms, Fall Webworms, Flecklock Chermes, Oak Looper, Oblongbanded Leafhoppers, Pear Slugs, Tent Caterpillars	1 1/2 fl. oz.*	3 pts.**

*1 fl. oz. = 2 tablespoons
**16 fl. oz. = 1 pint

Precaution: Do not use on Ferns, Poinsettia, Hibiscus, Peppers, Piles, and Gardenia, since injury to the plants may occur. Mimosa Webworms: Mix 1/2 fl. oz. in 3 gals. of water (or 1 pt. in 100 gals. of water) and apply as a thorough foliar spray when adults first appear. Make a second and third application at 4-5 week intervals.

Lawn Pest Control (Excluding Golf Courses and Sod Farms)

Precaution: Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label. Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York between November 1 and May 20. Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds, especially waterfowl. Avoid overlapping sprays. If waterfowl, i.e., ducks or geese, can be expected to enter the treated area after treatment (except spot treatments), water lawn with at least 1/2 inch water immediately after application, however, stop watering before puddling occurs.

To control pests listed below, apply recommended rates.

Pests	Rate per 1,000 sq. ft.		Remarks
	All Pro Diazinon AG500	Water	
Lawn Chinch Bugs	2-3 fl. oz.	3 gals.	Use higher rate for longer residual control and in lawns of dense growth such as St. Augustine lawns.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Crickets, Cutworms, Digger Wasps, Earwigs, Fire Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs), Springtails (Collembola)	2-3 fl. oz.	3 gals.	Spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk when wasps are not active. For fire flies, mow grass and water well before treatment, but delay application until grass is dry. For billbug control, apply when activity is first observed or when chewed or brown grass is present, indicating damage from this insect.
Brown Dog Ticks, Chiggers, Fleas	3 fl. oz.	3 gals.	Note: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevils	3 fl. oz.	3 gals.	Treat problem areas in mid-April and again in mid-May
Millipedes	3 fl. oz.	3 gals.	
Rhodesgrass Scales	3 fl. oz.	3 gals.	Apply when crawlers first emerge.
White Grubs (such as Japanese Beetle Larvae, European Chafar, Southern Chafar)	3 fl. oz.	3 gals.	Apply any time between late July and early October.

Nuisance Pests in Outdoor Areas

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pill Bugs), Spiders, Springtails (Collembola): Mix 3 fluid ounces in 3 gallons of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls and entranceways, walkways, garbage cans, fire trucks, and into cracks and other places where insects can hide. To prevent entry into houses spray a five foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary. Do not apply to animals. Do not treat plants (other than tree trunks) at this nuisance pest rate.

Corn Earworms	1-1½ qts. per acre	Apply when silks first appear. Repeat application every 7 days if necessary. Do not make more than 5 applications per season.
Corn Rootworm (Adults)	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Corn Leaf Aphids	1-2 pts. per acre	Apply in sufficient water to provide complete coverage to the upper portions of the plants. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Grasshoppers, Spider Mites, Flea Beetles	1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
Sap Beetles	2-2½ pts. per acre	Consult local agricultural authorities for proper timing of sprays. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.

Notes: (1) To protect bees, do not apply foliarly to corn during the pollen shed period. (2) Allow 7 days after the last application before grazing or cutting for forage. Forage may be fed to beef and dairy cattle and sheep. (3) Corn may be picked 7 days after the last application.

Sweet Potatoes	Wireworms, Flea Beetles	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
Swiss Chard	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Imported Cabbageworms, Diamondback Moths, Harlequin bugs	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur.
Tomatoes	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Dipterous Leafminers	½ pt. per acre	Apply as insects appear. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season.
	Banded Cucumber Beetles, Fall Armyworms, Southern Armyworms, Beet Armyworms	¾-1 pt. per acre	Do not apply within one day of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.
	Vinegar Flies (Drosophila spp.)	½-1½ pts. per acre	
Turnips, Roots and Tops	Cutworms	2-4 qts. per acre	Broadcast just before planting and immediately incorporate into the soil. (See Soil Incorporation Note.)
	Mole Crickets	1 qt. per acre	Apply to plant beds or fields 1-2 days before sowing of seed or transplanting and immediately incorporate into the top 1-2 inches of soil.
	Wireworms	3-4 qts. per acre	Broadcast just before planting and immediately incorporate into the top 4-8 inches of soil.
	Aphids, Flea Beetles, Dipterous Leafminers	½-1 pt. per acre	Apply as insects occur. Repeat application, as necessary, every 7 days. Do not make more than 5 applications per season. Do not apply within 14 days of harvest or illegal residues may occur. Note: Diazinon will not control organophosphate-resistant leafminers.

Brown Dog Ticks, Chiggers, Fleas	3 fl. oz.	3 gals.	grass is present, indicating damage from this insect.
Hyperoedae Weevils	3 fl. oz.	3 gals.	Note: Do not apply to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Millipedes	3 fl. oz.	3 gals.	These problem areas in mid-April and again in mid-May
Rhodesgrass Scales	3 fl. oz.	3 gals.	Apply when crawlers first emerge.
White Grubs (such as Japanese Beetle Larvae, European Chafar, Southern Chafar)	3 fl. oz.	3 gals.	Apply any time between late July and early October.

Nuisance Pests in Outside Areas

Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pill Bugs), Spiders, Springtails (Collembola): Mix 3 fluid ounces in 3 gallons of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls and entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.

To prevent entry into houses spray a five foot band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 feet. Repeat application as necessary. Do not apply to animals. Do not treat plants (other than tree trunks) at this nuisance pest rate.

AIIPRO

Manufactured by:
SureCo, Inc.
Fort Valley, Georgia 31030

16/17

COPY A

Do not use water from irrigated or flooded cranberry beds for drinking purposes or to irrigate crops other than those appearing on EPA approved diazinon labels.