

PM 19

769-835

Page 1 of 6
5515147 300
17

UNITED STATES ENVIRONMENTAL PROTECTION AGENCY

FEB 28 1997

Stacy A. Schmidt
SureCo
10012 N. Dale Mabry Hwy.
Suite 221
Tampa, FL 33618

Subject: Miller 1.75% Sevin® Dust
EPA Registration No. 769-835
Amendment dated November 12, 1996
Response to Agency letter dated 10/1/96

Dear Ms. Schmidt:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable provided that you:

- 1. Make the following changes to your label:
 - a) In the Precautionary Statements, delete the following sentences:
 - "Wash thoroughly after handling. Wear gloves and long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating or smoking."
 - "Keep children and pets out of treated area until dry."

These sentences are repeated in the required labeling in the User Safety Requirements box.

b) Move the 3 paragraphs that appear below the User Safety Requirements box "Wearing household latex or rubber gloves,....more than once a week" to the DIRECTIONS FOR USE section of your label as required in the Agency letter dated 10/1/96.

BEST AVAILABLE COPY

		CONCURRENCES					
SYMBOL							
SURNAME							
DATE							

Bedwards A: 769/835.1 2/26/97

- c) In the Directions for Use on CORN, delete the sentences "Repeat as necessary." and "Apply at 2 to 3 day intervals,..." The correct application interval is not more than once per week.
 - d) In the Directions for Use on LETTUCE; delete "5 to" in the last sentence.
 - e) In the Directions for Use on OKRA, delete "5 to" in the last sentence.
2. Submit one copy of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely yours,

/s/

Dennis H. Edwards, Jr.
 Product Manager (19)
 Insecticide-Rodenticide Branch
 Registration Division (7505C)

BEST AVAILABLE COPY

Enclosure

MILLER 1.75% SEVIN® DUST

AN INSECTICIDE
For Use On CERTAIN SMALL FRUITS
ORNAMENTALS and VEGETABLES

ACTIVE INGREDIENT:	
Carbaryl (1-Naphthyl Methylcarbamate)	1.75%
INERT INGREDIENTS:	98.25%
TOTAL:	100.00%

Sevin® is a registered trademark of Rhone Poulenc for carbaryl insecticide

EPA Reg. No. 769-835

EPA Est. No. 769-GA-1

KEEP OUT OF REACH OF CHILDREN CAUTION / CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Call a physician or poison control center immediately. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF IN EYES: Immediately flush eyes with plenty of water. Call a physician.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine administered intravenously is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured By:
SureCo, Inc.
Fort Valley, GA 31030


ACCEPTED
with COMMENTS
in EPA Letter Dated

Net Weight: _____

FEB 28 1997

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

769-835



PRECAUTIONARY STATEMENTS
Hazards To Humans and Domestic Animals
CAUTION

May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Wash thoroughly after handling. Wear gloves and long-sleeved work clothing. Change to clean clothing daily. Wash hands and face before eating or smoking. Do not contaminate food or feed. Avoid storage near food or feed products. Avoid contamination of feed, feeding troughs and watering receptacles. Food utensils such as teaspoons or tablespoons should not be used for food purposes after use in measuring pesticides. Do not use on humans or household pets. Keep children and pets out of treated area until dry.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. **MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.** Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application, or take equally effective precautions.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin.

Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

BEST AVAILABLE COPY

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in tightly closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

GENERAL INFORMATION

Sevin is an outstanding insecticide that controls a wide variety of insect pests. Sevin is not effective against mites; where mites are a problem, use a suitable miticide.

This product is compatible with commonly used miticides, fungicides and insecticides. Do not use with lime or other alkaline material.

FOR SMALL MEASUREMENTS

57 standard level tablespoonful = 1 lb.
 3 1/2 standard level tablespoonful = 1 oz.
 4 1/2 ounces = 1 cupful

ORNAMENTALS: Arborvitae, Azalea, Boxwood, Dogwood, Rose, Spirea, Yew: For Control of Japanese Beetle, Bagworm, Aphid, Flea Beetle, Leafhoppers, Tent Caterpillar and exposed Thrips: Apply 2 1/2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do not use on Boston Ivy or Virginia Creeper.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Sevin may be applied up to and including the day of harvest on Beans, Blueberries, Sweet Corn, Cucumbers, Okra, Summer Squash, Tomatoes, Eggplant, Peppers, Potatoes and within 1 day of harvest on Cranberries and Strawberries and within 5 days of harvest on Head Lettuce, or within 14 days on Leaf Lettuce. Allow 7 days between last application and harvest on Corn for use as forage and fodder as feed for dairy and meat animals. Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Sevin residues when Sevin is used in accordance with label directions. When Sevin has been applied in accordance with these directions for use, feeding of treated corn forage will not result in residues in milk or meat.

Apply this product when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results apply thoroughly to upper and lower leaf surfaces. If necessary, repeat applications every 7 to 10 days, or as indicated. Apply with conventional dusting equipment. Do not dust too heavily or crop injury may result.

BEANS: For control of Mexican Bean Beetle: Apply 2 lbs. per 1,000 sq. ft. For control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetles: Apply 1 1/2 lbs. per 1,000 sq. ft. For control of Corn Earworm, Armyworms and Velvet Bean Caterpillars: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 2 1.2 lbs. per 1,000 sq. ft.

BEST AVAILABLE COPY

CORN: For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetle: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. Repeat as necessary. For insects attacking Silks and Ears: Apply at 2 to 3 day intervals, starting when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential for effective control. May be applied up to harvest on Sweet Corn. Do not apply within 7 days of harvest on corn used for fodder or forage.

CUCUMBERS AND SUMMER SQUASH: For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs and Flea Beetles: Apply 1 1/2 lbs. per 1,000 sq. ft.

LETTUCE: For control of Armyworm, Cabbage Caterpillar, small Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus Bug and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 5 to 7 day intervals.

OKRA: For control of Corn Earworm and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 5 to 7 day intervals, or as necessary.

POTATOES, TOMATOES, EGGPLANTS AND PEPPERS: For control of Colorado Potato Beetle, Flea Beetles, and Leafhoppers: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus and Stink Bugs: Apply 2 1/2 lbs. per 1,000 sq. ft.

BLUEBERRIES: For control of Cherry and Cranberry Fruitworm: Apply 2 lbs. per 1,000 sq. ft.

CRANBERRIES: For control of Fruitworms, Fireworms, Leafhoppers and Cutworms: Apply 2 1/2 to 3 lbs. per 1,000 sq. ft.

STRAWBERRIES: For control of Meadow Spittlebug and Leaf Roller: Apply 2 1/2 lbs. per 1,000 sq. ft.

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product.
DISCLAIMER: SureCo warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. SureCo makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use, handling or application of this product.

BEST AVAILABLE COPY