

PM 04

769-835

3-3-98

10/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

MAR 3 1998

Stacey A. Schmidt  
SureCo  
7501 Harps Mill Road  
Raleigh, N.C. 27615-5422

Subject: Miller 1.75% Sevin Dust  
EPA Registration No. 769-835  
Deletion of Ornamentals  
Amendment dated March 20, 1997  
Federal Register (Vol. 62, No. 166) August 27, 1997

Dear Ms. Schmidt:

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, is acceptable provided that you:

1. Make the following changes to your label:

- a) The use for "ORNAMENTALS" was deleted from this label; however, immediately below the name of the product where it reads "AN INSECTICIDE For Use On CERTAIN SMALL FRUITS ORNAMENTALS and VEGETABLES" this use deletion was not taken into account. Remove the item "ORNAMENTALS" from this statement.
- b) In the Directions for Use on BEANS in the last sentence change the "Apply 2 1.2 lbs. per 1,000 sq. ft." to "Apply 2 1/2 lbs. per 1,000 sq. ft."
- c) In the Directions for Use on LETTUCE delete the word "small" from "small Cabbage Looper".

2. Submit two copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product bearing the amended labeling constitutes acceptance of these conditions.

A stamped copy of the labeling is enclosed for your records.

Sincerely,  
*George Tompkins*  
George Tompkins, Ph.D., Entomologist  
Insecticide-Rodenticide Branch  
Registration Division (7505C)

2075

ACCEPTED  
with COMMENTS  
in EPA Letter Dated:

# MILLER 1.75% SEVIN® DUST

AN INSECTICIDE  
For Use On CERTAIN SMALL FRUITS  
ORNAMENTALS and VEGETABLES

MAR 3 1998  
Under the Federal Insecticide,  
Fungicide, and Rodenticide Act  
as amended, for the pesticide  
registered under EPA Reg. No.

769-835

ACTIVE INGREDIENT:	
Carbaryl (1-Naphthyl Methylcarbamate) .....	1.75%
INERT INGREDIENTS: .....	98.25%
TOTAL:	100.00%

Sevin® is a registered trademark of Rhone Poulenc for carbaryl insecticide

EPA Reg. No. 769-835

EPA Est. No. 769-GA-1

## KEEP OUT OF REACH OF CHILDREN CAUTION / CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

### STATEMENT OF PRACTICAL TREATMENT

**IF SWALLOWED:** Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Call a physician or poison control center immediately. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

**IF INHALED:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

**IF IN EYES:** Immediately flush eyes with plenty of water. Call a physician.

**IF ON SKIN:** Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing.

**NOTE TO PHYSICIAN:** Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine administered intravenously is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured By:  
SureCo, Inc.  
Fort Valley, GA 31030

Net Weight: \_\_\_\_\_

» » »  
» » »  
» » »  
» » »  
» » »  
» » »  
» » »  
» » »  
» » »  
» » »



4075

### STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in tightly closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

### GENERAL INFORMATION

Sevin is an outstanding insecticide that controls a wide variety of insect pests. Sevin is not effective against mites; where mites are a problem, use a suitable miticide.

This product is compatible with commonly used miticides, fungicides and insecticides. Do not use with lime or other alkaline material.

### FOR SMALL MEASUREMENTS

- 57 standard level tablespoonful = 1 lb.
- 3 1/2 standard level tablespoonful = 1 oz.
- 4 1/2 ounces = 1 cupful

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

Sevin may be applied up to and including the day of harvest on Beans, Blueberries, Sweet Corn, Cucumbers, Okra, Summer Squash, Tomatoes, Eggplant, Peppers, Potatoes and within 1 day of harvest on Cranberries and Strawberries and within 5 days of harvest on Head Lettuce, or within 14 days on Leaf Lettuce. Allow 7 days between last application and harvest on Corn for use as forage and fodder as feed for dairy and meat animals. Tolerances established under the Federal Food, Drug and Cosmetic Act permit the sale of crops bearing probable Sevin residues when Sevin is used in accordance with label directions. When Sevin has been applied in accordance with these directions for use, feeding of treated corn forage will not result in residues in milk or meat.

Apply this product when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results apply thoroughly to upper and lower leaf surfaces. If necessary, repeat applications every 7 to 10 days, or as indicated. Apply with conventional dusting equipment. Do not dust too heavily or crop injury may result.

**BEANS:** For control of Mexican Bean Beetle: Apply 2 lbs. per 1,000 sq. ft. For control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetles: Apply 1 1/2 lbs. per 1,000 sq. ft. For control of Corn Earworm, Armyworms and Velvet Bean Caterpillars: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 2 1.2 lbs. per 1,000 sq. ft.

**CORN:** For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetle: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. For insects attacking Silks and Ears: Apply when first silks appear and continuing until silks begin to dry. Three or more applications may be required depending on severity of infestation. Timing and good coverage are absolutely essential.

