

PM 04

769-835

12/2/98

ACCEPTED 184

SEP 2

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the purpose of registration under EPA Reg. No. 769-835

MILLER 1.75% SEVIN® DUST

AN INSECTICIDE
 For Use On CERTAIN SMALL FRUITS
 ORNAMENTALS and VEGETABLES

ACTIVE INGREDIENT:

Carbaryl (1-Naphthyl Methylcarbamate)	1.75%
OTHER:	98.25%
TOTAL:	100.00%

Sevin® is a registered trademark of Rhone Poulenc for carbaryl insecticide

EPA Reg. No. 769-835

EPA Est. No. 769-GA-1

KEEP OUT OF REACH OF CHILDREN
CAUTION / CAUCIÓN

PRECAUCION AL USUARIO: Si usted no lee ingles, no use este producto hasta que la etiqueta haya sido explicado ampliamente.

FIRST AID

IF SWALLOWED: Drink one or two glasses of water and induce vomiting by touching back of the throat with finger. Call a physician or poison control center immediately. Do not induce vomiting or give anything by mouth to an unconscious or convulsing person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth.

IF IN EYES: Immediately flush eyes with plenty of water. Call a physician.

IF ON SKIN: Remove contaminated clothing. Wash affected area with soap and water. Launder clothes before reusing.

NOTE TO PHYSICIAN: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine administered intravenously is antidotal. Do not use 2-PAM, opiates, or cholinesterase inhibiting drugs.

SEE SIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

Manufactured By:
 SureCo, Inc.
 Fort Valley, GA 31030

Net Weight: _____

PRECAUTIONARY STATEMENTS
Hazards To Humans and Domestic Animals
CAUTION

May be harmful if swallowed. Avoid breathing vapors (dust or spray mist). Avoid contact with eyes, skin or clothing. Do not contaminate food or feed. Avoid storage near food or feed products. Avoid contamination of feed, feeding troughs and watering receptacles. Food utensils such as teaspoons or tablespoons should not be used for food purposes after use in measuring pesticides. Do not use on humans or household pets.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to aquatic and estuarine invertebrates. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. This product is highly toxic to bees exposed to direct treatment or residues on blooming crops or weeds. Do not apply this product or allow it to drift to blooming crops or weeds if bees are visiting the treatment area. **MAY KILL HONEYBEES IN SUBSTANTIAL NUMBERS.** Protective information may be obtained from your Cooperative Agricultural Extension Service. Do not use when value of bees as pollinators is more important than insect control. Warn beekeepers to locate hives beyond bee flight range until one week after application, or take equally effective precautions.

USER SAFETY REQUIREMENTS

- Wear long-sleeved shirt, long pants, shoes plus socks and household latex or rubber gloves when applying this product
- Change clothing as soon as possible after use.
- Wash the outside of gloves before removing. As with any pesticide product, wash hands thoroughly immediately after handling and before eating, smoking or using the toilet.
- Do not allow children or pets to contact treated area until dust has settled.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Wearing household latex or rubber gloves, use a dust applicator such as a shaker can, dust gun, puffer or other bulbous duster, rotary duster or other appropriate duster as available. Dust lightly to cover both upper and lower leaf surfaces with a thin, even film of dust. Apply when air is calm to avoid drift and contact with eyes and skin.

Start applying at the farthest corner of the treatment area and work backward to avoid contact with dusted surfaces. Allow dust to settle in treated areas before reentering.

For plants above chest height, consider using a liquid carbaryl product. Do not repeat application more than once a week.

STORAGE & DISPOSAL

Do not contaminate water, food or feed by storage or disposal. Store in tightly closed, original container in a locked area away from children and domestic animals. Do not reuse empty container. Securely wrap closed container in several layers of newspaper and discard in trash.

GENERAL INFORMATION

Sevin is an outstanding insecticide that controls a wide variety of insect pests. Sevin is not effective against mites; where mites are a problem, use a suitable miticide.

This product is compatible with commonly used miticides, fungicides and insecticides. Do not use with lime or other alkaline material.

FOR SMALL MEASUREMENTS

- 57 standard level tablespoonful = 1 lb.
- 3 1/2 standard level tablespoonful = 1 oz.
- 4 1/2 ounces = 1 cupful

ORNAMENTALS: Azalea, Boxwood, Rose, and Spirea,: For control of Japanese Beetle, Bagworm, Aphid, Flea Beetle, Leafhoppers, Tent Caterpillar and exposed Thrips: Apply 2 1/2 lbs. Per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 to 10 day intervals, as necessary. Do not use on Boston Ivy or Virginia Creeper.

Subsequent planting of food and feed crops in treated soil must be made with crops on this or other carbaryl label(s).

PREHARVEST USE LIMITATION

Allow 2 days between applications and harvest on sweet corn ears. Do not apply within 48 days of harvest of grain and fodder, or within 14 days of harvest or grazing of forage or silage. Allow 3 days between application and harvest for fresh beans, cucumbers, summer squash, tomatoes, eggplant, or peppers. Allow 7 days between application and harvest on potatoes and on small fruit and berry crops (including blueberries, cranberries, and strawberries). Allow 14 days between application and harvest on lettuce (both head and leaf lettuce).

Apply this product when insects first appear, preferably before damage from feeding occurs, or as indicated. For best results apply thoroughly to upper and lower leaf surfaces. If necessary, repeat applications every 7 to 10 days, or as indicated. Apply with conventional dusting equipment. Do not dust too heavily or crop injury may result.

BEANS: For control of Mexican Bean Beetle: Apply 2 lbs. per 1,000 sq. ft. For control of Leafhoppers, Bean Leaf Beetle, Cucumber Beetles, Western Bean Cutworm, Flea Beetles and Japanese Beetles: Apply 1 1/2 lbs. per 1,000 sq. ft. For control of Corn Earworm, Armyworms and Velvet Bean Caterpillars: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Lima Bean Pod Borer, Lygus and Stink Bugs: Apply 2 1/2 lbs. per 1,000 sq. ft.

CORN: For control of Corn Earworm, European Corn Borer, Fall Armyworm, Sap Beetles and Japanese Beetle: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Larvae in whorl (budworm damage) and foliage feeders, apply to entire plant. For insects attacking Silks and Ears: Apply when first silks appear and continuing until silks begin to dry. Three or more applications may be required

depending on severity of infestation. Timing and good coverage are absolutely essential for effective control.

CUCUMBERS AND SUMMER SQUASH: For control of Pickleworm, Melonworm, Cucumber Beetles, Squash Bugs and Flea Beetles: Apply 1 1/2 lbs. per 1,000 sq. ft.

LETTUCE: For control of Armyworm, Cabbage Caterpillar, Cabbage Looper, Corn Earworm, Flea Beetle, Harlequin Bug, Leafhopper, Six-spotted Leafhopper, Spittlebug, Lygus Bug and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat as needed at 7 day intervals.

OKRA: For control of Corn Earworm and Stink Bugs: Apply 2 lbs. per 1,000 sq. ft. Make first application when insects are first noted and repeat at 7 day intervals, or as necessary.

POTATOES, TOMATOES, EGGPLANTS AND PEPPERS: For control of Colorado Potato Beetle, Flea Beetles, and Leafhoppers: Apply 1 1/2 to 2 lbs. per 1,000 sq. ft. For control of Tomato Fruitworm, Fall Armyworm, Tomato Hornworm, European Corn Borer, Lygus and Stink Bugs: Apply 2 1/2 lbs. per 1,000 sq. ft.

BLUEBERRIES: For control of Cherry and Cranberry Fruitworm: Apply 2 lbs. per 1,000 sq. ft.

CRANBERRIES: For control of Fruitworms, Fireworms, Leafhoppers and Cutworms: Apply 2 1/2 to 3 lbs. per 1,000 sq. ft.

STRAWBERRIES: For control of Meadow Spittlebug and Leaf Roller: Apply 2 1/2 lbs. per 1,000 sq. ft.

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product. **DISCLAIMER:** SureCo warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. SureCo makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use, handling or application of this product.

For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticides Telecommunications Network at 1-800-858-7378.

per EPA letter 10/7/98