

515552

DEC - 9 1996

Ms. Stacy Schmidt
SureCo, Inc.
10012 N. Dale Mabry, Suite 221
Tampa, FL 33618

Dear Ms. Schmidt:

Subject: Amendment - Add Apples and Pears
Diazinon 22.4% Lawn & Garden Water Based Insecticide
EPA Registration Number 769-974
Your submission dated September 12, 1996

The amendment referred to above, submitted in connection with registration under FIFRA section 3(c)(7)(a), is acceptable provided that you:

- 1. Submit and/or cite all data or other material required for registration/reregistration of your product under FIFRA section 3(c)(5) or FIFRA section 4 when the Agency requires all registrants of similar products to submit such data.
- 2. Make the labeling changes listed below before you release the product for shipment bearing the amended labeling
 - a) On the container label, in the list of **Problem Area** and **Pests** for **Fruits & Nuts** after the entry for Walnuts add the following:

Apples, Pears	Aphids, Apple Maggots, Coddling Moths, Leafhoppers, Scale Crawlers, Spider Mites (Pears), Pear Psylla (Pears), and others
------------------	---

- b) In the booklet label, the last sentence of the first paragraph under the heading General Information should show the name of your product rather than Ciba's.
- c) In the booklet label, in the first entry for pears, move the pest listing Mealybugs through Tentiform leafminers up to after Coddling moths (be sure to add a comma after "Coddling moths").

2916
7

3) Submit three (3) copies of your final printed labeling before you release the product for shipment.

If these conditions are not complied with, the registration will be subject to cancellation in accordance with FIFRA section 6(e). Your release for shipment of the product constitutes acceptance of these conditions.

A stamped copy of the label is enclosed for your records.

Sincerely,

George T. LaRocca
Product Manager 13
Insecticide/Rodenticide Branch
Registration Division (7505C)

39 16

DIAZINON 22.4% LAWN & GARDEN WATER BASED INSECTICIDE

REC'D EPA/OPP/DPD1
96 OCT 16 AIO:42

Water-Based Concentrate

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

One Pint Net Contents

Active Ingredient:
Diazinon: O,O-Diethyl
O-(2-Isopropyl-6-
methyl-4-pyrimidinyl)
phosphoro-
thioate22.4%

**Inert
Ingredients: 77.6%**

Total: 100.0%

**KEEP OUT OF REACH
OF CHILDREN.**

CAUTION

See side panel for additional precautionary statements.

See full label in booklet under cap for complete directions for mixing and applying this product and a complete list of pests controlled.

Protects
Lawns
Vegetables
Fruits and Nuts
Roses, Flowers
Trees, Shrubs

Controls
Ants, Aphids
Chinch Bugs
Cutworms
Fire Ants¹
Flea Beetles
Grubs, Mites
Sod Webworms
...and many others
¹Aids in control

EPA Reg. No. 769-974

EPA Est. No. 769-GA-1

ACCEPTED
with COMMENTS
in EPA Letter Dated

DEC - 9 1996

Manufactured by:
SureCo, Inc.
Fort Valley, GA 31030

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

769-974

4916
7

- Concentrated
- Broad Spectrum Pest Control
- Less Smell
- Low Odor
- Reduced Odor Formulation
- Easy-to-Mix
- Easy-to-Apply
- Handles Easily
- Dilute and Spray
- Excellent Performance
- Controls soil, foliar, and sucking pests
- Convenient water-based formula
- Concentrated = Better Value
- Protects Home Lawns and Home Vegetable Gardens
- Controls pests in the home lawn and home garden
- Controls a wide spectrum of pests
- Soil and Foliar Insecticide with Broad Spectrum Control
- Spray on foliage or soil to control numerous pests
- Reliable control
- Covers up to _____ sq. ft. of lawn
- Fast acting
- Dependable Control
- Time Tested Insecticide
- Economical
- New Water-Based Formula*
- Use on home lawns, ornamentals, vegetables and fruits

* The term "new" may be used for no longer than 6 months following product introduction into the marketplace.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Diazinon 22.4% Lawn & Garden Water Based Insecticide is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. This product kills many outdoor pests which damage lawns, roses, flowers, trees, shrubs, vegetables and fruits, and insects which annoy pets and people.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, gardenia, or orchids, since plant injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply to food crops grown in greenhouses.

After spraying, wait the following number of days before harvesting fruits, nuts, or vegetables:

1 day	Tomatoes
3 days	Melons, Winter Squash
5 days	Peppers, Strawberries
7 days	Beans, Blackberries, Blueberries, Boysenberries, Broccoli, Cauliflower, Cucumbers, Dewberries, Loganberries, Peas, Raspberries, Summer Squash, Sweet Corn
10 days	Chinese Broccoli, Chinese Cabbage, Collards, Kale
14 days	Carrots, Endive (Escarole), Lettuce, Onions (Bulb and Green), Parsnips, Radishes, Rutabagas, Spinach, Turnips
21 days	Apricots, Cabbage, Cherries, Nectarines, Peaches, Plums, Prunes
28 days	Grapes
35 days	Potatoes (Irish)
45 days	Walnuts
*	Almonds (one dormant spray only)

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

* This table is only a partial listing of the pests controlled by this product. See booklet under cap for complete directions for use and pests controlled.

Problem Area	Pests*
Nuisance Pests in Outside Areas	Ants, Brown Dog Ticks, Chiggers, Cockroaches, Crickets, Fleas, Fire Ants, Spiders, and others*
Lawns	Chinch Bugs, Ants, Sod Webworms, Brown Dog Ticks, Fleas, White Grubs, Lawn Billbugs, and others*
Vegetables: Foliar Application	
Beans, Peas (except dried varieties)	Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites
Broccoli, Chinese Broccoli, Cabbage, Chinese Cabbage, Cauliflower, Collards, Kale	Aphids, Flea Beetles, Dipterous Leafminers, Diamondback Moths, Imported Cabbageworms
Endive (Escarole), Lettuce, Spinach	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites
Peppers, Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms (Tomatoes), Flea Beetles, Leafhoppers, Southern Armyworms (Tomatoes), Vinegar Flies - <i>Drosophila</i> spp. (Tomatoes)
Onions (Bulb and Green)	Onion Thrips
Carrots, Parsnips, Potatoes, Radishes, Rutabagas, Turnips	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Colorado Potato Beetles (Potatoes), Southern Armyworms (Potatoes)
Sweet Corn	Aphids, Flea Beetles, Grasshoppers, Spider Mites
Melons, Squash (Summer, Winter)	Aphids, Thrips, Spider Mites

6916
8

Vegetables: Preplant Soil Application

Beans, Cutworms, Mole Crickets, Wireworms (Snap Beans)

Peas (except dried varieties)

Beets, Red (Table) Cutworms, Mole Crickets

Cabbage Cutworms, Root Maggots, Wireworms

Carrots, Sweet Corn Cutworms, Mole Crickets, Wireworms

Lettuce, Tomatoes Cutworms, Wireworms

Radishes, Rutabagas, Turnips Cutworms

Fruits & Nuts

Almonds (CA only) Scale Crawlers

Walnuts (CA only) Aphids, Scale Crawlers, Mites, and others*

Apricots, Peaches, Nectannes Aphids, Apricot Mealybugs (Apricots), Spider Mites, Scale Crawlers

Blueberries Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Two-Spotted Spider Mites, Thrips

Cherries Black Cherry Aphids, Cherry Fruitflies, Cherry Rust Mites, Scale Crawlers, and others*

Plums, Prunes Aphids, Spider Mites

Caneberries, Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries (CA, OR, WA only - all crops) Aphids, Leafhoppers, Two-Spotted Spider Mites, Thrips, Raspberry Fruitworms

Strawberries Aphids, Two-Spotted Spider Mites, Strawberry Leafrollers

Grapes Aphids, Grape Berry Moths, Grape Leafrollers, Leafhoppers

Roses & Flowers

Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses Aphids, Flea Beetles, Dipterous Leafminers, Leafhoppers, Mealybugs, Two-spotted Spider Mites, Thrips, Whiteflies, and others*

Ornamental Trees & Shrubs

Arborvitae, Azaleas, Aphids, Mites, Scale Crawlers, Whiteflies, Tent Caterpillars, and others*

Birch, Boxwood,
 Camellias, Crab Apple,
 Dogwood, Douglas Fir,
 Elm, Euonymus,
 Hawthorn Holly,
 Honeysuckle,
 Honey Locust,
 Juniper, Lilac,
 Locust, Maple,
 Mimosa, Oak, Pine,
 Ornamental Plum,
 Poplar,
 Rhododendron,
 Scotch Pine, Spirea,
 Spruce, Sycamore,
 Wax Plant (Hoya),
 Willow, Yellowwood,
 Yew

Storage and Disposal

Storage

Store in original container in an area inaccessible to children or pets. If product freezes, place container in a warm room (68°F or above) inaccessible to children or pets and allow to thaw. Shake well before using.

Disposal

Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by

injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where

watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

DIAZINON 22.4% LAWN & GARDEN WATER BASED INSECTICIDE

Water-Based Concentrate

This product must not be used on golf courses and sod farms.

For outdoor use on the home lawn and the home garden only.

Spray concentrate. Contains diazinon insecticide.

One Pint

Net Contents

Active Ingredient:
Diazinon: O, O-Diethyl
O-(2-isopropyl-6-
methyl-4-pyrimidinyl)
phosphoro-
thioate 22.4%

Inert
Ingredients: 77.6%

Total: 100.0%

KEEP OUT OF REACH
OF CHILDREN.

CAUTION

See end of booklet for
additional precautionary
statements.

EPA Reg. No. 769-974

EPA Est. No. 769-GA-1

Protects

Lawns
Vegetables
Fruits and Nuts
Roses, Flowers
Trees, Shrubs

Controls

Ants, Aphids
Chinch Bugs
Cutworms
Fire Ants¹
Flea Beetles
Grubs, Mites
Sod Webworms
...and many others

¹Aids in control

Manufactured by:
SureCo, Inc.
Fort Valley, GA 31030

IMPORTANT: Read these entire Directions and Conditions of Sale before using this product. **DISCLAIMER:** SureCo warrants that this product conforms to the chemical description on the label and is reasonably fit for the specific purposes referred to in the Directions For Use, subject to the inherent risks referred to above. SureCo makes no other express or implied warranty of fitness or merchantability nor any other express or implied warranty. In no case shall SureCo be liable for consequential, special, or indirect damages resulting from the use, handling or application of this product.

- Do not exceed maximum permitted label rates, since rates above those recommended significantly increase potential hazards to birds and waterfowl.
 - Avoid overlapping of sprays.
 - Where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs.
- Use on Bent, Bermuda, Bluegrass, Dichondra, Fescue, Merion, and St. Augustinegrass. To control pests listed below, apply recommended rates.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR INSECT CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

General Information

Water Based Insecticide

Diazinon 22.4% Lawn & Garden is a water-based concentrate product which must be diluted with water before application. To avoid having spray mixture left over, mix only the amount needed. Refer to the section **How to Mix and Apply**. This product kills many outdoor insects which damage lawns, roses, flowers, trees, shrubs, vegetables, fruits and nuts, and insects which annoy pets and people. Contains diazinon insecticide which provides positive control of many "hard-to-kill" sucking and chewing insects. Diazinon Lawn & Garden WBC works fast and applications should be repeated only as directed to maintain control.

Notes: (1) Do not use inside the home. (2) Do not use on ferns, poinsettia, hibiscus, papaya, pilea, geranium, or orchids, since injury may occur. (3) Use this product on succulent peas and beans only. Do not use on dried pea or bean varieties. (4) Do not apply to food crops grown in greenhouses.

Precautions: (1) Wear a long-sleeved shirt and long-legged pants when spraying. (2) Spray with the wind to your back. (3) Do not spray on windy days. (4) If clothing becomes wet from spray during use, remove clothing after spraying, wash affected body areas thoroughly with soap and water, and launder clothing before wearing again.

How to Mix and Apply

Step #1: Determine size of spraying job. For lawns and vegetables, measure the area to be sprayed. For all other applications, apply as a thorough cover spray. Make applications when insects first appear. Applications should be repeated only as directed to maintain control.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Diazinon 22.4% Lawn & Garden Water Based Insecticide and water as indicated in the rate chart.

Step #3: Spray as directed in rate chart. Thorough coverage is important.

See chart for complete list of use sites, crops, and insects controlled. Use the rate given in the chart for the type of insect you want to control on the site or crop you are spraying.

Step #4: Flush sprayer with clean water after each use.

Measurement Conversions

- 1 tablespoon = 3 teaspoons
- 1 fl. oz. = 6 teaspoons or 2 tablespoons
- 6 fl. oz. = 3/4 cup
- 1 1/2 tablespoons in 3 gals. of water = 1 1/2 teaspoons in 1 gal. of water
- 2 tablespoons in 3 gals. of water = 2 teaspoons in 1 gal. of water

Lawn Pest Control

Precautions

- Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Apply this product only as specified on this label.
- Because of the migratory habits of certain waterfowl, do not apply in Nassau County, New York, between November 1 and May 20.

Pests	Rate		Remarks
	Diazinon 22.4%	Water	
Lawn Chinch Bugs	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Thoroughly wet down grass a few hours before applying. In the North, if Chinch Bugs are the only problem, a lower rate of application can be used: 3 teaspoons in 3 gals. of water covers 125 sq. ft. or 4 fl. oz. in 24 gals. of water per 1,000 sq. ft.
Ants, Armyworms, Bermudagrass Mites, Clover Mites, Springtails (Collembola), Crickets, Cutworms, Digger Wasps, Earwigs, Frit Flies, Lawn Billbugs, Leafhoppers, Sod Webworms (Lawn Moths), Sowbugs (Pillbugs)	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Thoroughly wet down grass a few hours before applying. For ants and digger wasps, spot spray ant hills and wasp ground nest openings. For digger wasps, apply at dusk, when wasps are not active. Billbug control: Apply when activity is first observed or when chewed or brown grass is present indicating damage from the insect. For frit flies, mow grass and water thoroughly before treatment, but delay application until grass is dry.
Brown Dog Ticks, Chiggers, Fleas	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Note: Do not apply this product to animals. For brown dog ticks, spray grass and under shrubbery, particularly near house.
Hyperodes Weevil	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Treat problem areas in mid-April and again in mid-May. Water grass thoroughly after application.
Millipedes	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Thoroughly wet grass a few hours before applying.
Rhodesgrass Scales	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Apply when crawlers first emerge.
White Grubs of Japanese Beetle, European Chafer, Southern Chafer	1 1/2 tablespoons per 125 sq. ft. or 6 fl. oz. per 1,000 sq. ft.	3 gals. or 24 gals.	Apply any time between late July and early October. Water grass thoroughly after application.

10/16

Nuisance Pests Outside the Home

<p>Ants, Boxelder Bugs, Brown Dog Ticks, Chiggers, Clover Mites, Cockroaches, Crickets, Earwigs, Fleas, Flies, Millipedes, Sowbugs (Pillbugs), Spiders, Springtails (<i>Collembola</i>)</p>	<p>Mix 2 fl. oz. in 1 gal. of water and thoroughly spray window and door screens, sills, foundations, porch and patio walls, entranceways, walks, garbage cans, tree trunks, and into cracks and other places where insects can hide.</p> <p>To prevent entry into houses, spray a 5-ft. band of soil around the house next to the foundation and spray the foundation of the house to a height of 2-3 ft. Repeat application as necessary.</p> <p>Do not treat animals with this product. Do not treat plants (other than tree trunks) at this nuisance pest rate.</p>
<p>Fire Ants</p>	<p>To aid in the control of fire ants in home lawns and other outdoor home recreation areas, apply 2 tablespoons in a minimum of 1 gal. of water as a drench. Apply over and 6 inches around each mound. Apply gently to avoid disturbing ants. Reapply as necessary. Larger quantities of solution can be prepared for multiple mound treatments. Use equipment capable of delivering the diluted product as a gentle rain at the rate of 1 gal. per mound. High pressure sprays may disturb the ants and cause migration, reducing product effectiveness.</p> <p>Note: For best results, apply in cool weather 65-80°F, or in early morning or late evening hours. Treat new mounds as they appear.</p>

Insect Control on Ornamental Trees, Shrubs, and Flowers

Spray when listed pests first appear. Repeat as necessary.

Trees and Shrubs	Insects	Rate per 1 Gal. of Water	Timing/Special Directions
<p>Arborvitae, Azaleas, Birch, Boxwood, Camellias, Crab Apple, Dogwood, Douglas Fir, Elm, Euonymus, Hawthorn, Holly, Honey Locust, Honeysuckle, Juniper, Lilac, Locust, Maple, Mimosa, Oak, Pine, Ornamental Plum, Poplar, Rhododendron, Scotch Pine, Spirea, Spruce, Sycamore, Wax Plant (Hoya), Willow, Yellowwood, Yew</p>	<p>Aphids, Apple-and-Thorn Skeletonizers, Bagworms, European Pine Sawfly, European Pine Shoot Moths, Holly Bud Moths, Dipterous Leafminers (Arborvitae, Azalea, Birch, Boxwood, Holly, Oak), Hemlock Charmes, Leafhoppers, Mealybugs, Mites (Clover, Cyclamen, European Red, Privet, Two-spotted Spider), Oak Loopers, Oblique-banded Leafrollers, Obscure Root Weevils, Omnivorous Leaf-tiers, Pear Slugs, Scale Crawlers (Cottony-cushion, Euonymus, Lecanium, Pine Needle, Pit, San Jose, Soft), Tent Caterpillars, Webworms (Cotoneaster, Fall, Juniper, Mimosa), Whiteflies</p>	<p>2 tea-spoons</p>	<p>Apply as a thorough foliar spray. Wet leaves and branches to the dripping point. Try to penetrate dense foliage with spray.</p> <p>Mimosa Webworms: apply when adults first appear. Make 2 additional applications at 4 to 5-week intervals.</p>

See Copy A

Roses and Flowers	Insects	Rate per 1 Gal. of Water	Timing/Special Directions
<p>Azaleas, Camellias, Chrysanthemums, Carnations, Gladioli, Lilac, Lilies, Marigolds, Pansies, Petunias, Rhododendron, Roses</p>	<p>Aphids, Dipterous Leafminers, Euonymus Scale Crawlers, Flea Beetles, Leafhoppers, Mealybugs, Mites (Carnation Bud, Carnation Shoot, Cyclamen, European Red, Two-spotted Spider), Thrips, Whiteflies</p>	<p>2 tea-spoons</p>	<p>Apply as a thorough foliar spray. Try to spray underside of leaves and penetrate dense foliage.</p>

Fruits and Nuts

Start spraying after blooming. Applications should be made according to infestation or when aphids, mites, and crawler stages of scale insects are first noted.

Observe stated time intervals between last application and harvest of the crop, as well as intervals between applications and total number of applications that can be made per season.

Crop	PHI*
Almonds	**
Apricots	21
Blackberries	7
Blueberries	7
Boysenberries	7
Cherries	21
Dewberries	7
Grapes	28
Loganberries	7
Nectarines	21
Peaches	21
Plums	21
Prunes	21
Raspberries	7
Strawberries	5
Walnuts	45

* PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

** Dormant spray only.

Note: To protect bees, do not apply to fruit trees when trees or substantial numbers of weeds around the trees are in bloom.

Crop	Pests	Rate	Timing/Special Directions
<p>Almonds (CA only)</p>	<p>San Jose Scale, Parlatoria Scale</p>	<p>2 teaspoons in 1 gal. of water</p>	<p>Apply only as a dormant spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard.</p> <p>This product may be used on almonds only in CA. Do not use this product on almonds grown in other states.</p>
<p>Apricots</p>	<p>Aphids, Apricot Mealybugs, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)</p>	<p>2 teaspoons in 1 gal. of water</p>	<p>Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest.</p>

Blueberries	Aphids, Blueberry Maggots, Cherry Fruitworms, Cranberry Fruitworms, Thrips, Two-spotted Spider Mites	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of blueberry patch. Do not make more than 5 applications per season, a minimum of 14 days apart. Do not apply within 7 days of harvest.
Caneberries (CA, OR, WA only): Blackberries, Boysenberries, Dewberries, Loganberries, Raspberries	Aphids, Leafhoppers, Raspberry Fruitworms, Two-spotted Spider Mites, Thrips	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 5 gals. per 500 sq. ft. of caneberry patch. Do not make more than 5 applications per season, a minimum of 14 days apart. Do not apply within 7 days of harvest. This product may be used on caneberries only in CA, OR, and WA. Do not use this product on caneberries grown in other states.
Cherries	Black Cherry Aphids, Cherry Fruitflies, Cherry Fruitworms (Maggots), Fruitree Leafrollers, Mites (Cherry Rust), Scale Crawlers (San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 21 days of harvest.
Grapes	Aphids, Grape Berry Moths, Grape Leafhoppers, Leafhoppers	1 1/2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of home vineyard. Do not make more than 5 applications per season, a minimum of 7 days apart. Do not apply within 28 days of harvest.
Nectarines	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest.
Peaches	Aphids, Mites (Brown, Two-spotted Spider), Scale Crawlers (Olive, San Jose)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest.

Plums and Prunes	Aphids (Leafcurl Plum, Mealy Plum, Thistle), Mites (Brown, European Red)	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 21 days of harvest.
Strawberries	Aphids, Two-spotted Spider Mites, Strawberry Leafrollers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 3 gals. per 500 sq. ft. of strawberry patch. Do not make more than 3 applications per season, a minimum of 7 days apart. Do not apply within 5 days of harvest.
Walnuts (CA only)	Aphids, Scale Crawlers (Walnut, Italian Pear, Frosted), Spider Mites, Twig Borers	2 teaspoons in 1 gal. of water	Apply as a full cover spray. Do not apply more than 6 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply after husks open or split. Do not apply within 45 days of harvest. This product may be used on walnuts only in CA. Do not use this product on walnuts grown in other states.

→ See Copy B

Vegetables

For foliar or soil applications, **Diazinon 22.4%** should be applied in the volume of water recommended in the table below to assure thorough coverage of the foliage or soil.

Observe stated time intervals between last application and harvest as well as intervals between applications and maximum numbers of applications per season. Do not apply to food crops grown in greenhouses.

Pre-Harvest Intervals (PHI*) for Foliar Applications

Crop	PHI* (Days)
Beans ¹ (succulent varieties only: Lima, Pole and Snap)	7
Broccoli	7
Cabbage	21
Cantaloupes	3
Carrots	14
Casabas	3
Cauliflower	7
Chinese Broccoli, Chinese Cabbage, Collards	10
Crenshaws	3
Cucumbers	7
Endive (Escarole)	14
Honeydew Melons	3
Kale	10
Lettuce	14
Muskmelons	3
Onions (Bulb and Green), Parsnips	14
Peas ¹	7
Peppers	5
Persian Melons	3
Potatoes (Irish)	35
Radishes, Rutabagas, Spinach	14
Squash, Summer	7
Squash, Winter	3
Sweet Corn	7
Tomatoes	1
Turnips (Roots and Tops)	14
Watermelons	3

*PHI = Pre-Harvest Interval, or the number of days between the last application and harvest.

¹ Do not use this product on dried pea or bean varieties, such as pinto beans, dried limas, split peas, or dried black-eyed peas.

Vegetables

For application to the foliage, thoroughly spray all plant parts. Application to the soil should be mixed into the soil immediately after application.

Crop	Pests	Rate	Timing/Special Directions
Beans, Lima, Pole, and Snap (succulent varieties only)	Aphids, Cucumber beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 5 days up to 3 applications per season. Do not apply within 7 days of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms (snap beans only)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Notes: (1) Do not apply this product to dried bean varieties, such as pinto beans or dried limas. (2) To protect bees, do not apply bean foliage treatments if the beans or weeds in the treatment area are in bloom.			
Beets, Red (Table)	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Cabbage	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 21 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Cabbage (cont.)	Root Maggots	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Carrots	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Cauliflower	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Broccoli	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Chinese Cabbage (Bok Choy and Napa)	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.

Collards	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Corn (see sweet corn)			
Cucumbers	Aphids, Thrips, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 7 days of harvest.
Endive (Escarole)	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Kale	Aphids, Diamondback Moths, Flea Beetles, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 10 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
Lettuce	Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.

<p>Melons (Cantaloupes, Casabas, Cran-shaws, Honey-dews, Musk-melons, Persians, and Hybrids of these) and Water-melons</p>	<p>Aphids, Thrips, Spider Mites</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 3 days of harvest.</p>
<p>Onions (Bulb and Green)</p>	<p>Onion Thrips</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 14 days of harvest.</p>
<p>Parsnips</p>	<p>Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat applications every 7 days, up to 5 applica- tions per season. Do not apply within 14 days of harvest. Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.</p>
<p>Peas (succulent only)</p>	<p>Aphids, Cucumber Beetles, Grasshoppers, Leafhoppers, Mexican Bean Beetles, Spider Mites</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 5 days, up to 3 applica- tions per season. Allow at least 7 days between the last ap- plication and harvest.</p>
	<p>Cutworms (surface, subterranean)</p>	<p>6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water</p>	<p>Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.</p>
	<p>Mole Crickets</p>	<p>2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water</p>	<p>Spray the soil evenly 1-2 days before plant- ing. Immediately work into soil 1-2 inches.</p>
	<p>Wireworms</p>	<p>6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water</p>	<p>Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.</p>
<p>Notes: (1) Do not apply this product to dried pea varieties, such as split peas or dried black-eyed peas. (2) To protect bees, do not apply foliage treatments if peas or weeds in the treatment area are in bloom.</p>			
<p>Peppers</p>	<p>Aphids, Colorado Potato Beetles, Flea Beetles, Leafhoppers</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 5 days of harvest.</p>

<p>Potatoes</p>	<p>Aphids, Flea Beetles, Leafhoppers, Colorado Potato Beetles, Southern Armyworms, Dipterous Leafminers</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 35 days of harvest. Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.</p>
<p>Radishes</p>	<p>Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 3 applica- tions per season. Do not apply within 14 days of harvest. Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.</p>
	<p>Cutworms (surface, subterranean)</p>	<p>6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water</p>	<p>Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.</p>
<p>Rutabagas</p>	<p>Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 14 days of harvest. Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.</p>
	<p>Cutworms (surface, subterranean)</p>	<p>6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water</p>	<p>Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for sur- face cutworms, 3-6 inches for subterra- nean cutworms.</p>
<p>Spinach</p>	<p>Aphids, Flea Beetles, Leafhoppers, Dipterous Leafminers, Spider Mites</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 14 days of harvest. Note: Resistant leaf- miners will not be con- trolled by diazinon or similar type insecti- cides.</p>
<p>Squash, Summer</p>	<p>Aphids, Thrips, Spider Mites</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 7 days of harvest.</p>
<p>Squash, Winter</p>	<p>Aphids, Thrips, Spider Mites</p>	<p>1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden</p>	<p>Thoroughly spray all plant parts. Repeat application every 7 days, up to 5 applica- tions per season. Do not apply within 3 days of harvest.</p>

Sweet Corn	Aphids, Thrips, Flea Beetles, Grasshoppers, Spider Mites	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Corn may be picked 7 days following last application.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Mole Crickets	2 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly 1-2 days before planting. Immediately work into soil 1-2 inches.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Note: To protect bees, do not apply to corn foliage during the pollen shed period.			
Tomatoes	Aphids, Colorado Potato Beetles, Fall Armyworms, Flea Beetles, Leafhoppers, Southern Armyworms, Vinegar Flies (<i>Drosophila</i> spp.)	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 1 day of harvest.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.
	Wireworms	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 4-8 inches.
Turnips and Turnip Tops	Aphids, Flea Beetles, Leafhoppers, Imported Cabbageworms, Dipterous Leafminers	1 teaspoon in 1 gal. of water to cover 500 sq. ft. of garden	Thoroughly spray all plant parts. Repeat application every 7 days up to 5 applications per season. Do not apply within 14 days of harvest. Note: Resistant leafminers will not be controlled by diazinon or similar type insecticides.
	Cutworms (surface, subterranean)	6 tablespoons per 500 sq. ft. in a minimum of 1 gal. of water	Spray the soil evenly just before planting. Immediately work into soil 2-3 inches for surface cutworms, 3-6 inches for subterranean cutworms.

Storage and Disposal

Storage

Store in original container in an area inaccessible to children and pets. If product freezes, place container in a warm room (68°F or above) inaccessible to children or pets and allow to thaw. Shake well before using.

Disposal

Do not pour the concentrate, the spray mixture, or the rinse water down the drain or sewer. Do not reuse empty container. Triple rinse the empty container, pour the rinse water into the sprayer, then spray the area to be treated. Securely wrap empty container in several layers of newspaper and discard in trash. If disposal of a partially full container is necessary, make sure cap is tightly closed, securely wrap the container in several layers of newspaper and discard in trash.

Precautionary Statements

Hazards to Humans and Domestic Animals

CAUTION

Harmful if swallowed, inhaled, or absorbed through skin. Avoid contact with eyes, skin, or clothing. Avoid breathing spray mist. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse. Do not use inside the home. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not allow children or pets to go onto a treated area until the treated area, such as grass or soil, has dried. Do not contaminate ornamental fish ponds. Food utensils, such as spoons and measuring cups, must not be used for food purposes after use in measuring pesticides.

Statement of Practical Treatment

If swallowed: Call a physician or Poison Control Center immediately. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth if person is unconscious or convulsing.

If in eyes: Flush eyes with plenty of water. Call a physician if irritation persists.

If inhaled: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

If on skin: Wash with plenty of soap and water. Get medical attention.

Note to Physician: This product is an organophosphate insecticide. If symptoms of cholinesterase inhibition are present, atropine sulfate by injection is antidotal. 2-PAM is also antidotal and may be administered, but only in conjunction with atropine. If ingested, induce emesis or lavage stomach.

Environmental Hazards

This product is highly toxic to birds, fish, and other wildlife. Birds, especially waterfowl, feeding or drinking on treated areas may be killed. Because of the migratory habits of certain Atlantic Coast waterfowl, do not apply this product to lawns in Nassau County, New York, between November 1 and May 20. Do not exceed maximum permitted label rates. Application rates above those recommended significantly increase potential hazards to birds and waterfowl. Avoid overlapping of sprays. For lawns, where watering is recommended in the directions, or if waterfowl (ducks or geese) can be expected in the treated area after treatment (except spot treatment), water lawn with at least 1/4 inch of water immediately after spraying this product; however, stop watering before puddling occurs. Keep out of lakes, streams, ponds, tidal marshes, and estuaries. Do not apply directly to water. Runoff may be hazardous to aquatic organisms in neighboring areas. Shrimp and crab may be killed at application rates recommended on this label. Do not apply where fish, shrimp, crab, and other aquatic life are important resources. Do not contaminate water by cleaning of equipment or disposal of equipment wash waters.

This pesticide is highly toxic to bees exposed to direct treatment or to residues on blooming crops or weeds. Do not apply this pesticide or allow it to drift to blooming crops or weeds if bees are visiting the treatment area.

COPY A

Crop	Pests	Rate	Timing/Special Directions
Apples	Aphids, Apple maggots, Codling moths, Eyespotted budmoths, Fruit Tree Leafrollers, Leafhoppers, Mealybugs, Scale crawlers (San Jose Forbes), Spider Mites	2 teaspoons in 1 gallon of water	Apply as a full cover spray. Do not apply more than 4 gal. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots may not be controlled by diazinon or similar type insecticides.
Pears	Aphids, Apple maggots, Codling moths Mealybugs, Scale crawlers (San Jose Forbes), Spider mites, Tentiform leafminers	2 teaspoons in 1 gallon of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season, a minimum of 14 days apart. Do not apply within 21 days of harvest. Note: Resistant apple maggots will not be controlled by diazinon or similar type insecticides.
	Pear Psylla	1-2 teaspoons in 1 gallon of water	Apply as a full cover spray. Do not apply more than 4 gals. per 500 sq. ft. of home orchard. Do not make more than 3 applications per season a minimum of 14 days apart. Do not apply within 21 days of harvest.

COPY B