


MICROFILM COPY

Clinafarm® EC Emulsifiable Concentrate

For control of *Aspergillus fumigatus* contamination in poultry hatchery equipment

ACTIVE INGREDIENT:
Imazal (feniconazole) 1-(2-(2,4-Dichlorophenyl)-2-(2-propenyl)oxy)ethyl-1H-imidazole 13.8% w/w
INERT INGREDIENTS 86.2%
100.0% w/w



**KEEP OUT OF REACH OF CHILDREN
DANGER**

See Side Panel for Additional Precautionary Statements

TM

STATEMENT OF PRACTICAL TREATMENT

IF IN EYES: Hold eyelids open and flush with steady, gentle stream of water for 15 minutes. Call physician.

IF SWALLOWED: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger. Do not induce vomiting or give anything by mouth to an unconscious person.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth to mouth. Get medical attention.

Sterwin Laboratories Inc., A Pitman Moore Company, Millsboro, DE 19966. Made in U.S.A. A product of Janssen Research

Sterwin
Laboratories Inc.
A Pitman Moore Company

Net Contents: 750 mL (1.58 U.S. Pints)
EPA Reg. No. 773-55
EPA Est. No. 6175-LA-1

**NOTIFICATION
LABEL NOT REVIEWED
PER FR NOTICE 99-9
DATE 10-16-90**

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

DANGER
Corrosive, causes irreversible eye damage. Harmful or fatal if swallowed. Harmful if absorbed through skin or inhaled. Avoid breathing spray mist and contact with skin. Do not get in eyes, wear goggles or face shield. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. SHAKE WELL BEFORE USING.

For the control of *Aspergillus* organisms in cleaned poultry and turkey hatchery equipment prior to introduction of the egg. Equipment would include empty hatcher cabinets, setters, coolers, storerooms, handling equipment, etc. CLINAFARM EC will dramatically reduce the levels of infectious organisms and spores in treated areas.

To make 1 gallon of a 1:100 dilution, add 1.25 fl. oz. (36 mL) of CLINAFARM EC to one gallon of clean water to provide an antifungal concentration of 0.15% active ingredient. Do not use product undiluted.

To assure proper efficacy, the area or equipment to be treated should be clean and closed off from outside air currents before treatment. Ventilation ducts may be treated but fans must be shut off.

Fog or spray area to be treated at the rate of 1/2 fl. oz. diluted product per 150 cu. ft. (1 ml per 10 cu. ft.). Fog or spray to insure good contact with surface areas. Do not re-enter treatment areas for two hours.

CLINAFARM EC is compatible with other commonly used disinfectants such as quaternary ammonium compounds, glutaraldehyde, formaldehyde and phenolic compounds.

PHYSICAL/CHEMICAL HAZARDS: Do not use or store near heat or open flame.

STORAGE AND DISPOSAL

STORAGE: Store at room temperature.

PESTICIDE DISPOSAL: Do not contaminate water, food or feed by storage or disposal. Do not contaminate water when disposing of equipment washwaters. Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent), then offer for recycling or conditioning, or puncture and dispose of in a sanitary landfill, by incineration or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

CLINAFARM is a registered trademark of Janssen Pharmaceutica NV. (F-15)

