PM 13 773-73 P 1 4 EXspot® (w/ Amblyomma americanum) Inside Panel, Front Revised 2/5/97

DEFEND® EXspot® INSECTICIDE FOR DOGS

FOR USE ON DOGS AND PUPPIES

KILLS & REPELS*

Kills & Repels Fleas up to 4 Weeks

Kills & Repels Deer Ticks (Ixodes scapularis), Vector of LYME DISEASE, up to 4 Weeks

Kills & Repels Lone Star Ticks (Amblyomma americanum) for 2-3 Weeks

Kills & Repels Mosquitoes

Aids in Prevention of Blood Feeding by Mosquitoes,

Vector of HEARTWORM, up to 4 Weeks

Kills & Repels Brown Dog Ticks (Rhipicephalus sanguineus) up to 4 Weeks

Kills & Repels American Dog Ticks (Dermacentor variabilis) for 2 to 3 Weeks

Contains: Two 1cc Applicators

Active Ingredients

Permethrin: (3-phenoxyphenyl) methyl (±)-cis, trans-3-(2,2-dichloroethenyl)-2, 2-dimethylcyclopropanecarboxylate**

65%

Inert Ingredients

<u>35%</u>

Total (wt/wt) 100%

**cis/trans ratio: Max. 55% (±) cis and min. 45% (±) trans

CCEPTED

APR - 1 1997

Inche des Pederal Insecticide, Tagitade, and Rodenticide Act is emended, for the pesticide whitered and at 172, 72 KEEP OUT OF REACH OF CHILDREN
CAUTION

SEE INSIDE PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS

EXspot® (w/o Ámblyomma americanum) Outside Panel, Back Revised 2/5/97

DEFEND® EXspot® INSECTICIDE FOR DOGS

READ ENTIRE LABEL BEFORE EACH USE **USE ONLY ON DOGS**

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. DO NOT USE ON CATS. May be toxic if applied to or ingested by cats. Do not use this product in or on electrical equipment due to the possibility of shock hazard. Do not reapply for 7 days. Do not use on puppies under 4 weeks of age.

use direction picture 1	use direction picture 2	use direction picture 3

HOW TO APPLY

Hold card in upright position. Point tip of tube cap up and away from the user's face and body, then cut along dotted line. Invert card over dog and use point of card to part the hair of dog. Squeeze the tube firmly to apply all of the solution to the dog's skin. Wrap container and put in trash.

use direction picture 4	use direction picture 5

Dosage for Dogs Weighing Less Than 33 Pounds:

Apply 1 (one) cc of Defend® EXspot® insecticide solution to the skin of the dog's back between the shoulder blades.

Dosage for Dogs Weighing More Than 33 Pounds:

Apply 1 (one) cc of Defend® EXspot® insecticide solution to the skin of the dog's back between the shoulder blades and 1 (one) cc of Defend® EXspot® insecticide solution to the skin of the dog's back directly in front of the base of the tail.

For information concerning proper product use or actions to take in the event of an emergency.2 phone:

Product use - 1-800-TIC-FLEA or 1-800-842-3532.

Emergency - 1-800-613-4357

Mallinckrodt

Veterinary Mallinckrodt Veterinary, Inc. 421 E. Hawley Street Mundelein, IL 60060 U.S.A. 0000000A R1296

ı	ot	No.	
ı.	£ 7 £	INU.	

Kills & Repels* U.S. Patent No. 4,20,181, 5,236,954& other Patent Pending

EPA Reg. No. 773-73 EPA Est. No. 6175-LA-1

bar code

EXspot® (w/o Amblyomma americanum) Inside Panel Revised 2/5/97

applicator tube

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION

Harmful if swallowed or absorbed through the skin. Causes moderate eye irritation. Avoid contact with skin, eyes or clothing. Wash throughly with soap and water after handling. DO NOT USE ON CATS. Cats which actively groom or engage in close physical contact with recently treated dogs may be at risk of toxic exposure. Do not use on sick or billitated dogs. Consult with a veterinarian before using on aged or medicated dogs. 3 or on dogs with known organ dysfunction.

FIRST AID

If Swallowed: Call a physician or Poison Control Center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or, if available, by administration of syrup of ipecac. DO NOT INDUCE VOMITING OR GIVE ANYTHING BY MOUTH TO AN UNCONSCIOUS PERSON. If On Skin: Remove contaminated clothing and wash with plenty of soap and water. Call a physician if irritation persists. If In Eyes: Flush with plenty of water. Call a physician if irritation persists. Adverse Reaction: Some animals may be sensitive to ingredients in this product. Reported reactions in dogs have included skin Instituty, increased itchiness, redness, rash, hair discoloration or hair loss at the application site. Dogs may also show lethargy. Observe the dog following treatment. If sensitivity to Defend® EXspot® For Dogs occurs, discontinue use and bathe dog with mild soap and rinse thoroughly. Seek Veterinary advice.4

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of washwaters.

KEEP OUT OF REACH OF CHILDREN
CAUTION

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Store in a cool, dry place. Protect from freezing. Pesticide Disposal: Securely wrap original container in several layers of newspaper and discard in trash. Container Disposal: Do not reuse empty container. Wrap container and put in trash.

Defend® EXspot ® insecticide is an effective and easy to use product. Defend® EXspot® insecticide demonstrates greater than 92% control of fleas within 3 days of application. As with all flea and tick control products, Defend® EXspot® insecticide should be used as part of a control program aimed at reducing flea populations in the dog's environment (bedding, carpets, kennel, yard). Consult your veterinarian or entomologist for program recommendations. Repeat applications may be made as necessary, but do not apply at intervals less than 7 days. DO NOT USE ON CATS.

Exspot® (w/o Amblyomma americanum)
Inside Panel (continued)
Revised 2/5/97

NOTICE OF WARRANTY

MALLINCKRODT VETERINARY, INC. MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE, EXPRESSED OR IMPLIED CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THE PRODUCT UNDER NORMAL CONDITIONS IN ACCORD WITH THE STATEMENTS MADE ON THIS LABEL.

applicator tube