

Volume 1
Page 10

Saber™ Pour-On Insecticide

For Beef Cattle & Calves

For Control of Horn Flies and Lice

Animal
Graphic
Of Cattle

Bottle
Graphic

Graphic
For
Dosing

KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE BACK PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS

ACCEPTED
JAN 3 2001
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 773-74


EPA Reg. No. 773-74
EPA Est. No. 6175-LA-1

Net Contents: 16.9 fl oz (500mL)
Or
Net Contents: 30 fl oz (900mL)
Or
Net Contents: 1 gallon (3.785 L)

Treats 50, 90 or
378
550 lb Cattle

SP Schering-Plough Animal Health

Bar Code



FIRST AID	
If inhaled	<ul style="list-style-type: none"> Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes	<ul style="list-style-type: none"> Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> Call poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact Rocky Mountain Poison Center 303-595-4869 for emergency medical treatment information.	

- 5 -

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

NOTICE OF WARRANTY

SCHERING-PLOUGH ANIMAL HEALTH CORPORATION MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THIS PRODUCT UNDER NORMAL CONDITIONS AND IN ACCORDANCE WITH THE STATEMENTS ON THIS LABEL.

- 6 -

SABER™

POUR-ON INSECTICIDE FOR BEEF CATTLE & CALVES

ACTIVE INGREDIENT	%(W/W)
Lambdacyhalothrin:	
[1α-(S),3α(Z)]-(±)-cyano-	
(3-phenoxyphenyl)-methyl-3-	
(2-chloro-3,3,3-trifluoro-1-propenyl)-	
2,2-dimethylcyclopropanecarboxylate	1.0%
OTHER INGREDIENTS	99.0%
TOTAL	100.0%

 Schering-Plough Animal Health
1095 Morris Ave., Union, NJ 07083
Copyright © 1998, 1999,
Schering-Plough Animal Health Corp.
All rights reserved.

00000000

Rev. 3/00

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

READY TO USE - No dilution necessary. Do not apply to lactating or dry dairy cows. Do not apply this product to face of beef cattle or calves.

Apply To	Target Insects	Dosage
Beef Cattle and Calves	Lice, Horn flies	Apply product down the backline at the rate of: Cattle less than 600 lbs = 10 mL (1/3 fl oz) Cattle greater than 600 lbs = 15 mL (1/2 fl oz)

Repeat treatment as needed; however, do not apply more than once every 2 weeks and do not apply more often than four times within any 6-month period. For reduction of sucking lice, two treatments at a 14-day interval are recommended.

Special Note: SABER™ Pour-On is not recommended for use on veal calves. SABER Pour-On is not effective in controlling cattle grubs; therefore, this product can be used on cattle at any time of the year without fear of host-parasite reactions commonly associated with grub treatment products. SABER Pour-On should be used in an integrated pest management system, which may involve repeat treatments and the use of other pest control practices. Continual exposure of horn flies to a single class of insecticide (eg, pyrethroids or organo-phosphates) may lead to the development of resistance to that class of insecticide. In order to reduce the possibility of horn flies developing resistance, it is important to rotate, on a seasonal basis, the class of insecticide used and/or the method of horn fly control. For advice concerning current control practices with relation to specific local conditions, consult resources in resistance management programs and your Cooperative Agricultural Extension Service.

- 3 -

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION - Harmful if swallowed, absorbed through the skin, or inhaled. Causes moderate eye irritation. Avoid breathing vapors or spray mist. Avoid contact with skin, eyes, or clothing. Remove contaminated clothing and wash before reuse. Wash thoroughly with soap and water after handling. Avoid contact with eyes. Wear rubber or non-permeable protective gloves when applying this product. Data indicate this product to be a contact allergenic. Do not apply product to face of cattle.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying in areas adjacent to any body of water. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of rinse water. Apply this product only as specified on the label.

PHYSICAL OR CHEMICAL HAZARDS
Do not use or store near heat or open flame.

- 4 -

11

2/2