



POUR-ON INSECTICIDE FOR BEEF & DAIRY CATTLE & SHEEP

Squeeze 'N' Measure Bottle

May be used with Automatic Dosing System

ACTIVE INGREDIENT

INERT INGREDIENTS

Keep Out of Reach of Children
CAUTION

SEE BACK FOR PRECAUTIONARY STATEMENTS

Net Contents:
One US Quart (0.946 L)



Schering-Plough
Animal Health



DIRECTIONS FOR USE
It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
READY TO USE — No dilution necessary.

Apply To	Target Species	Dosage
Lactating and Non-Lactating Dairy Cattle, Beef Cattle, and Calves	Cows, Fem. Heifers, Steers, Bulls, and Heifers, and Bulls	Apply 1.5 ml per 100 lbs body weight of animal as a maximum of 30 ml for any one animal. Pour along back and above head.
Sheep	Sheep heads, Lice	Pour along back. Apply 1.5 ml per 50 lbs of body weight of animal up to a maximum of 18 ml for any one animal.

For cattle and sheep, repeat treatment as needed, but not more than once every 2 weeks. For optimum lice control, two treatments at 14-day intervals are recommended.
SPECIAL NOTE: POUR-ON Insecticide is not effective against cattle gnats. POUR-ON Insecticide should be used in an integrated pest management system which may involve repeat treatments and the use of other pest control practices.

PRECAUTIONARY STATEMENTS

HAZARD TO HUMANS AND DOMESTIC ANIMALS

CAUTION — Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Absorption of frequently repeated skin contact may cause allergic reactions in some individuals.

STATEMENT OF PRACTICAL TREATMENT

Harmful if swallowed. Call a physician immediately. DO NOT INDUCE VOMITING UNLESS UNDER MEDICAL ATTENTION. Hints To Physicians: Solvent presents aspiration hazard. Gavage length is indicated if material was taken internally. If in eyes, immediately flush eyes with plenty of water. Get medical attention if discomfort persists.

If On Skin: Wash skin with soap and water. Get medical attention.

ENVIRONMENTAL HAZARDS

This product is extremely toxic to fish and other aquatic invertebrates. Do not add directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on label.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal.

Residue: Do not contaminate water and in use. Do not store near food or feed.

Residue: Repeat treatment meeting with the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: High dose for application. Then offer for recycling or reconditioning, or purchase and dispose of in a sanitary landfill or incinerator, as directed by State and local authorities, by burning, if allowed, out of site.

NOTICE OF WARRANTY

SCHERING-PLOUGH ANIMAL HEALTH CORPORATION MAKES NO WARRANTY OF MERCHANTABILITY, FITNESS FOR ANY PARTICULAR PURPOSE, OR OTHERWISE EXPRESSED OR IMPLIED, CONCERNING THIS PRODUCT OR ITS USES WHICH EXTEND BEYOND THE USE OF THIS PRODUCT UNDER NORMAL CONDITIONS AND IN ACCORDANCE WITH THE STATEMENTS ON THIS LABEL.

SP Schering-Plough Animal Health
Copyright © 1981, Schering-Plough Animal Health Corp.,
1500 Morris Ave., Kenilworth, NJ 07033.
All rights reserved. 23416409 Rev. 9/79

ACCEPTED
JAN 28 2002
Under the Federal Insecticide, Fungicide, and Rodenticide Act as amended, for law purposes registered under EPA Reg. No. 2-13-82