

773-84

03/01/2006

1/3

FRONT PANEL

ULTRA BOSS®

Pour-On Insecticide For Cattle, Sheep, Goats and Horses
Take Control of Flies and Lice
Season-Long Lice Control with One Application†
Squeeze 'N' Measure™ Bottle
May Be Used With Automatic Dosing System

ACCEPTED
MAR 1 2006
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg. No. 773-84

ACTIVE INGREDIENTS

Permethrin (3-Phenoxyphenyl) methyl (±) cis, trans-3-(2,2-dichloroethenyl) -2,2-dimethylcyclo-propanecarboxylate*	5.00%
Piperonyl Butoxide Technical**	5.00%
OTHER INGREDIENTS***	90.00%
Total.....	100.00%

*cis/trans ratio: Min. 35%(±) cis and max 65%(±) trans
**Equivalent to 4.5% (butylcarbityl) (6-propylpiperonyl) ether and 0.5% related compounds
***contains petroleum distillates

**KEEP OUT OF REACH OF CHILDREN
CAUTION
SEE SIDE PANEL FOR ADDITIONAL
PRECAUTIONARY STATEMENTS**

Net Contents: One U.S. Quart
(0.946 L)

EPA Reg. No. 773-84
EPA Est. No. 6175-LA-01



Schering-Plough Animal Health Corporation
Union, NJ 07083

Treats 57
550 lb. Cattle

NOTE: For One Gallon containers:

Net Contents: One U.S. Gallon
(3.785 L)

Reference to Squeeze 'N' Measure™ Bottle
does not apply to One Gallon container.

Treats 229
550 lb. Cattle

SIDE PANELS

DIRECTIONS FOR USE		
It is a violation of Federal law to use this product in a manner inconsistent with its labeling. READY TO USE – No dilution necessary.		
Apply To	Target Species	Dosage
Lactating and Non-Lactating Dairy Cattle; Beef Cattle; Calves; Horses (Not intended for human consumption) Lactating and Non-Lactating Dairy Goats;	Lice, Horn flies, Face flies Aids in control of Horse flies, Stable flies, Mosquitos, Black flies and Ticks	Apply 3 mL per 100 lbs. body weight of animal up to a maximum of 30 mL for any one animal. Pour along back and down face. Back Rubber Use: Mix 100 mL per gallon of #2 diesel oil or mineral oil. Keep rubbing device charged. Results improved by daily forced use. Apply 1.5 mL per 50 lbs. body weight of animal up to a maximum of 18 mL for any one animal. Pour along back and down face.
Sheep	Sheep keds, Lice	Pour along back. Apply 1.5 mL per 50 lbs. of body weight of animal up to a maximum of 18 mL for any one animal.

For cattle, horses, sheep and goats, repeat treatment as needed but not more than once every two weeks.

For horses, do not saddle horse for 6 hours after application and refrain from washing down area of application.

For optimum lice control, two treatments at 14-day intervals are recommended.

SPECIAL NOTE: Ultra Boss® Pour-On Insecticide is not effective against cattle grubs.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS CAUTION
Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling. Prolonged or frequently repeated skin contact may cause allergic reactions in some individuals.
FIRST AID
If on Skin or Clothing <ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15- 20 minutes. • Call a poison control center or doctor for treatment advice.
If swallowed <ul style="list-style-type: none"> • Call poison control center or doctor immediately for treatment advice. • Do not induce vomiting unless told to do so by the poison control center or doctor. • Do not give any liquid to the person. • Do not give anything by mouth to an unconscious person.

If in Eyes

- Hold eyes open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

HOT LINE NUMBER

Have the product container or label with you when calling the poison control center or doctor or going for treatment. You may also contact the Rocky Mountain Poison Control Center at 1-303-595-4869 for emergency medical treatment information.

Note to Physician: Contains petroleum distillates. Vomiting may cause aspiration pneumonia.

ENVIRONMENTAL HAZARDS

This pesticide is extremely toxic to fish. Use with care when applying to areas adjacent to any body of water. Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

Storage: Keep container sealed when not in use. Do not store near food or feed.

Pesticide Disposal: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

Container Disposal: Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incinerator, or if allowed by State and local authorities, by burning. If burned, stay out of the smoke.

NOTICE OF WARRANTY

Schering-Plough Animal Health Corp. makes no warranty of merchantability, fitness for any particular purpose or otherwise expressed or implied concerning this product or its uses which extend beyond the use of the product under normal conditions in accord with the statements made on this label.

† Season long lice control from January to April.