




VirKon® S

BROAD SPECTRUM DISINFECTANT

For Use in Cleaning and Disinfecting Industrial
Animal and Agricultural Facilities

EFFECTIVE AGAINST:

BACTERIA	VIRUSES OF	FUNGI
Streptococcus pyogenes Campylobacter pyloridis Klebsiella pneumoniae Escherichia coli Salmonella typhimurium Salmonella choleraesuis Pseudomonas aeruginosa Staphylococcus aureus Staphylococcus epidermidis Mycoplasma gallisepticum	Newcastle Disease Infectious Bronchitis Infectious Bursal Disease Avian Laryngotracheitis Avian Influenza Marek's Disease	Aspergillus fumigatus Aspergillus fumigatus Candida albicans

ACTIVE INGREDIENTS

Potassium peroxymonosulfate 20.4%
Sodium Chloride 1.5%

INERT INGREDIENTS 78.1%

TOTAL 100.0%

Equivalent to 9.75% Available Chlorine

**KEEP OUT OF REACH OF CHILDREN
DANGER:**

See Back Panel for Additional Precautions

11 lbs (5kg) Net Weight

DISTRIBUTED BY

AGRI-BIO®

Agri-Bio Corp., a subsidiary of A.H. Robins Company,
966 Dorsey St. Gainesville, GA 30501, phone 1-800-AGRI-BIO
(247-4246)

In Georgia call collect 404-536-0111

VirKon S is a registered trademark of and manufactured
by Antec International Limited U.S. Patent No. 4822512

VirKon.S

BROAD SPECTRUM DISINFECTANT

VIRKON S is effective against numerous microorganisms affecting animals: viruses, gram positive and gram negative bacteria, fungi (molds and yeasts), and mycoplasma. Efficacy was determined in the presence of hard water and organic material.

CONTROLS

Viruses of - Newcastle Disease, Infectious Bronchitis, Infectious Bursal Disease, Laryngotracheitis, Avian Influenza, and Marek's Disease.

Fungi (molds and yeasts) - *Aspergillus flavus*, *Aspergillus fumigatus*, and *Candida albicans*.

Bacteria - *Streptococcus pyogenes*, *Campylobacter pyloridis*, *Klebsiella pneumoniae*, *Escherichia coli*, *Salmonella typhimurium*, *Salmonella choleraesuis*, *Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Staphylococcus epidermidis*, *Mycoplasma gallisepticum*.

DIRECTIONS FOR USE

Federal law prohibits use of this product in a manner inconsistent with its labeling.

This powder formula is easily diluted for use in manual or machine operations.

VIRKON S is recommended for use in hatcheries, chicken houses, and processing plants.

HATCHERIES: VIRKON S at 1% is generally suitable for cleaning and disinfecting hatcheries, setters, evaporative coolers, humidifying systems, ceiling fans, chicken houses, transfer trucks, trays, and plastic chick boxes (use 2% for *Mycoplasma*, *Aspergillus flavus*, and *Aspergillus fumigatus*).

BROILER/BREEDER HOUSES: Following removal of chickens, feed, and water troughs, remove manure and litter and spray floors and side walls with VIRKON S at 1%. Thoroughly wash waterers and feeders with a 1% solution of VIRKON S (use 2% for *Mycoplasma*, *Aspergillus flavus*, and *Aspergillus fumigatus*). After contact for 10 minutes, rinse with potable water. Do not use to house poultry or employ equipment until treatment has dried.

PROCESSING PLANTS: For disinfecting and cleaning walls, ceilings, and floors spray at a dilution of 1%.


Virkon S is recommended for use in fogging (wet misting) operations as an adjunct either preceding or following regular cleaning and disinfecting procedures. Virkon S at 1-2% is suitable. Fog (wet mist) until the area is moist using automatic foggers according to manufacturer's use directions.

For air sanitizing: Virkon S at 1-2%; fog the atmosphere until surfaces are moist; allow at least 2 hours before entering area that has been fogged.

VirKon.S

DIUTION CHART

QUANTITY OF FINAL SOLUTION	1% SOLUTION	2% SOLUTION
American		
1 gallon	1.3 ounces	2.7 ounces
10 gallons	13.4 ounces	26.7 ounces
50 gallons	66.8 ounces	133.5 ounces
Metric		
10 liters	100 grams	200 grams
50 liters	500 grams	1000 grams
200 liters	2 kgs	4 kgs



FOR DISINFECTING use a rate of 7.5 gallons/1000 sq. ft.
For CLEANING AND DISINFECTING use a rate of 15 gallons/1000 sq. ft.

STORAGE AND DISPOSAL - Store in a cool, dry place in tightly closed container. Always replace lid after use. Wash empty container thoroughly and dispose of in trash.

PRECAUTIONARY STATEMENTS

Hazard to Humans and Domestic Animals - DANGER, Corrosive. Causes skin burns and irreversible eye damage. Harmful if swallowed, absorbed through skin, or inhaled. Do not get in eyes, on skin, or on clothing. Wear protective clothing and rubber gloves. Avoid breathing dust. Wear goggles, face shield, or safety glasses. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

Practical Treatment

IF IN EYES: Hold eyelids open and flush with a steady, gentle stream of water for 15 minutes. Get prompt medical attention.

IF ON SKIN: Wash with plenty of soap and water. Get medical attention if irritation persists.

IF INHALED: or symptoms of coughing, choking, or wheezing are troublesome, remove to fresh air and seek medical attention.

IF SWALLOWED: Drink promptly a large quantity of water. No oil alcohol. Get immediate medical attention.

NOTICE TO PHYSICIAN: Probable mucosal damage may contraindicate the use of gastric lavage.

Manufactured for
Agri-Bio Corporation, 946 Dorsey Street, Gainesville, Georgia 30501 U.S.A.
EPA Reg. No. 778-91-43490. EPA Est. No. 42482-BM-001