

CAUTION: Keep out of reach of Children. Read Directions and Cautions before using.

ACCEPTED May 13, 1971

Chlordane is toxic and Avoid inhaling dust a and goggles if necess free from residues. A animals, or in or arc Toxic to fish, birds treated areas may be

> CRABGRASS SEEDLIN areas should be app A temporary browning 15 lbs. per 7½ lbs. per > 11/2 lbs. per

To control wireworm onions and cabbage v treats 1200 to 1400 so

To control cutworms, . 1200 square feet of a moth) apply liberally to get material into I

CAUTION: Keep ch into the soil.

To control grasshopp dust directly on the !

To control cutworms, cutting ants on flower the plants. Remains 400-600 square feet.

SOLD I


Seller warrants that th the purpose stated on particular purpose, expr tions, or under abnorma buyer assumes the risk

CHLORDANE 10% DUST

CHLORDANE

• 10% DUST


CHLORDANE **10% DUST** 



KIL GARDEN AND SOIL INSECTS: FOR CONTROL OF JAPANESE BELTLE GRUBS. CUTWORMS, CRICKETS, ANTS, WHITE GRUBS. CONTROLS CRABGRASS SEEDLINGS



reach of Children. tions before using.



# DANE UST

CHLORDANE 10% DUST



TROL OF JAPANESE BELTLE GRUBS, S, WHITE GRUBS. SEEDLINGS

## CAUTION

#### HARMFUL IF SWALLOWED

Chlordane is toxic and may be absorbed through the skin in dry form or from solutions. Avoid inhaling dust and mists from sprays or getting them in the eyes. Use respirator and goggles if necessary for protection. Avoid unnecessary skin contact. Keep clothing free from residues. Avoid contamination of feeds and foodstuffs. Do not use on dairy animals, or in or around dairy buildings, or on animals being finished for slaughter.

Toxic to fish, birds and mammals. Do not contaminate fish ponds. Birds feeding on treated areas may Le harmed.

# DIRECTIONS FOR USE

CRABGRASS SEEDLINGS: MUST BE APPLIED before crabgrass seed germinates – in most areas should be applied before April 1st. WATER IN WELL down onto the soil surface. A temporary browning sometimes occurs.

15 lbs. per 1000 sq. ft. lawn area — 11¼ lbs. per 750 sq. ft. lawn area 7½ lbs. per 500 sq. ft. lawn area — 3¾ lbs. per 250 sq. ft. lawn area > 1½ lbs. per 100 sq. ft. lawn area

## FOR GARDEN SOIL

abbage work into top 2 or 3 inches of soil when preparing seed beds. I pound 1400 square feet.

# FOR LAWNS

square feet of area. For white grubs, Japanese Beetle grubs and sod webworm (lawn-inoth) apply liberally at the rate of 1 pound to 500 to 600 square feet. Water well with hose to get material into nests and living places.

CAUTION: Keep children and pets off treated lawn until insecticide has been washed into the soil.

#### FOR FOLIAGE DUST

To control grasshoppers, thrips and plant bugs on flowers, shrubs, and other ornamentals dust directly on the plant and insect. Repeat as necessary to prevent damage.

#### IN THE GARDEN

To control cutworms, earwigs, mole crickets, sow bugs, millipedes, slugs, snails and leaf cutting ants on flowers, shrubs and other ornamentals, dust liberally on the ground, around the plants. Remains effective at least 10 days. Re-Treat as necessary. 1 pound treats 400-600 square feet.

# SOLD FOR AGRICULTURAL INSECTICIDAL PURPOSES

## Do NOT store with foodstuffs

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty not any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Reg. No. 779-31

FAESY & BLADE FO