

U.S.D.A. Reg. No. 779-43

Do not reuse empty container. Destroy by burying with waste or burning. Stay away from smoke or fumes.

This product is Highly Toxic to Bees Exposed to Direct Treatment or Residues on Crops. Protective Information May Be Obtained from Your Cooperative Agricultural Extension Service.

**CAUTION: Keep out of reach of Children.**  
**Read Directions and Cautions before using.**  
2 POUNDS NET

F&B

ACCEPTED

JUL 22 1971

UNDER THE FEDERAL INSECTICIDE FUNGICIDE AND RODENTICIDE ACT FOR ECONOMIC PEST CONTROL REGISTERED UNDER REG. NO. 779-43 TO ACCEPTED COMMENTS.

**SEVIN\***  
WETTABLE POWDER

779-43

**KILLS JAPANESE BEETLES**

Bag Worms  
Birch Leaf Miner  
Flea Beetles  
Leaf Hopper  
Larvae and certain other insects on Ornamentals  
Fruits and Vegetables



**ORNAMENTAL INSECT CONTROL**

Mosses, ferns, and other plants are attractive to Japanese beetles. The beetles feed on the leaves, stems, and flowers of these plants. They also feed on the leaves of many other plants, including roses, peonies, and hydrangeas. The damage caused by Japanese beetles is often severe, and it is important to take steps to control their numbers. SEVIN is an effective insecticide for controlling Japanese beetles on ornamental plants. It is easy to use and provides long-lasting protection.

SEVIN can be used safely on a wide variety of ornamental plants, including azalea, begonia, boxwood, camellia, geranium, Chinese elm, clematis, dahlia, dogwood, elm, euonymus, gladiolus, geranium, hickory, hydrangea, iris, laurel, magnolia, orchid, privet, rhododendron, rose, yew. DO NOT USE ON BOSTON IVY.

**LAWN INSECT CONTROL**

Japanese beetles are a common pest of lawns. They feed on the grass blades, causing damage and discoloration. SEVIN is an effective insecticide for controlling Japanese beetles on lawns. It is easy to use and provides long-lasting protection. To use SEVIN on lawns, mix the powder with water and spray it onto the grass. Repeat the application every 2 to 3 weeks if necessary.

**Close to Harvest Use and Residue Tolerances**

**FRUIT INSECT CONTROL**

**VEGETABLE INSECT CONTROL**

SEVIN is an effective insecticide for controlling a wide variety of insects on fruits and vegetables. It is easy to use and provides long-lasting protection. To use SEVIN on fruits and vegetables, mix the powder with water and spray it onto the plants. Repeat the application every 7 to 10 days if necessary. SEVIN is also effective for controlling insects on ornamental plants and lawns.

FAESY & BETHOFF, INC., Edgewater, N. J.

**ACCEPTED**  
 OCT 4 1971  
 UNDER THE FEDERAL  
 FUNGICIDE AND  
 FOR ECONOMIC PEST  
 ED UNDER NO. 277-1-1-1  
 EQ ATTACHED

CAUTION

DIRECTIONS

ORNAMENTALS

LAWNS

VEGETABLES

**BEANS:** Apply to foliage and soil. For leafhoppers, apply 1/2 cup per 100 sq. ft. per day of infestation.

**CORN:** Apply to foliage if necessary for control of insects that begin to dry plants for human consumption, fodder and storage.

**CUCUMBER (SUMMER):** Beetles. Pickle beetles. Apply to foliage and soil to control their damage to plants.

**TOMATO, EGGPLANT:** Beetles. Apply to foliage and soil to control insects on the plants and including dirt on the plants.

**POTATOES:** Apply to foliage and soil to control insects on the plants per 100 sq. ft. per day of infestation. Apply 7 to 10 lbs. per 100 sq. ft.

**5% SEVIN\***  
 VEGETABLE AND GARDEN DUST

**5% SEVIN\***  
 VEGETABLE AND GARDEN DUST

**5%**  
 VEGETAB

**ANOTHER  
 F&B PLANNED  
 PACKAGE  
 NO FUSS  
 MUSS  
 WASTE**



**ANOTHER  
 F&B PLANNED  
 PACKAGE  
 NO FUSS  
 MUSS  
 WASTE**



FOUR PO

IS  
 Do not reuse  
 away from  
 ORNAMENTAL  
 Sweet Corn  
 Avoid contact  
 with eyes  
 Wash hands  
 thoroughly  
 after use  
 Keep out of  
 reach of children  
 and animals  
 For more information  
 contact your  
 local health  
 department  
 or the  
 manufacturer

Avoid contacting  
 supplies, street  
 additional caut

CAUTION  
 Read Dire

ANOTHER  
**F&B** PLANNED  
 PACKAGE  
**NO** FUSS  
 MUSS  
 WASTE



Begin to apply... for human consumption... toddler and furego...  
**CUCUMBER, SUMMER CUCURBIT, CABBAGE, SQUASH, PEAS, BEANS, TOMATOES, EGGPLANT, PEPPERS, POTATOES, GRAPES:** Beetles, Pickleworm, Hornworm, Stem Borer, and other insects or their damage...  
**POTATOES:** Apply... apply 1/2 to 3/4 cup per 1000 sq. ft. For every 7 to 10 days...  
**GRAPES:** For control of first brood... control of second and third broods... applied up to and including day of harvest...  
 Seller warrants that the product conforms to its... the purpose stated on the label... of use, but neither this particular product... buyer assumed to be...

USDA REG. NO. 119-18

FAESY & BESTHOFF, INC.  
 EDGEWATER, N. J. 07020

ANOTHER  
**F&B** PLANNED  
 PACKAGE  
**NO** FUSS  
 MUSS  
 WASTE



**FOUR POINT**  
 ACTIVE INGREDIENT  
 Permethrin (1-naphthyl piperonyl butyrate)  
 INERT INGREDIENT  
 Total . . . . .  
 Net Wt. of C. C. Co. (1-naphthyl N-methyl piperonyl butyrate)

FOR INSECT CONTROL



OCT 24 1971  
 UNDER THE FEDERAL INSECTICIDE ACT  
 FUNGICIDE AND PESTICIDE ACT  
 FOR ECONOMIC PEST CONTROL  
 REGISTERED UNDER NO. 779-45757  
 EQ ATTACHED

**CAUTION**

**DANGEROUS IF INHALED OR SWALLOWED.** Do not breathe dust. Do not get internally. Skin contact with any insecticide may be harmful. Avoid unnecessary contact. Wash hands before eating. Take shower or bath after use. Wear regular long sleeved work clothing. Change to clean clothing

For Physician: Carbaryl is a moderate reversible cholinesterase inhibitor. Atropine is antidote.

KEEP OUT OF REACH OF CHILDREN AND ANIMALS

**DIRECTIONS**

Apply at first sign of insect or damage and repeat on a regular 5 day schedule or as necessary for good insect control.

**ORNAMENTALS**

Control of Apple Aphid, Bagworm, Birch Leaf Miner, Blister Beetle, Boxelder Bug, Elm Leaf Miner, Elm Leaf Aphid, Elm Leaf Beetle, Flea Beetle, Japanese Beetle, Leafhopper, Leaf Roller, Oak Leaf Miner, Orange Tortrix, Periodical Cicada, Psyllid, Rose Slug, Rose Aphid, Scale insects in the crawler stage, Tent Caterpillar, Willow Leaf Beetle and exposed Thrips — at first sign of damage apply to infested plants. Scale Control — apply in Spring and early Summer when scales are present. **DO NOT USE ON BOSTON IVY.**

**LAWNS**

Apply 1/2 cup per 1000 sq. ft. for control of Leafhopper, Fall Armyworm, Sod Webworm. Results apply after rain or watering and do not water for at least 2 days after application. Repeat in 2 to 3 weeks if necessary.

**VEGETABLES**

Apply thoroughly to upper and lower leaf surfaces at first sign of insect damage before feeding occurs. For Mexican Bean Beetle apply 1/2 cup per 1000 sq. ft., for Flea Beetles apply 1/4 cup per 1000 sq. ft.; for Lygus and Stink Bugs apply 1 1/2 cups per 1000 sq. ft. If needed, repeat applications every 7 to 10 days up to and including day of harvest.

For control of Corn Earworm use 1/4 to 1 1/2 cups per 1000 sq. ft. Repeat as needed for insects attacking silks and ears; apply at 2 to 3 day intervals until silks are dry. May be applied up to and including day of harvest on sweet corn ears for consumption. Do not apply within 7 days before harvest on corn for use as forage for dairy and meat animals.

**FOR SUMMER SQUASH:** For control of Squash Bugs, Cucumber Beetles, Flea Beetles and Melonworm — apply 3/4 cup per 1000 sq. ft. when insects or damage appear. Repeat at 7 to 10 day intervals or as necessary. May be applied up to and including day of harvest.

**FOR PLANT PROTECTOR:** For control of Tomato Fruitworm, Fall Armyworm, Flea

**ORNAMENTAL USE:** Be sure to cover upper and lower leaf surfaces thoroughly.

**SWEET CORN USE:** (See center-rear panel); Application to tassel zone of corn during pollen shed period will seriously reduce bee population.

"This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service."

See rear-panel for actual names of insects controlled by this product. One cup of 5% Sevin weighs about 8 ounces.

**CAUTION**

Avoid contamination of food, animals, feed, water supplies, streams and ponds. See rear-panel for additional cautions.



**CAUTION: Keep out of reach of Children. Read Directions and Cautions before using.**

**5% SEVIN\* VEGETABLE AND GARDEN DUST**

**5% SEVIN\* VEGETABLE AND GARDEN DUST**

ANOTHER



**FOUR POUNDS NET**

ACTIVE INGREDIENT:  
 Carbaryl (1-naphthyl N-methylcarbamate) . . . . . 5.00%  
 INERT INGREDIENTS . . . . . 95.00%

