



CHLORDANE 10% GRANULAR

ACTIVE INGREDIENT:

*Technical Chlordane _____

INERT INGREDIENTS: _____

*equivalent to 6% Octachloro-4,7-Methanotetrahydroindane and 4% related compounds

FOR CRABGRASS CONTROL AND INSECT CONTROL OF CERTAIN SOIL-INHABITING INSECTS

RECOMMENDATIONS

CRABGRASS CONTROL—Apply 15 lbs. per 1,000 sq. ft. of lawns and ornamental turf to suppress crabgrass seedlings before germination or sprouting occurs. It is for use as an early season preventative of established crabgrass plants. Timing of application is important and should be applied before any germination which in many areas is during the period April 15 to May 1. This treatment will not burn or injure established lawns.

DO NOT USE ON NEWLY SEEDED LAWNS

SPOT TREATMENT: Late winter application to infested patches may be as effective as spring application. For spot treatment apply 1 1/2 lbs. per 100 sq. ft.

HELPFUL HINTS: After crabgrass seedlings have been suppressed by the Chlordane treatment, control may be assisted by:

1. Mowing the lawn to 1 1/2 to 2 inches.
2. Using an adequate quantity of fertilizer to maintain active growth.

After treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children or pets.

CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass and into the soil by drenching or sprinkling with water and grass has completely dried.

each 1,000 sq. ft.) to cover surface uniformly, using a lawn fertilizer spreader. Treatment should be applied in the Spring or when insects are observed to be troublesome. Do not treat soil when it is wet. Apply before rain or sprinkling.


INSECT CONTROL ON HOME LAWNS: For the control of the above listed insects on home lawns apply 50 lbs. per acre (1 1/4 lbs. per 1,000 sq. ft.) to cover surface uniformly, using a lawn fertilizer spreader. Treatment should be applied in the spring, or when insects are observed to be troublesome. Immediately after treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children and pets. Do not treat when soil is wet. Apply before rain or sprinkling.

CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water and grass has completely dried. Do not reuse bag. Destroy by burning. Stay away from smoke.

CAUTION

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash with soap and water after exposure. Avoid contamination of feed and foodstuffs.

Toxic to Fish & Wildlife. Keep out of lakes, streams, ponds. Do not apply where runoff can occur. Do not contaminate



assisted by:

1. Mowing the lawn to 1 1/2 to 2 inches.
2. Using an adequate quantity of fertilizer to maintain active growth.

After treatment, drench lawn with water or sprinkle heavily to make treatment more effective and to reduce the hazard to children or pets.

CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass and into the soil by drenching or sprinkling with water and grass has completely dried.

INSECT CONTROL ON TURF, NURSERY AND GREENHOUSE SOILS: For the control of Ants, Armyworms, Boxelder Bugs, Chiggers, Chinch Bugs, Crickets, European Earwigs, Grasshoppers, Japanese Beetle Grubs, Rose Chafer, Sod Webworm, Spiders, Ticks, White-fringed Beetle, White Grubs and Wireworms in Turf (such as golf courses, athletic fields, large estate nurseries, cemeteries and park sod) and in nursery and greenhouse soils where ornamental plants are grown, use at the rate of 50 lbs. per acre (1 1/4 lbs. to

CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water and grass has completely dried. Do not reuse bag. Destroy by burning. Stay away from smoke.

CAUTION

Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash with soap and water after exposure. Avoid contamination of feed and foodstuffs.

Toxic to Fish & Wildlife. Keep out of lakes, streams, ponds. Do not apply where runoff can occur. Do not contaminate water by cleaning equipment or waste disposal. Birds feeding in treated areas may be killed.

DO NOT STORE WITH FOODSTUFFS

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CHLORDANE 10% GRANULAR

SPREADER

APPROXIMATE SPREADER SETTING

SCOTT
CYCLONE

CRABGRASS CONTROL	LAWN INSECT CONTROL
9 1/2	3 3/4
6 1/4	2 3/4

NOTICE: Variables inherent in equipment and materials, as well as accuracy of machine adjustment, may cause actual application rates to deviate slightly. Therefore, Faesy & Besthoff, Inc. makes no warranty as to the uniformity of coverage obtained from the setting listed above.

CAUTION: Keep Out of Reach of Children

Read Directions and Cautions before Using

E.P.A. Reg. #779-49

172

FAESY & BESTHOFF, INC.
EDGEWATER, N.J. 07020
25 LBS NET