

FLY CONTROL AROUND BUILDINGS YARDS OUT OF DOORS For Residual Spray mix 1 cup in 2 1/2 gals. water. Spray where flies alight or congregate at rates of 1 gal. per 1000 sq. ft. on painted surface or 2 gals. per 1000 sq. ft. on unpainted areas. 1 cup of sugar may be added to act as a bait to attract flies. Make regular removal of materials such as manure, garbage, and other waste products, where flies breed. NOTE This product may cause spotting on automobile paint finish. Cars should not be sprayed directly. If accidental exposure occurs, the car should be washed immediately.

CAUTION

Harmful if swallowed. AVOID prolonged breathing of spray mist, repeated contact with skin, contamination of feed and foodstuffs. Wash thoroughly after using.

If poisoning should occur, CALL A PHYSICIAN. Meanwhile, take 1 or 2 1/2 100 grain atropine tablets; repeat in 1 hour if necessary. NOTE TO PHYSICIANS Malathion upon repeated prolonged, careless use may cause cholinesterase inhibition. Atropine is antidotal.

Do not use or store near heat or open flame.

Do not use in dwellings or rooms where edible products are stored or handled.

Highly toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Reg. No. 779-66

1 FLUID QUART NET



CYTHION

50% SPRAY

**The Premium Grade Malathion
FOR USE IN AND AROUND
HOME ORCHARDS AND GARDENS**

ACTIVE INGREDIENTS:

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) -----	50.9%
Aromatic Petroleum derivative solvent ---	44.1%
INERT INGREDIENTS -----	5.0%

CAUTION:
Keep out of Reach of Children. See Side Panel for Additional Cautions.

Contains 4.5 lbs. Malathion per gallon.
*Reg. T.M. American Cyanamid Co.

Contains 4.5 lbs. Malathion per gallon.
*Reg. T.M. American Cyanamid Co.

FAESY & BESTHOFF, INC., Edgewater, N. J. 07020

Max. quantities per plant completely

Teaspoonful

1	Apples and
1 1/2	Apples and
1 1/2	Pears, pe
1 1/2	Grapes
1 1/2	Strawberry
2	Peaches
2	Apples and
1	Peppers an
1 1/2	Tomato and
1 1/2	Peas
1 1/2	Beans
1 1/2	Cabbage, B
1 1/2	aphids
1 1/2	Onion
2	Lettuce
2	Spinach

ORNAMENTS

1	Oyster shel
1 1/2	Aphids, me
	(adults)
2	Birch leafh
	young s

NOTE For best kill crawl

CAUTION Injury and canker; ju injury has also; petunias, small drought and d slight damage t

PRECAUTIONS
within 1 day of plants, broccoli, cabbage, cauliflower 7 days of harvest within 7 days of vines are not to applied on day the northeast varieties when

Oct 30, 1969

779-66

FLY CONTROL AROUND BUILDINGS, YARDS - OUT OF DOORS: For Residual Spray — mix 1 cup in 2½ gals. water. Spray where flies alight or congregate at rates of 1 gal. per 1000 sq. ft. on painted surface or 2 gals. per 1000 sq. ft. on unpainted areas. 1 cup of sugar may be added to act as a bait to attract flies. Make regular removal of materials such as manure, garbage, and other waste products, where flies breed. **NOTE:** This product may cause spotting on automobile paint finish. Cars should not be sprayed directly. If accidental exposure occurs, the car should be washed immediately.

CAUTION

Harmful if swallowed! AVOID prolonged breathing of spray mist; repeated contact with skin; contamination of feed and foodstuffs. Wash thoroughly after using.

If poisoning should occur, CALL A PHYSICIAN. Meanwhile, take 1 or 2 1/100 grain atropine tablets; repeat in 1 hour if necessary. **NOTE TO PHYSICIANS:** Malathion, upon repeated, prolonged, careless use may cause cholinesterase inhibition. Atropine is antidotal.

Do not use or store near heat or open flame.

Do not use in dwellings or rooms where edible products are stored or handled.

Highly toxic to fish. Do not contaminate any body of water by direct application, cleaning of equipment or disposal of wastes and containers.

NOTICE

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

USDA Reg. No. 779-66

Oct 30, 1969

779-66

1 FLUID QUART NET



CYTHION

50% SPRAY

**The Premium Grade Malathion
FOR USE IN AND AROUND
HOME ORCHARDS AND GARDENS**

ACTIVE INGREDIENTS:
 Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) ----- 50.9%
 Aromatic Petroleum derivative solvent ---- 44.1%
INERT INGREDIENTS ----- 5.0%

CAUTION: Keep out of Reach of Children. See Side Panel for Additional Cautions.

Contains 4.5 lbs. Malathion per gallon.
*Reg. T.M. American Cyanamid Co.

FAESY & BESTHOFF, INC., Edgewater, N. J. 07020

Mix quantities of plants completely

Teaspoonful

- 1 Apples and
- 1½ Apples and
- 1½ Pears
- 1½ Grapes
- 1½ Strawberries
- 2 Peaches
- 2 Apples and
- 1 Peppers and
- 1½ Tomato and
- 1½ Peas — appl
- 1½ Beans — M
- 1½ Cabbage, B

- aphids
- 1½ Onions — t
- 2 Lettuce
- 2 Spinach —

ORNAMENT

- 1 Oyster shel
- 1½ Aphids, me
- (adults).
- 2 Birch leafm
- young sc

NOTE For best kill craw

CAUTION: Injury and canaerti ju injury has also l petunias, small-l drought and di slight damage to

PRECAUTIONS: within 1 day of plants, broccoli, cabbage, cauliflo 7 days of harve within 7 days of vines are not to applied on day the northeast, varieties when t

F&B

1 FLUID QUART NET

CYTHON

50% SPRAY

The Premium Grade Malathion
FOR USE IN AND AROUND
HOME ORCHARDS AND GARDENS

ACTIVE INGREDIENTS:

Malathion (0,0-dimethyl dithiophosphate of diethyl mercaptosuccinate) ----- 50.9%

Aromatic Petroleum derivative solvent ---- 44.1%

INERT INGREDIENTS ----- 5.0%

CAUTION:

Keep out of Reach of
Children. See Side Panel
for Additional Cautions.

Contains 4.5 lbs.
Malathion per gallon.

*Reg. T.M. American
Cyanamid Co.

FAESY & BESTHOFF, INC., Edgewater, N. J. 07020

DIRECTIONS FOR USE

Mix quantities given in table with 1 gal. of water and spray to wet plants completely without run off

Teaspoonful

- 1 Apples and Pears wooly aphid and bud moth
- 1 1/2 Apples and Pears aphid and mealy bugs
- 1 1/2 Pears pear psylla
- 1 1/2 Grapes spider mites and mealy bugs
- 1 1/2 Strawberries aphids and mites
- 2 Peaches — spider mites, oriental fruit moth
- 2 Apples and Pears leaf rollers, mites and codling moth
- 1 Peppers and Potatoes aphids
- 1 1/2 Tomato and Eggplant spider mites and aphids
- 1 1/2 Peas - aphids
- 1 1/2 Beans - Mexican bean beetle and spider mites
- 1 1/2 Cabbage, Broccoli, Cauliflower, Brussels Sprouts, Turnip aphids
- 1 1/2 Onions - thrips
- 2 Lettuce - aphids, mites, leafhoppers
- 2 Spinach - aphids

ORNAMENTALS

- 1 Oyster shell scale, lace bugs
- 1 1/2 Aphids, mealy bugs, spider mites, whitefly, Japanese beetle (adults), leafhopper, tarnished plant bug, thrips
- 2 Birch leafminer, boxwood leafminer, bagworm, tent caterpillar, young scales such as magnolia, azalea juniper

NOTE For best results with scale insects, apply in May and June to kill crawlers and again one month later

CAUTION: Injury may occur on ferns, hickory, viburnum, lantana, crassula and canaert juniper following the use of emulsifiable liquid. Slight injury has also been reported on Boston, Pteris, and Maidenhair ferns, petunias, small-leaf spirea, white pine and maples. Under extreme heat, drought and disease conditions the emulsifiable liquids may cause slight damage to elms

PRECAUTIONS: Make no application on beans, pears and tomatoes within 1 day of harvest; on apples, grapes, strawberries, peppers, egg plants, broccoli, turnips and onions within 3 days of harvest; on peaches, cabbage, cauliflower, brussels sprouts, spinach and head lettuce within 7 days of harvest; on leaf lettuce within 14 days of harvest; on peas within 7 days of harvest if vines are to be used for animal feeds. If vines are not to be fed, within 3 days of harvest, on potatoes, may be applied on day of harvest. Injury may occur on Bosc pears grown in the northeast, and on grapes of Ribier, Italia, Cardinal and Almeria varieties when this product is applied after the clusters appear

DOORS: For
re flies alight
ed surface or
sugar may be
al of materials
e flies breed.
paint finish.
osure occurs.

ray mist; re-
lstuffs. Wash

take 1 or 2
ry. NOTE TO
less use may

are stored or

ater by direct
nd containers.

al description
el when used
of use, but
hantability or
xtends to the
der abnormal
ble to seller.

1969

79-66