


TO OPEN BAG, LIFT  
END OF PAPER FILTER  
CORD AND PULL  
EAM-Y-AIDE




CONTENTS OF THIS BAG WILL  
TREAT 2,500 SQUARE FEET  
(50' x 50')



# LAWN AND TURF FUNGICIDE

Controls Many  
Grass Diseases  
Contains DYRENE<sup>®</sup> and MANEB with Zinc



## LAWN & TURF FUNGICIDE

Contains Dyrene<sup>®</sup> and Maneb with Zinc

A Broad Spectrum Fungicide for Prevention and Control of Most Turf Diseases Such as:

SNOW MOLD, (Typhula Blight and Fusarium Patch), RUST, Sclerotinia DOLLAR SPOT, COPPER SCOT, RED THREAD, Rhizoctonia BROWN PATCH, Fusarium Roseum Blight, and Helminthosporium induced diseases (Helminthosporium Blight in Fescue, and Helminthosporium LEAF SPOT in Fescue, Bent & Blue Grass, WILTING OUT in Kentucky Blue Grass, RED LEAF SPOT in Bent Grass and BROWN BLIGHT in Ryegrass, ZONATE EYESPOT in Fescue)

### DIRECTIONS FOR USE

A preventive or an eradicant treatment for the diseases listed above will give best results if application is made to a slightly moistened lawn. Application after a light shower or watering on grass blades is therefore recommended. DO NOT APPLY WHEN RAINFALL IS EXPECTED WITHIN 24 HOURS AND DO NOT WATER LAWN FOR 24 HOURS AFTER TREATMENT.

**PREVENTIVE USE:** Warm weather preventive treatments can be started from early Spring to early Summer, as previous disease experience dictates, and repeated at 10 to 14 day intervals as long as cool, wet weather continues or until improvement is noted. (Set spreader at preventive rate—4 pounds per 1000 square feet.)

**ERADICANT USE:** If lawn disease is already prevalent, make application at 10 day intervals until improvement is noted. (Set spreader at eradicant rate—6 pounds per 1000 square feet.)

**SNOW MOLD USE:** Apply at eradicant rate (6 pounds per 1000 square feet) after first killing frost and again in midwinter when ground is free of snow.

### CAUTION

KEEP OUT OF REACH OF CHILDREN

Hazard if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Wash thoroughly after use. Avoid contamination of feed, foodstuffs, streams, lakes, or fishponds. Keep children and pets off treated areas until material has washed into soil and grass is dry. Do not reuse empty container. Destroy it by burning. Stay away from smoke and fumes.

### ANTIDOTE

**INTERNAL:** Give tablespoon of mustard in glass of water. Get medical attention.

### APPROXIMATE SPREADER SETTINGS

	PREVENTIVE	ERADICANT
SCOTT	6	7
LAWN BEAUTY	7	8

ACCEPTED

5-11-70  
779-68

UNDE

# LAWN AND TURF FUNGICIDE

## Controls Many Grass Diseases

Combines DYRENE® and MANEB with  
Zinc in Granular Form for Easy Application

### WARNING

**KEEP OUT OF REACH OF CHILDREN  
READ ALL DIRECTIONS BEFORE USING**

See Rear Panel for Additional Precautions

<b>ACTIVE INGREDIENTS:</b>	
2,4-Dichloro-6-(o-chlorophenyl)-s-triazine .....	4.0%
Maneb (Manganese Ethylene Bisdithiocarbamate) .....	4.0%
Zinc, as Metallic .....	0.1%
<b>INERT INGREDIENTS:</b> .....	91.9%

**10 LBS. NET. WT.**

**FAESY & BESTHOFF, INC.**

Edgewater, N.J. 07020

U.S.D.A. Reg. No. 779-68  
DYRENE is a Registered Trademark in U.S. Patent Office by Ciba Corp.  
U.S. Patent No. 2,720,489

ACCEPTED

5-11-70  
779-68

UNDER THE FEDERAL INSECTICIDE ACT  
FUNGICIDE AND FERTILIZER ACT  
FOR ECONOMIC CONTROL OF WEEDS  
AND UNDER NO. 100-100000-1  
ATTACHED TO SUBJECT

induced diseases (Rhynchosporium Blight in Fescue, and Helminthosporium LEAF SPOT in Fescue, Bent & Blue Grass, MELTING OUT in Kentucky Blue Grass, RED LEAF SPOT in Bent Grass and BROWN BLIGHT in Ryegrass, ZONATE EYESPOT in Fescue.

### DIRECTIONS FOR USE

A preventive or an eradicant treatment for the diseases listed above will give best results if application is made to a slightly moistened lawn. Application after a light shower or when dew is on grass blades is therefore recommended. DO NOT APPLY WHEN RAINFALL IS EXPECTED WITHIN 24 HOURS AND DO NOT WATER LAWN FOR 24 HOURS AFTER TREATMENT.

**PREVENTIVE USE:** Warm weather preventive treatments should be started from early Spring to early Summer, as previous disease experience dictates, and repeated at 10 to 14 day intervals as long as cool, wet weather continues or until improvement is noted. (Set spreader at preventive rate—4 pounds per 1000 square feet.)

**ERADICANT USE:** If lawn disease is already prevalent, make application at 10 day intervals until improvement is noted. (Set spreader at eradicant rate—6 pounds per 1000 square feet.)

**SNOW MOULD USE:** Apply at eradicant rate (6 pounds per 1000 square feet) after first killing frost and again in midwinter when ground is free of snow.

### CAUTION

#### KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed or inhaled. Avoid contact with eyes, skin, or clothing. Wash thoroughly after use. Avoid contamination of feed, foodstuff, streams, lakes, or ponds. Keep children and pets off treated areas until material has washed into soil and grass is dry. Do not reuse empty container. Destroy it by burning. Stay away from smoke and fumes.

### ANTIDOTE

**INTERNAL:** One tablespoon of mustard in glass of water. Get medical attention.

### APPROXIMATE SPREADER SETTINGS

	PREVENTIVE	ERADICANT
SCOTT .....	6	7
LAWN BEAUTY .....	7	8
CYCLONE .....	4	5
SEARS .....	7½	8½

**NOTICE:** Normal differences in equipment and materials, as well as condition of machine, adjustment, and walking speed may cause variation in application rate. Therefore, Faesy & Besthoff, Inc. makes no warranty as to the uniformity of coverage obtained from the above spreader settings. Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

**10 LBS. NET. WT.**

**FAESY & BESTHOFF, INC.**

Edgewater, N.J. 07020

STACK AIDE

FRONT SHIPPING BAG  
of  
FAESY & BESTHOFF, INC.  
Edgewater, N.J. 07020

CS