



7-22-70

Treats 1500 Sq. Ft. (30' x 50')

F & B BROAD SPECTRUM LAWN INSECTICIDE SPRAY

ASPON* - CHLORDANE 1.03 - 1.18 - EC*

WARNING

May be fatal if swallowed. Avoid breathing spray mist. Do not get in eyes, on skin or on clothing. In case of contact, immediately flush eyes with plenty of water. For skin, wash with soap and water. Avoid contamination of feed and food. Avoid drift to adjoining food and forage crops. Wash off.

This product is toxic to fish and wildlife. Birds feeding on treated areas may be killed. Keep out of any body of water. Apply this product only as specified on this label.

GENERAL USE PRECAUTIONS

Clean up all spills on walks or patios. Aspon* - Chlordane 1.03-1.18-EC should be used only for recommended purposes at the recommended rates. Do not allow children or pets on treated areas until this material has been washed into the soil and the lawn is dry.

Crush and burn empty container.

F & B Warranty

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

U.S.D.A. Reg. No. 77172

NE 987

KILLS CHINCH BUGS, SOD WEBWORMS, ANTS, JAPANESE BEETLE LARVAE, LAWN MOTHS, CUTWORMS, ARMYWORMS, Harvester Ants, Leaf Cutting Ants, Fire Ants, Asiatic Garden Beetle Larvae, Boxelder Bugs, Chiggers, Earwigs, European Chafer Larvae, Green June Beetle Larvae, Mosquitoes, Adult Ticks, White Grubs, Wireworms, Mole Crickets, Spiders, Centipedes.

ACTIVE INGREDIENTS:

O, O, O, O - tetrapropyl dithiopyrophosphate	12.7%
Technical Chlordane (1)	14.6%
Aromatic Petroleum Solvent	69.5%
INERT INGREDIENTS	3.2%

* Aspon is a trademark of Stauffer Chemical Company for an insecticide. Total 100.0%

(1) Equivalent to 876 octachloro-4,7-methanotetrahydrochlordane and 584 insecticidally active related compounds.

**WARNING KEEP OUT OF THE REACH OF CHILDREN
SEE SIDE PANEL FOR ADDITIONAL CAUTIONS**

1 Fluid Quart Net

FAESY & BESTHOFF, INC. EDGEWATER, N. J. 07020

Lawns and Ornamental Turf: Wet lawn thoroughly before spraying. Use at the rate of 10 1/2 ounces (1 1/2 cupfuls) to 15 gallons of water per 500 square feet of lawn (1 quart covers 1,500 square feet). Apply with pressure sprayer, sprinkler can, or garden hose gun. Cover area in two directions to insure complete coverage.

Consult State Agricultural Experiment Station or State Extension Service Specialists for timing of application for your region.

Aspon* Chlordane may be used in accordance with the directions given above during the active growing season to control the following insects:

Chinch Bugs, Sod Webworms, Ants, Japanese Beetle Larvae, Lawn Moths, Cutworms, Armyworms, Harvester Ants, Leaf Cutting Ants, Fire Ants, Asiatic Garden Beetle Larvae, Boxelder Bugs, Chiggers, Earwigs, European Chafer Larvae, Green June Beetle Larvae, Mosquitoes, Adult Ticks, White Grubs, Wireworms, Mole Crickets, Spiders, Centipedes.

In cooler climates, state areas, make application in June or when insects first appear. Should give protection for an entire season. Due to the very long growing season, a 2, 3, or 4 applications may be necessary at 6 week intervals beginning in April or May to protect a lawn for the entire season.



7-22-70

Treats 1500 Sq. Ft. (30' x 50')

F & B BROAD SPECTRUM LAWN INSECTICIDE SPRAY

ASPON* - CHLORDANE 1.03 - 1.18 - EC*

WARNING

Allowed. Avoid breathing spray mist. Do not get in eyes, clothing, or skin. In case of contact, immediately flush eyes with water. If on skin, wash with soap and water. Avoid contamination of food and forage crops. Wash off.

Toxic to fish and wildlife. Birds feeding on treated areas should be kept out of any body of water. Apply this product only as directed.

GENERAL USE PRECAUTIONS

Use on walks or patios. **Aspon*-Chlordane 1.03-1.18-EC** is for recommended purposes at the recommended rates. Do not allow children or pets on treated areas until this material has been washed and the lawn is dry. Store in original container.

F & B Warranty

This product conforms to its chemical description and to the purpose stated on the label when used in accordance with the instructions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose or implied, extends to the use of this product under conditions, or under abnormal conditions, or under conditions reasonably foreseeable to seller, and buyer assumes the

KILLS CHINCH BUGS, SOD WEBWORMS, ANTS, JAPANESE BEETLE LARVAE, LAWN MOTHS, CUTWORMS, ARMYWORMS, Harvester Ants, Leaf Cutting Ants, Fire Ants, Asiatic Garden Beetle Larvae, Boxelder Bugs, Chiggers, Earwigs, European Chafer Larvae, Green June Beetle Larvae, Mosquitoes (Adult), Ticks, White Grubs, Wireworms, Mole Crickets, Spiders, Centipedes.

ACTIVE INGREDIENTS:

O, O, O, O - tetrapropyl dithiopyrophosphate	12.7%
Technical Chlordane (1)	14.6%
Aromatic Petroleum Solvent	69.5%
INERT INGREDIENTS	3.2%

* Aspon is a trademark of Stauffer Chemical Company for an insecticide. **Total 100.0%**

(1) Equivalent to 8.76% — octachloro-4, 7-methanotetrahydroindane and 5.84% insecticidally active related compounds.

WARNING KEEP OUT OF THE REACH OF CHILDREN SEE SIDE PANEL FOR ADDITIONAL CAUTIONS

1 Fluid Quart Net

FAESY & BESTHOFF, INC. EDGEWATER, N. J. 07020

Lawns and Ornamental Turf: Wet lawn thoroughly at the rate of 10½ ounces (1½ cupfuls) to 15 ounces per 1,000 square feet of lawn (1 quart covers 1,500 square feet) using a pressure sprayer, sprinkler can, or garden hose according to directions to insure complete coverage.

Consult State Agricultural Experiment Station or Extension Specialists for timing of application for your region.

Aspon*-Chlordane may be used in accordance with directions above during the active growing season to control lawn insects.

Chinch Bugs, Sod Webworms, Ants, Japanese Beetle Larvae, Cutworms, Armyworms, Harvester Ants, Leaf Cutting Ants, Garden Beetle Larvae, Boxelder Bugs, Chiggers, Earwigs, European Chafer Larvae, Green June Beetle Larvae, Mosquitoes (Adult), Ticks, White Grubs, Wireworms, Mole Crickets, Spiders, Centipedes.

In cooler (Mid Atlantic state) areas a single application of insects first appear should give protection for a long season. In the very long growing season (Florida, Southern states) a second application may be necessary at 6 week intervals beginning in late May or early June. Water a lawn for the entire season.