

CHLORDANE 5% GRANULAR

ACTIVE INGREDIENT:		5%
*Technical Chlordane	_____	95%
INERT INGREDIENTS:	_____	100%

*equivalent to 3% Octachloro-4,7-Methasotetrahydroindane and 2% related compounds

FOR THE CONTROL OF CERTAIN SOIL-INHABITING INSECTS

RECOMMENDATIONS

INSECT CONTROL ON TURF (Golf Courses, Athletic Fields, Large Estate Nurseries, Cemeteries and Parks), NURSERY AND GREENHOUSE SOILS:

For the control of Ants, Armyworms, Chiggers, Chinch-bugs, Crickets, Grasshoppers, Rose Chafers, Spiders and Ticks — apply uniformly at the rate of 100 lbs. per acre (2½ lbs. per 1000 sq. ft.) using a lawn fertilizer spreader. Treatment should be made in the spring or when insects are observed to be troublesome. Apply before rain or sprinkling. Do not treat when soil is wet.

For the control of Sod Webworms, Japanese Beetle Larvae, White Grubs and Wireworms — apply at the rate of 100 lbs. per acre (2½ lbs. per 1000-sq. ft) using a lawn fertilizer spreader. Apply just before a rain or use irrigation or sprinklers to wash the material into the top few inches of soil or into established turf. Do not treat when soil is wet.

For the control of Earwigs apply at the rates listed above to infested outside areas — especially along fences, walks and buildings.

For the control of Box Elder Bugs — make applications at the rate of 100 lbs. per acre (2½ lbs. per 1000 sq. ft.) to the ground where the insects crawl. Water into the established turf.

CAUTION: Do not permit children or pets to go on treated lawn until the insecticide has been washed off the grass into the soil by drenching or sprinkling with water and grass has completely dried. Do not reuse bag. Destroy by burning. Stay away from smoke. Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash with soap and water after exposure. Avoid contamination of feed and foodstuffs.

Toxic to Fish & Wildlife. Keep out of lakes, streams, ponds. Don't apply where runoff can occur. Don't contaminate water by cleaning equipment or waste disposal. Birds feeding in treated areas may be killed.

SPREADER	APPROXIMATE SPREADER SETTINGS
SCOTT	4½
CYCLONE	4

NOTICE: Variables inherent in equipment and materials, as well as accuracy of machine adjustment, may cause actual application rates to deviate slightly. Therefore, F&B & Besthoff, Inc. makes no warranty as to the uniformity of coverage obtained from the settings listed above.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to seller and buyer assumes the risk of any such use.

CAUTION: Keep Out of Reach of Children
Read Directions and Cautions before Using

APPROVED
2-7-72
278-79

AND GREENHOUSE SOILS:

For the control of Ants, Armyworms, Chiggers, Chinch-bugs, Crickets, Grasshoppers, Rose Chafers, Spiders and Ticks — apply uniformly at the rate of 100 lbs. per acre (2½ lbs. per 1000 sq. ft.) using a lawn fertilizer spreader. Treatment should be made in the spring or when insects are observed to be troublesome. Apply before rain or sprinkling. Do not treat when soil is wet.

For the control of Sod Webworms, Japanese Beetle Larvae, White Grubs and Wireworms — apply at the rate of 100 lbs. per acre (2½ lbs. per 1000 sq. ft.) using a lawn fertilizer spreader. Apply just before a rain or use irrigation or sprinklers to wash the material into the top few inches of soil or into established turf. Do not treat when soil is wet.

For the control of Earwigs apply at the rates listed above to infested outside areas — especially along fences, walks and buildings.

For the control of Box Elder Bugs — make applications at the rate of 100 lbs. per acre (2½ lbs. per 1000 sq. ft.) to the ground where the insects crawl. Water into the established turf.

Put the grass into the sun by spreading on a dry water and grass has completely dried. Do not reuse bag. Destroy by burning. Stay away from smoke. Harmful if swallowed. Avoid breathing of dust. Avoid contact with skin. Wash with soap and water after exposure. Avoid contamination of feed and foodstuffs.


Toxic to Fish & Wildlife. Keep out of lakes, streams, ponds. Don't apply where runoff can occur. Don't contaminate water by cleaning equipment or waste disposal. Birds feeding in treated areas may be killed.

SPREADER	APPROXIMATE SPREADER SETTINGS
SCOTT	4 ½
CYCLONE	4

NOTICE: Variables inherent in equipment and materials, as well as accuracy of machine adjustment, may cause actual application rates to deviate slightly. Therefore, Faesy & Besthoff, Inc. makes no warranty as to the uniformity of coverage obtained from the settings listed above.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

CAUTION: Keep Out of Reach of Children
Read Directions and Cautions before Using



E.P.A. Registration No. 779-79

FAESY & BESTHOFF, INC
EDGEWATER, N.J. 07020
25 LBS NET