



PRE-EMERGENCE CRABGRASS CONTROL

For Use: 2500 Sq.Ft. established lawn or 5,000 Sq.Ft. new lawn

CAUTION: Keep Out of Reach of Children and away from animals. Do not use on lawns that do not apply to areas used for food or feed purposes.

May cause irritation to the eyes, nose, throat and skin. If irritation occurs, wash skin or eyes with water and for eyes get medical attention. Wash hands thoroughly with soap and water after handling this material and before eating, drinking, or smoking. DO NOT INHALE DUST. Do not reuse empty containers. Do not burn or incinerate containers as waste or burning. Stay away from smoke or fumes.

This product is toxic to fish. Keep out of lakes, streams, ponds, etc. where runoff is likely to occur. Do not contaminate water by cleaning, maintenance or disposal of waste. Apply this product only as specified on this label.

PESTICIDE INGREDIENT STATEMENT

ACTIVE INGREDIENT:	
1-(2-Methylcyclohexyl)-3-phenylurea	3.8%
INERT INGREDIENTS	94.20%
TOTAL INGREDIENTS	100.00%

E.P.A. Reg. No. 779-85

*Also known as Turberson

© E.I. DuPont de Nemours, Inc.



Highly selective action kills crabgrass as it germinates without harming desirable lawn grasses.

APPROXIMATE SPREADER SETTINGS

Established Lawns (2500 Sq.Ft. Coverage)	Spring Seeding (5,000 Sq.Ft. Coverage)	Established Lawns (2500 Sq.Ft. Coverage)	Spring Seed- ing (5,000 Sq. Coverage)
--	--	--	---

Agri-Cred
Lawn Equity
Sports
Cyclone

Nov 7, 1973

Modern
Cross Country (Sears)
Jackson

GENERAL INFORMATION

Pre-Emergence Crabgrass Control is highly effective for controlling crabgrass, goosegrass, and grass (smooth and hairy), foxtail and several other newly seeded, established plantings of the following grasses: Bahia, Bermuda, Bunchgrass, Centipede, St. Augustine, Zoysia, and other lawn grasses. It is also effective for controlling crabgrass, goosegrass, and grass (smooth and hairy), foxtail and several other newly seeded, established plantings of the following grasses: Bahia, Bermuda, Bunchgrass, Centipede, St. Augustine, Zoysia, and other lawn grasses. It is also effective for controlling crabgrass, goosegrass, and grass (smooth and hairy), foxtail and several other newly seeded, established plantings of the following grasses: Bahia, Bermuda, Bunchgrass, Centipede, St. Augustine, Zoysia, and other lawn grasses.

PESTICIDE INGREDIENT STATEMENT

ACTIVE INGREDIENT:

Fluorenon 1-(2-Methylcyclohexyl)-3-phenylurea* 5.80%
OTHER INGREDIENTS 94.20%
TOTAL INGREDIENTS 100.00%



E.P.A. Reg. No. 779-85

*Also known as Turpersan
(R) E.I. DuPont de Nemours, Inc.

Highly selective lawn killer crabgrass as it germinates. It is safe for use on established lawns.

APPROXIMATE SPREADER SETTINGS

Established Lawns (2500 Sq. Ft. Coverage)	Spring Seeding (5,000 Sq. Ft. Coverage)	Established Lawns (2500 Sq. Ft. Coverage)	Spring Seeding (5,000 Sq. Ft. Coverage)

T.M.
 Agri-Sored
 Lawn Beauty
 Scotts
 Cyclone

Nov 7, 1973

Modern
 Cross Country (Scars)
 Jackson

GENERAL INFORMATION

Pre-Emergence Crabgrass Control is highly effective for controlling annual weed grasses such as crabgrass (smooth and hairy), foxtail and barnyard grass, in newly seeded or established plantings of the following grasses only: bluegrass, fescue, reedtop, smooth brome, perennial ryegrass, orchardgrass, zoysia and the following strains of bentgrass: Penncrest, Seaside, Highland, Astoria, Nimisila, C-1, C-7, and C-19. Do not use on other bentgrass strains or on bermuda grass as injury to turf might result. The effect on golf course greens may vary with turf management practices and environmental conditions; therefore, its use on golf greens is not recommended as injury to turf may result. Treatment will control annual bluegrass (Poa Annua), clover, or most broadleaf weeds.

DIRECTIONS FOR USE

1. Rake lawn to remove leaves and other debris.
2. ESTABLISHED LAWN-For established cool-season lawns, apply F&B Pre-Emergence Crabgrass Control at the rate of 4 lbs. per 1,000 Sq. Ft. (10 lb. bag covers 2500 Sq. Ft.). Apply after the 1st or 2nd mowing in Spring before the crabgrass and other annual grass weeds emerge. Highly effective for lawn seeded the previous fall as well as older grass areas. If it does not rain for 7 days after the application, water the treated area thoroughly for best results.
3. NEW SPRING SEEDINGS-For new Spring grass seedings, apply F&B Pre-Emergence Crabgrass Control at the rate of 2 lbs. per 1,000 Sq. Ft. (10 lb. bag covers 5,000 Sq. Ft.). Apply immediately after seeding which should be early in the spring before emergence of other annual grass weeds emerges. If it does not rain for 7 days after application, water the treated area. The early application is essential.

CAUTION

Do not apply to lawns where children or pets play. Do not apply to lawns where children or pets play. Do not apply to lawns where children or pets play. Do not apply to lawns where children or pets play. Do not apply to lawns where children or pets play.

