

F & B DIAZINON

25% Spray Concentrate

WARNING: KEEP OUT OF REACH OF CHILDREN

May be fatal if swallowed. May be absorbed through skin, and clothing. Wash thoroughly after handling. Avoid contamination of feed and foodstuffs. Do not use on household pets or humans. Do not permit children or pets to go onto the sprayed grass until the spray has completely dried. Keep out of reach of domestic animals. Do not store or use near heat or open flame. Do not reuse empty container. Destroy when empty.

This product is highly toxic to bees exposed to direct treatment, or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service.

This product is toxic to fish, birds and other wildlife. Birds and other wildlife in treated areas may be killed. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment or disposal of wastes. Apply this product only as specified on this label.

ACTIVE INGREDIENTS
 O, O-diethyl O-(2-isopropyl-4-methyl-6-pyrimidinyl) phosphorothioate 25%
 Aromatic Petroleum Derivative Solvent 55.78%
INERT INGREDIENTS 19.02%
Total 100.00%

NET CONTENTS: 1 QUART

Faesy & Besthoff, Inc. Edgewater, New Jersey 07020

PROBLEM AREA	PESTS	RATE & AREA COVERED	NOTES
Roses & Flowers: Roses, Azaleas, Chrysanthemums, Gladiolus, Cattails	Aphids, Mites, Scale, Leafhoppers, Leaf Miners, Fire Ants	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray	Apply when insects first appear. Repeat applications as needed or as noted in the rate chart.
Trees & Shrubs: Maple, Elm, Holly, Birch, Pine, Spruce, Fir	Aphids, Mites, Scale, Leafhoppers, Leaf Miners, Fire Ants	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a thorough foliar spray	Apply when insects first appear. Repeat applications as needed or as noted in the rate chart.
Fruits: Apples, Peaches, Oranges, Plums, Grapes, Cherries, Strawberries	Leafhoppers, Mites, Aphids, Scale, Fire Ants	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Apply when insects first appear. Repeat applications as needed or as noted in the rate chart.
Vegetables: Cucumbers, Squash, Tomatoes, Beans, Cauliflower, Cabbage, Broccoli, Carrots, Potatoes, Peas, Corn	Mites, Aphids, Leafhoppers, Scale, Fire Ants, Caterpillars, Cutworms, Beetles	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Apply when insects first appear. Repeat applications as needed or as noted in the rate chart.
Lawns: Turf, Sod, Sod Grass, Bermuda, St. Augustine, Centipede, Zoysia	Chinch Bugs, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs, Lawn Billbugs	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Apply when insects first appear. Repeat applications as needed or as noted in the rate chart.
Nuisance Pests in Outside Areas:	Fire Ants, Caterpillars, Boxelder Bug, Leafhoppers, Cicadas, Fleas, Chiggers, Fleas	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Apply when insects first appear. Repeat applications as needed or as noted in the rate chart.

E.P.A. Reg. No. 779-87-574 E.P.A. Est. No. 779-NJ-1
 LOT #674

HOW TO MIX AND APPLY

Step #1: Determine size of spraying job. For lawns and vegetables measure the area to be sprayed. For all other applications apply as a thorough cover spray. Make applications when insects first appear, or in advance of an expected infestation. Repeat applications as needed or as noted in the rate chart.

Step #2: Use clean sprayer. Carefully measure and mix the amount of Diazinon and water as suggested in the chart below.

Step #3: Spray as directed in rate chart. Use coarse drenching sprays for lawns, soil, and other surface applications. Use medium-fine mist sprays for trees and plants. Thorough coverage is important. Flush sprayer with clean water after each use.

See Chart for complete Insect List—use dosages shown unless otherwise indicated

2 tablespoons 1 fluid ounce or 6 teaspoons 1 cup 8 fluid ounces or 1 pint 2 tablespoons in 3 gals. water 2 teaspoons in 1 gal. water or 1/2 pint in 24 gals.

(Note: Do not use on certain ferns, poinsettia, hibiscus, papaya, pilea and gardenia, and orchids since injury may occur.)

CAUTION: Combustible Mixture

Rate Chart for Applying F & B Diazinon 25% Spray Concentrate

PROBLEM AREA	PESTS	RATE & AREA COVERED	NOTES
Lawns: Bermuda, St. Augustine, Centipede, Zoysia	Chinch Bugs*, Sod Webworms, Bermuda Mites, Ants, Ticks, Earwigs, Lawn Billbugs	2 tbsps. (1 fl. oz.) per 3 gals. of water. Apply as a full cover spray.	Thoroughly wet down grass a few hrs. before applying. For lawn Billbugs, water lawn thoroughly after treatment. *In the north, if Chinch Bugs are the only insect problem a lower rate of application can be used, 2 tbsps. in 3 gals. of water covers 250 sq. ft.

779-87

