

F & B Liquid Sevin*
PRODUCT INFORMATION

Liquid Sevin is an extremely fine dispersion of Sevin in water. **SHAKE WELL BEFORE USING. PROTECT FROM FREEZING.** Keep cover on opened container to prevent drying. Pour required amount of material into nearly filled spray tank or pre-mix material in another container half full of water and pour mixture into spray tank.

Add balance of water. Agitate while mixing and using. Spray all plant surfaces thoroughly. Use coarse spray to minimize drift.

USE PRECAUTIONS

CAUTION: HARMFUL IF SWALLOWED, INHALED OR ABSORBED THROUGH THE SKIN. Avoid breathing spray. Do Not Take Internally. Skin Contact May Be Harmful. Avoid Skin Contact. Wash after handling. Keep children and animals off treated lawns until these areas are dry. Food utensils such as teaspoons and measuring cups should not be used for food purposes after use with pesticides. Note For Physician: Carbaryl is a moderate, reversible, Cholinesterase inhibitor. Atropine is antidotal. **AVOID CONTAMINATION OF FOOD.**

To avoid possible fruit thinning on apples and foliage injury on McIntosh, delay use until at least 30 days after bloom. Dispose of container promptly when empty. This product is highly toxic to bees exposed to direct treatment or residues on crops. Protective information may be obtained from your Cooperative Agricultural Extension Service. Avoid spray drift to adjoining food or foliage crops or crops in bloom. Do not use with Lime or Alkaline materials.

F & B WARRANTY

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

No 2040

MARCH 74



LIQUID SEVIN*

Controls insects on ornamentals, flowers, lawns, fruit trees and vegetables. Kills Gypsy Moths, Japanese Beetles, Army Worms, Chinch Bugs, Lawn Moths, Ants, Mosquitoes and certain other insects.

ACTIVE INGREDIENTS

Carbaryl: (1-Naphthyl N-Methylcarbamate)22.5%

INERT INGREDIENTS77.5%

Contains 2 lbs. Sevin per gallon


CAUTION:

Keep out of Reach of Children. See Side Panel for Additional Cautions.

*Trade Mark of Union Carbide for the Active Ingredient. U.S. Pat. No. 2,903,478

1 FLUID QUART NET

FAESY & BESTHOFF, INC., Edgewater, N.J. 07020



LIQUID SEVIN*

Controls insects on ornamentals, flowers, lawns, fruit trees and vegetables. Kills Gypsy Moths, Japanese Beetles, Army Worms, Chinch Bugs, Lawn Moths, Ants, Mosquitoes and certain other insects.

ACTIVE INGREDIENTS

Carbaryl: (1-Naphthyl N-Methylcarbamate) ...22.5%

INERT INGREDIENTS77.5%

Contains 2 lbs. Sevin per gallon

CAUTION:

Keep out of Reach of Children. See Side Panel for Additional Cautions.

*Trade Mark of Union Carbide for the Active Ingredient. U.S. Pat. No. 2,903,478

1 FLUID QUART NET

FAESY & BESTHOFF, INC., Edgewater, N.J. 07020

F & B Liquid Sevin may be used in all type sprayers. Shake this container well before using. Mix 2 pints in 50 gallons of water (4 teaspoons per gallon or 2 ounces in 3 gallons).

RECOMMENDATIONS:

ORNAMENTAL PLANTS, FLOWERS, SHRUBS AND SHADE TREES including rose, carnation, gladiolus, zinnia, chrysanthemum, lilac arbor vitae, juniper, hydrangea, azalea, elm, maple, oak, dogwood, birch and pines (Do not use on Boston Ivy). Apply when insects or damage appear. Repeat weekly or as needed to control: apple aphid, bagworm, birch leaf miner, blister beetles, box-elder bug, boxwood leaf miner, elm leaf beetle, flea beetles, gypsy moth, Japanese beetle, June beetles, lace bugs, leafhoppers, leaf rollers, mealy bugs, mimosa webworm, oak leaf miners, orange tortrix, periodical cicada, plant bugs, puss caterpillars, psyllids, rose aphid, roseslug, scale insects, tent caterpillars, thorn bugs, thripes, (exposed), willow leaf beetles.

LAWN: Ants, chinch bugs, earwigs, fall armyworms, fleas, leafhoppers, mosquitoes, sod webworms (lawn moths), millipedes. Apply finished spray solution at rate of 1 gallon per 40 sq. ft of established lawn. For best results, mow lawn and apply immediately after rain or watering. Repeat in 2 or 3 weeks if necessary.

VEGETABLE GARDEN SPRAY: Apply when insects or damage appear and repeat at 7 to 10 day intervals if necessary. When used at the above rates F & B Liquid Sevin may be applied to the following vegetables right up to the day of Harvest.

LIMA BEANS AND SNAP BEANS—for control of Japanese and Mexican Bean Beetles, and Tarnished Plant Bugs. **TOMATO, EGGPLANT AND PEPPER**—for Leafhopper, Lace Bugs and Tomato fruit worms. **MELON, SQUASH, CUCUMBER AND PUMPKIN**—Flea Beetles, Melonworms, Cucumber beetles and Squash Bugs. Avoid excessive applications. Some leaf injury may occur on tender foliage when prolonged rain or humidity follows spraying. Do not use on Watermelons in Florida.

FRUIT TREES (APPLES, PEARS AND PEACHES): Apple Maggot, Plum Curculio, Apple Aphid, Leaf Rollers, Peach Tree Borer, Oriental Fruit Moth, Codling Moth—Make one application between bud break and bloom, a second application at petal fall, and then at 3 to 4 week intervals for 3 to 4 months as necessary. Apple Maggot—For best control, apply at 10 day intervals. Rosy Apple Aphid—Apply before leaves are curled. Do not apply within 1 day of harvest.

E.P.A. Reg. #779-92 EFA Est. 779-NJ-1