



LAWN FUNGICIDE

Contains[®] Dyrene

GRANULAR

GUARANTEED ANALYSIS

ACTIVE INGREDIENT:	
4,6-Dichloro-N-(2-chlorophenyl)-1,3,5-triazin-2-amine	5%
INERT INGREDIENTS:	
Total	100%

DYRENE is a registered trademark of Mobay Chemical Corporation

MANUFACTURED BY:

FAESY & BESTHOFF, INC.
EDGEWATER, NEW JERSEY 07020

E. P. A. EST. NO. 779-NJ-1
E. P. A. REGISTRATION NO. 779-97

A BROAD SPECTRUM FUNGICIDE FOR CONTROL OF IMPORTANT FUNGUS DISEASES OF TURFGRASSES.

CONTROLS THE FOLLOWING DISEASES:

BROWN PATCH, COPPER SPOT, DOLLAR SPOT, HELMINTHOSPORIUM LEAF SPOTS, MELTING-OUT RUSTS AND TYPHULA SNOW MOLD

SUGGESTED TRIAL SPREADER SETTING

	2.4 lbs. per thousand sq. ft.	5.4 lbs. per thousand sq. ft.
Scotts	4½	6¼
Cyclone.....	2	4

NOTICE: Normal differences in equipment and materials, as well as condition of machine, adjustment, and walking speed may cause variation in application rate. Therefore, Faesy & Besthoff, Inc. makes no warranty as to the uniformity of coverage from the above spreader settings.

Seller warrants that the product conforms to its chemical description and is reasonably fit for the purpose stated on the label when used in accordance with directions under normal conditions of use, but neither this warranty nor any other warranty of merchantability or fitness for a particular purpose, express or implied, extends to the use of this product contrary to label instructions, or under abnormal conditions, or under conditions not reasonably foreseeable to seller, and buyer assumes the risk of any such use.

DIRECTIONS FOR USE

For best results, the lawn should be moist when application. Just after a light shower in the morning when dew is on the grass is ideal. LAWN FUNGICIDE should not be applied when rain is expected within 24 hours. Do not water lawn after application.

Apply from 2.5 to 5.4 lbs. of F&B Granular Lawn Fungicide per thousand square feet for control of turf diseases indicated on this label. Apply at 7 to 14 day intervals through spring and summer.

For Typhula Snow Mold: Apply 2½ lbs. per thousand square feet, beginning after first killing frost in fall. Repeat when grass is free of snow in mid-winter and again after final snow melts in spring.

Lawn spreaders suitable for application of Granular Turf products should be used in making this application.

Calibrate your spreader as follows:

1. Accurately measure an area on your lawn 10 ft. long by 10 ft. wide.
2. Using the Suggested Trial Spreader Settings listed herein, accurately weigh ¼ lb. F&B Granular Lawn Fungicide into spreader compartment.
3. Apply at normal working speed to the measured area.
4. Adjust spreader accordingly to deliver more or less material as required.

CAUTION

KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed.

Avoid inhalation.

Avoid contact with skin, eyes and clothing. In case of skin contact, wash with soap and water; for eyes, get medical attention.

Avoid Contamination of feed and foodstuffs.

Do not graze treated area. Do not feed clippings to livestock.

Do not reuse empty container. Destroy by burying with wastes or burning. Stay away from smoke or fumes.

FISH AND WILDLIFE CAUTION

This product is toxic to fish.

Keep out of lakes, streams or ponds.

Do not contaminate water by cleaning of equipment, or disposal of wastes.

Apply this product only as specified on this label.

ACCEPTED
NET WEIGHT (10 POUNDS)

JUN 23 1975

Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under E.P.A. Reg. No. 779-97

6/75