

Camerasman's Note
Poor Copy

Diazinon[®] 14G

Granular Insecticide

Active Ingredient:
O,O-diethyl O-(2-iso-
propyl-6-methyl-4-
pyrimidinyl) phos-
phorothioate 14.0%
Inert
Ingredients: 86.0%
Total: 100.0%

For control of soil
insects including
wireworms, cut-
worms, and other
crop pests (as
listed in label
directions)

Caution: KEEP OUT OF REACH OF CHILDREN

Harmful if swallowed. Avoid contamination of
food and feed. Do not store near food or feed
products. Avoid inhalation of dust. Avoid contact
with skin, eyes, and clothing. Wash thoroughly
after handling. May be absorbed through skin.

Fish and Wildlife Caution

This product is toxic to fish and wildlife. Keep out of
streams, ponds, lakes, and other bodies of water.
Do not apply directly to water.

ACCEPTED
JUN 6 1975
Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 779-99

DIRECTIONS FOR USE AND CONDITIONS OF SALE AND WARRANTY

IMPORTANT: Read the entire Directions for Use
and the Conditions of Sale and Warranty before
using this product.

Conditions of Sale and Warranty

Follow directions on label for use and application.
Do not use on crops or plants not listed on label.
Do not use on crops or plants if they are under
treatment with another insecticide.
Do not use on crops or plants if they are under
treatment with a fungicide.
Do not use on crops or plants if they are under
treatment with a rodenticide.
Do not use on crops or plants if they are under
treatment with a nematocide.

Directions for Use

Apply to the soil in the row or bed
the top few inches of soil
with a rotary tillage implement
or a similar implement.

Field and Forage Insects

Peas (Reno).
Apply to the soil in the row or bed
the top few inches of soil
with a rotary tillage implement
or a similar implement.

Corn
Corn Rootworm, European Cornborer, and
Southwestern Cornborer.

Southwestern Corn Borers. Apply 14 lbs per
acre over the corn so granules fall into whorls
when second-generation borers first appear.
Repeat application one week later.

Note: Do not feed treated feed to dairy or beef
cattle or sheep for 10 days following applica-
tion. Do not feed treated feed to dairy or beef
cattle or sheep for 10 days following applica-
tion of the granules to the soil.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Cutworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Peas (Reno). Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Peas (Reno). Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Corn
Corn Rootworm, European Cornborer, and
Southwestern Cornborer.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Sugarcane

Sugarcane Borers. Broadcast 14 lbs per acre
after plants have begun to form and within
east 500 ft of plants still with borers feeding
under the leaf sheath. Do not broadcast
into the soil. Additional applications may be
made if necessary.

Note: Do not use more than 14 lbs per acre
per growing season. Allow 28 days between
applications. Do not apply to soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Tobacco

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Wireworms. Broadcast 14 lbs per acre before
planting. Apply to the soil in the row or bed
the top few inches of soil with a rotary tillage
implement or a similar implement.

Directions for Use

Applications to the soil should be worked into the top few inches immediately after application with a rotary hoe, cultivator, disk, or other suitable means for greatest effectiveness.

Field and Forage Insects

Beans (Pinto)

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high.

Corn

Corn Rootworm. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time.

Apply 4 1/2 lbs. per acre in a 5-inch band over a handover row at planting time or when the plants are 10 to 15 inches high, but not later than July 14. Make a second application July 14 to 15. Make a third application August 14 to 15. Apply 1.4 lbs. per acre in a 5-inch band over the treated band when the plants are 10 to 15 inches high.

Apply amount per acre recommended according to row spacing and other spacing treatment.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Apply amount per acre recommended according to row spacing and other spacing treatment.

Suggested Rates According to Corn Row Spacing

Row spacing	Linear feet of corn row per acre	Amount of Dieldrin 14G per acre	Postemergence basal treatment
48	1,280	1.4 lbs.	1/2 inch of soil
42	1,429	1.4 lbs.	1/2 inch of soil
40	1,571	1.4 lbs.	1/2 inch of soil
39	1,633	1.4 lbs.	1/2 inch of soil
36	1,778	1.4 lbs.	1/2 inch of soil
30	2,240	1.4 lbs.	1/2 inch of soil
24	2,800	1.4 lbs.	1/2 inch of soil
20	3,360	1.4 lbs.	1/2 inch of soil
15	4,480	1.4 lbs.	1/2 inch of soil

Note: For any spacing, the amount of insecticide does not exceed 1.4 lbs. per acre.

European Corn Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Flazop 14.5. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

6-1 Armyworms. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Some 8 inches. This preplant treatment will also control other soil insects such as cutworms, surface and subterranean and wireworms.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Peanuts

Summer Corn Rootworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Sorghum

Cutworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Soybeans

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Sugar Beets

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Mole Crickets. Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 12 days prior to sowing of seed or transplanting and work into soil 1 1/2 inches.

Seed Corn Maggots. Broadcast 1 1/2 lbs. per acre 4 weeks before transplanting. Work into soil 2 inches.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre 4 weeks before transplanting. Work into soil 4 to 6 inches.

Forage Crops (Alfalfa, Clover, Cowpeas, and Lespedeza)

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Vegetable Crop Insects

Broccoli, Brussels Sprouts, Cabbage, and Cauliflower

Root Maggots. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Onions

Onion Maggots. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Radishes

Root Maggots. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Sweet Corn

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Tomatoes

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Other Vegetable Insects

Cutworms (Surface and Subterranean)

Species	Rate (lbs. per acre)	Application
Black Cutworm	1.4	5-inch band
Common Nighthawk	1.4	5-inch band
Flourish Cutworm	1.4	5-inch band
Green Cutworm	1.4	5-inch band
Large Cutworm	1.4	5-inch band
Small Cutworm	1.4	5-inch band
White Cutworm	1.4	5-inch band

Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borers

Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms

Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.

Small Fruit Insects

Strawberries

Maggots. Apply 7 lbs. per acre when insects are present. Make application to plant beds or fields 12 days prior to transplanting and work into soil 1 1/2 inches.

Lawn Insects

Lawn Insect Bugs. Apply 1.4 lbs. per 1,000 sq. ft. or 14 lbs. per 10,000 sq. ft. and crush down to work into soil down to the soil surface. Water down thoroughly after application. Repeat application 10 to 15 days thereafter. For longer periods of control, and on dense growing lawns, use 2.8 lbs. per 1,000 sq. ft. or 28 lbs. per 10,000 sq. ft.

Firebugs, White Grubs such as Japanese Beetle Larva, European Chaffer and Southern Chaffer. Apply 1.4 lbs. per 1,000 sq. ft. or 14 lbs. per 10,000 sq. ft. after application. For longer periods of control, and on dense growing lawns, use 2.8 lbs. per 1,000 sq. ft. or 28 lbs. per 10,000 sq. ft.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Hyperbaena. Apply 1.4 lbs. per 1,000 sq. ft. or 14 lbs. per 10,000 sq. ft. after application. For longer periods of control, and on dense growing lawns, use 2.8 lbs. per 1,000 sq. ft. or 28 lbs. per 10,000 sq. ft.

Wireworms. Broadcast 2 1/2 to 2 3/4 lbs. per acre before planting. Work into soil 4 to 6 inches.

Lesser Cornstalk Borer. Apply 1.4 lbs. per acre in a 5-inch band over the row at planting time or when the plants are 10 to 15 inches high. Make additional applications at planting time or when the plants are 10 to 15 inches high. Cover the treated band with 1/2 inch of soil.