

#### HOSPITAL DISINFECTION

Use for disinfecting and deodorizing bed frames, mattress covers, rubber sheets, urinals, bed pans, catheter bottles, diaper pails and wheel chairs. Excellent for general cleaning and disinfecting of sick rooms, operating rooms and bathrooms. To disinfect walls, floors, furniture and other hard surfaces in hospitals and nursing homes, mop or brush surface to be treated with a solution containing 2 ounces of TRIPLE D per gallon of water. Those surfaces that are heavily soiled with blood, fecal matter or exudates should be thoroughly scrubbed with a solution containing 2 ounces per gallon of water before disinfection. This solution will kill *Pseudomonas aeruginosa*. Excellent for odor control in bathroom and incontinent areas. In the event of urine spills, these areas when mopped are deodorized and disinfected.

#### FOR CLEANING

Lightly soiled areas—surfaces to be cleaned should be mopped or brushed with a solution containing 1 ounce of TRIPLE D per gallon of water. Rinse cleaned area with fresh water. Heavily soiled area—mop or brush surface with a solution containing 2-3 ounces of TRIPLE D per gallon of water. Rinse cleaned surface with fresh water.

#### DEODORIZATION

To deodorize surface during cleaning, mop or brush surface with a solution containing 2 ounces of TRIPLE D per gallon of water. TRIPLE D is an effective antibacterial cleaner designed for use in household, industrial and institutional cleaning. When used as directed, TRIPLE D will disinfect hard surfaces, such as floors, walls, counter tops and furniture. It may be relied upon to deodorize bathroom areas, garbage pails, food storage areas, kennel areas and other areas where obnoxious odors may develop. This product may be used as a general purpose antibacterial detergent in schools, restaurants, hospitals, nursing homes, public rooms, food processing plants and in other places where efficient cleaning and antimicrobial action is desired.

# Triple D

Disinfectant - Cleaner - Sanitizer  
Fungicide - Deodorant  
FOR HOSPITAL USE

ACTIVE INGREDIENTS  
n-Alkyl (50% C<sub>12</sub>, 40% C<sub>14</sub>, 10% C<sub>16</sub>) dimethylbenzyl ammonium chloride 5.0%  
Tetrasodium salt of ethylene diamine tetraacetic acid 2.3%  
Sodium carbonate 2.0%  
INERT INGREDIENTS 90.7%  
PHENOL COEFFICIENT  
S. TYPHOSA 45 S. AURLUS 24

#### DANGER

EPA Reg. No. 784-87

#### KEEP OUT OF REACH OF CHILDREN


See First Aid statement and other precautions on right panel.  
Container Disposal: When empty, rinse container thoroughly with water and destroy it by crushing or burying in a safe place.

NET CONTENTS: ONE GALLON



Consolidated Chemical, Inc.  
St. Louis, Missouri 63110

EPA EST. No. 00784-Mo.-1



#### GENERAL PURPOSE DISINFECTION

To disinfect walls, floors and other hard surfaces in schools, institutions, and public buildings; mop or brush surface to be treated with a solution containing 1 ounce TRIPLE D per gallon of water. Surfaces that are heavily soiled should be cleaned with 2 ounces of TRIPLE D per gallon of water and then disinfected. Those surfaces which may later come in contact with food must be rinsed with clear water of drinking quality. To disinfect locker rooms, shower stalls, walkways around pools and other areas where fungicidal action is required, mop area with a solution containing 2 ounces in 3 gallons of water. This solution will destroy *Trichophyton interdigitale*, a causative agent of Athlete's foot.

#### DANGER

#### KEEP OUT OF REACH OF CHILDREN

**CORROSIVE.** Causes eye damage and skin irritation. Do not get in eyes, on skin or on clothing. Protect eyes and skin when handling. Harmful if swallowed. Avoid contamination of food. **FIRST AID:** In case of contact, immediately flush eyes or skin with plenty of water for at least 15 minutes. For eyes, call a physician. Remove and wash contaminated clothing before reuse. If swallowed, drink promptly a large quantity of milk, egg whites, gelatin solution or if these are not available, drink large quantities of water. Call a physician immediately. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage. Measures against circulatory shock, respiratory depression and convulsion may be needed.