

Front Panel ...

Miller's

LIQUID LIME SULPHUR

San Jose Scale - Scab - Powdery Mildew

on Apples - Pears

Clean up Spray - Roses

ACTIVE INGREDIENT

Calcium Polysulfide. 29%

INERT INGREDIENTS 71%

Density - Baume, at 60°F31°

Contains calcium and sulfur expressed as gypsum -

3.0 lbs. per gallon. Other combined sulfur 1.9 lbs.

per gal. Product Weight 10.59 lbs. per gal. at 68°F.

E.P.A. Reg. No. 802-1-ZA

CAUTION - Keep out of reach of children

See right side panel for additional cautions.

NET CONTENTS _____ GALLONS

Manufactured for:

THE CHAS. H. LILLY CO.

PORTLAND, OREGON

EPA Est. 802-OR-1

802-1
26

DILUTION TO USE

Number	Amount to Use
1	11 gals. Miller's Liquid Lime Sulphur to 100 gallons water; or 2-2/3 pint to 3 gallons water.
2	3 gals. Miller's Liquid Lime Sulphur plus 1-1/2 to 2 gals. Miller's Superior Type Spray Oil to 100 gallons of water; or 1-1/2 cups Miller's Liquid Lime Sulphur plus 3/4 to 1 cup Miller's Superior Type Spray Oil to 3 gallons of water.
3	2-1/2 to 3 gals. Miller's Liquid Lime Sulphur to 100 gallons of water; or 3/4 pint to 3 gallons of water.
4	2-1/2 gals. Miller's Liquid Lime Sulphur to 100 gallons of water; or 1-1/4 cup to 3 gallons of water.
5	2 to 2-1/2 gals. Miller's Liquid Lime Sulphur to 100 gallons of water; or 1-1/4 cups to 3 gallons of water.
6	12-1/2 gals. Miller's Liquid Lime Sulphur to 100 gallons of water; or 3 pints to 3 gallons of water.
7	8-1/3 gals. Miller's Liquid Lime Sulphur to 100 gallons of water; or 1 qt. to 3 gallons of water.

CAUTION

May cause skin irritation. Avoid contact with skin or eyes. In case of contact with skin or eyes, flush immediately with plenty of water for at least 15 minutes.

Do not spray straight or diluted Miller's Liquid Lime Sulphur against painted surfaces. Walls can be protected with heavy paper or a tarpaulin. Do not use on Apples or Pears immediately before or after an oil spray. Do not use on sulphur-susceptible varieties of Pears and Apples after the Pink Sprays. Use with caution on Apples in the Pink Sprays. Consult State Agricultural Experiment Station or State Extension Service Authorities for exact timing and spacing of sprays in your area.

Rinse equipment and containers and dispose of wastes by burying in non-crop lands away from water supplies. Containers should be disposed of by breaking or punching holes in them and burying with wastes.

SPRAY CALENDAR

Crop and Pest	Time of Application	Dilution to Use
APPLES San Jose Scale	Dormant: As winter buds swell, just before opening.	1
San Jose Scale Oyster Shell Scale Blister Mite	Dormant: As winter buds swell, just before opening.	2
Scab and Powdery Mildew	Prepink, or green bud spray: Leaves separating just exposing blossom bud cluster.	3
Scab and Powdery Mildew	Pink, or preblossom spray: Just before blossoms open.	4
Scab and Powdery Mildew	Calyx, when $\frac{3}{4}$ petals have fallen.	5
PEARS San Jose Scale	Dormant: As winter buds swell, just before opening.	1
Scab and Powdery Mildew	Prepink: Blossom bud clusters just exposed.	3
Scab and Powdery Mildew	Pink, or preblossom spray: Just before blossoms open.	4
ROSES Clean-up Spray	Dormant: As winter buds swell—before opening. December or January.	6
	If roses not sprayed before buds swell—February.	7

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals and specifically recommended and other influencing factors in the use of this product are to be the control of the user. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

MP-9J