

ACCEPTED



MICROCOP

(Micronized Tribasic Copper Sulphate)
WETTABLE POWDER

FIXED COPPER SPRAY
For Use in place of
BORDEAUX

for Controlling Certain Diseases

on
FRUITS NUTS SMALL FRUITS
VEGETABLES FLOWERS SHRUBS

ACTIVE INGREDIENT
*Copper expressed as metallic 50.0%
INERT INGREDIENTS 50.0%
*Equivalent to 95% Micronized Tribasic Copper Sulphate, Commercial

E.P.A. Reg. No. 802-12-AA

CAUTION - Keep out of reach of children
See back panel for additional cautions

NET WEIGHT 4 LBS.

MP-12K

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218
EPA Est. 802-OR-1



Miller's MICROCOP

PLEASE READ ENTIRE LABEL. USE ONLY AS DIRECTED.
NOTING ALL CAUTIONS AND WARNINGS.

Miller's MICROCOP is a "Fixed" Copper Spray used in place of Bordeaux Mixture. It is very stable and does not deteriorate rapidly like Bordeaux Mixture when exposed to moist air.

MIXING DIRECTIONS

1. For Large Spray Tanks — fill tank about 1/4 full with water and, with agitator running, add the required amount of Miller's MICROCOP. When the tank is 1/2 full add spreader, if desired, and fill the tank. Keep agitator running during filling and spraying operations.
2. For small amounts — mix the required amount of Miller's MICROCOP with a small amount of water to form a smooth creamy paste. Next add this paste to the required amount of water. Shake sprayer frequently while spraying to insure uniformity of solution.

SPECIFIC DIRECTIONS

- PEACHES — LEAF CURL**
Use 6 lbs. to 100 gals. of water (4 tablespoons to 1 gal.) Apply in late fall or early winter before winter rains or snow. Repeat in late winter or early spring before buds break.
- PEACHES, APRICOTS — Coryneum Blight**
Use 4 lbs. to 100 gals. of water (2 1/2 tablespoons to 1 gal.). Apply after harvest and again before fall rains.
- APPLES, PEARS — Bulls-eye Rot**
Use 4 lbs. Miller's MICROCOP plus 1 qt. Miller's SPRA-OIL to 100 gals. of water (2 1/2 tablespoons Miller's MICROCOP plus 2 teaspoons Miller's SPRA-OIL to 1 gal.). Apply as a wetting spray after fruit harvest.
- SMALL FRUITS**
Use 6 lbs. to 100 gals. of water (4 tablespoons to 1 gal.) for the following:
Blackberry Leaf and Cone Spot. Apply sprays about September 15th, March 15th and April 1st.
Gooseberry and Currants — Anthracnose and Leafspot. Apply sprays at full bloom stage repeat in two weeks and re-apply after harvest.
- WALNUTS — Blight**
Use 3 lbs. to 100 gals. of water (2 tablespoons to 1 gal.). Apply in early pre-bloom stage Repeat in late pre-bloom and early post-bloom stages.
- ALMONDS (in California) — Brown Rot**
Use 4 to 5 lbs. per 100 gals. of water. Apply to thoroughly cover entire tree at Pink Bud Stage. Repeat at Popcorn Stage.
- CELERY, POTATOES, TOMATOES — Late Blight**
Use 4 lbs. to 100 gals. of water (2 1/2 tablespoons to 1 gal.). Apply at first sign of blight. Repeat at 4 to 5 day intervals until under control.
- AZALEA — Leaf Gall**
Use 4 lbs. to 100 gals. water (2 1/2 tablespoons to 1 gal.). Apply to entire plant before buds break in the spring. Repeat 2 to 3 weeks later. Should any galled leaves appear, remove and burn.
- LILAC — Bacterial Blight**
Use 4 lbs. to 100 gals. of water (2 1/2 tablespoons to 1 gal.). Apply thoroughly in September and again before fall rains. Prune out and burn affected parts of the bush as they appear.
- FLOWERING SHRUBS — certain Leaf Spots**
Use 4 lbs. to 100 gals. of water (2 1/2 tablespoons to 1 gal.) Spray at first appearance of small discolored areas on leaves. Repeat sprays as necessary.
- LILIES, PEONIES — Botrytis Blight**
Use 4 lbs. to 100 gals. of water (2 1/2 tablespoons to 1 gal.) Apply as a thorough coverage spray when young plants emerge. Repeat every 7 to 10 days until blooming. Remove and burn all debris and affected plant parts from growing beds.
- ARBOR VITAE — Berckman's Blight**
Use 3 lbs. to 100 gals. of water (2 tablespoons to 1 gal.). Spray twice in fall. First spray late September or early October and second spray in early November.

CAUTION

Harmful if swallowed! Avoid inhaling dust or spray mist. Keep out of reach of children.
This product is toxic to fish. Keep out of lakes, streams or ponds. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.
Consult your State Agricultural Experiment Station or State Extension Service Authorities for additional information as the timing, number and rate of applications will vary with local conditions.

NOTICE: Follow directions carefully. Time and method of application, weather and crop conditions are with other chemicals not specifically recommended and their use is at the user's risk. The information on this label are based on the date of the label being printed. It is the user's responsibility to check for any changes in the label information with the date of purchase.

MP-12K