

Front Panel

Lilly Miller  
MICROCCOP  
Fungicide  
(Micronized Tribasic Copper Sulphate)

ACCEPTED  
SEP 27 1982  
Federal Insecticide,  
Fungicide and Rodenticide Act,  
for the pesticide  
E.C. No. 803-72

FIXED COPPER SPRAY FOR  
USE IN PLACE OF BORDEAUX

Fruits Nuts Small Fruits  
Vegetables Flowers  
Shrubs

Keep out of reach of children — 12 pt.  
CAUTION — 18 pt.  
See side panel for additional precautionary statements

ACTIVE INGREDIENT  
\*Copper, expressed as  
metallic . . . . . 50.0%  
INERT INGREDIENTS . . . . . 50.0%

\*Equivalent to 95% Micronized Tribasic Copper  
Sulphate, Commercial

NET WT. 1 LB.

*Left Side Panel*

PRECAUTIONARY STATEMENTS

*Hazards to Humans and Domestic Animals*

CAUTION

*Keep out of reach of children. Avoid inhaling dust or spray mist. Wash thoroughly after use.*

ENVIRONMENTAL HAZARDS

*This product is toxic to fish. Do not apply directly to water. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from areas treated. Do not contaminate water by cleaning of equipment, or disposal of wastes. Apply this product only as specified on this label.*

DIRECTIONS FOR USE

*It is a violation of Federal law to use this product in a manner inconsistent with its labeling.*

STORAGE AND DISPOSAL 12 pt.

*Do not reuse container. Wrap container and place in trash collection.*

**SPECIFIC DIRECTIONS**  
Not for use in greenhouses

**AZALEA** Leaf Gall

1. Use 1 lb. to 25 gals. of water (2 2/3 tablespoons to 1 gal.).
2. Apply to entire plant before buds break in the spring. Repeat 2 or 3 weeks later.
3. Should any galled leaves appear, remove and burn.

**LILAC** Bacterial Blight

1. Use 1 lb. to 25 gals. of water (2 2/3 tablespoons to 1 gal.).
2. Apply thoroughly in September and again before fall rains.
3. Prune out and burn affected parts of the bush as they appear.

**FLOWERING SHRUBS** certain Leaf Spots (*Cercospora*, *Septoria*, *Botrytis*)

1. Use 1 lb. to 25 gals. of water (2 2/3 tablespoons to 1 gal.).
2. Spray at first appearance of small discolored areas on leaves.
3. Repeat sprays as necessary.

**ARBOR VITAE** Berckman's Blight

1. Use 1 lb. to 33 gals. of water (2 tablespoons to 1 gal.).
2. Spray twice in fall. First spray in late September or early October and second spray in early November.

**LILIES, PEONIES** Botrytis Blight

1. Use 1 lb. to 25 gals. of water (2 2/3 tablespoons to 1 gal.).
2. Apply as a thorough coverage spray when young plants emerge. Repeat every 7 to 10 days until blooming.
3. Remove and burn all debris and affected plant parts from growing beds.

**NOTICE:** Seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Buyer assumes all risk of use and of handling of this material when such use and/or handling is contrary to label instructions.

15-R

THE CHAS. H. LILLY CO.  
Portland, Oregon 97218

EPA EST. NO. 3000-K-1

EPA Reg. No. 802.12

Right Side Panel

MICROCOP FUNGICIDE is a "Fixed" Copper Spray used in place of Bordeaux Mixture. It is very stable and does not deteriorate rapidly like Bordeaux Mixture when exposed to moist air.

MIXING DIRECTIONS

1. For Large Spray Tanks - Fill tank about 1/4 full when water and with agitator running, add the required amount of MICROCOP FUNGICIDE. When the tank is 2/3 full add spreader, if desired, and fill the tank. Keep agitator running during the filling and spraying operations.
2. For small amounts - Mix the required amount of MICROCOP FUNGICIDE with a small amount of water to form a smooth creamy paste. Next add this paste to the required amount of water. Shake sprayer frequently while spraying to insure uniformity of solution.

Lilly/Miller Sta-Stuk "m" extends the effective life of Lilly/Miller by resisting the washing effects of rains and snows.

Consult your State Agricultural Experiment Station or State Extension Service Authorities for additional information as the timing, number and rate of application will vary with local conditions.

SPECIFIC DIRECTIONS

Not for use in greenhouses

PEACHES - Leaf Curl

1. Use 1 lb. to 16 gals of water (4 tablespoons to 1 gal.).
2. Apply in late fall or early winter before winter rains or snow.
3. Repeat in late winter or early spring before buds break.

PEACHES, APRICOTS - Coryneum Blight

1. Use 1 lb. to 25 gals. of water (2 2/3 tablespoons to 1 gal.).
2. Apply after harvest before fall rains.

APPLES, PEARS - Anthracnose, Perennial Canker

1. Use 1 lb. MICROCOP FUNGICIDE plus 1 cup superior type oil to 25 gals of water (2 2/3 tablespoons MICROCOP FUNGICIDE plus 2 teaspoons superior type oil to 1 gal.)
2. Apply as a wetting spray after fruit harvest.

SMALL FRUITS

1. Use 1 lb. to 16 gals. of water (4 tablespoons to 1 gal. for the following).
2. Blackberry Leaf and Cane Spot. Apply sprays after September 15th, March 15th, and April 1st.
3. Gooseberry and Currants - Anthracnose and Leafspot. Apply sprays at full bloom stage. Repeat in two weeks and reapply after harvest.

Right Side Panel Continued

**WALNUT - Blight**

1. Use 1 lb. to 33 gals. of water (2 tablespoons to 1 gal.)
2. Apply in early pre-bloom stage. Repeat in late pre-bloom and early post-bloom stages.

**CELERY, POTATOES, TOMATOES - Late Blight**

1. Use 1 lb. to 25 gals. of water (2 2/3 tablespoons to 1 gal.).
2. Apply at first sign of blight. Repeat at 4 to 5 day intervals until under control.

LAB L COPY as appears on litho'd containers . . . . .

LILLY/MILLER  
**Sta-stuk**  
"m"



**PROTECT FROM FREEZING**

**DIRECTIONS FOR USE**

1. For use only with Lilly/Miller MICROCOP FUNGICIDE or other fixed copper sprays
2. Use 4 teaspoons per gallon of spray (4 tablespoons per 3 gallons of spray)
3. Add the required amount to the sprayer or spray tank **last**. Stir or shake well to insure thorough mixing
4. On producing fruit trees or caneberryes use only with post-harvest and dormant copper sprays

Lilly/Miller STA-STUK "m" extends the effective life of Lilly/Miller MICROCOP FUNGICIDE sprays by resisting the washing effects of rains and snows

Lilly/Miller STA-STUK "m" should be used on producing fruit trees and caneberryes only with post-harvest or dormant copper sprays.

**PLEASE READ ENTIRE LABEL.  
USE ONLY AS DIRECTED  
NOTING ALL CAUTIONS.**

**CAUTION:** May cause skin irritation. Avoid contact with skin, eyes or clothing. In case of contact, flush with plenty of water, for eyes, get medical attention. Destroy empty container. Do not re-use for any purpose.

**IMPORTANT** - After spraying be sure to wash out sprayer with water to prevent Lilly/Miller STA-STUK "m" from drying and plugging orifices. Do not allow spray mixture to stand in sprayer overnight.

**NOTICE:** Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

**LILLY MILLER STA-STUK "m" sticks Lilly/Miller MICROCOP FUNGICIDE sprays on trees and shrubs through all kinds of weather.**

**THE CHAS. H. LILLY CO.**  
PORTLAND, OREGON 97218

Copyright © 1962