

U.S. ENVIRONMENTAL PROTECTION AGENCY
OFFICE OF PESTICIDES PROGRAMS
REGISTRATION DIVISION (WH-567)
WASHINGTON, D.C. 20460

EPA REGISTRATION NO.

102-16

DATE OF ISSUANCE
MAY 14 1985

TERM OF ISSUANCE

REREGISTERED

NAME OF PESTICIDE PRODUCT

NOTICE OF PESTICIDE: REGISTRATION
 REREGISTRATION

(Under the Federal Insecticide, Fungicide,
and Rodenticide Act, as amended)

NAME AND ADDRESS OF REGISTRANT (Include ZIP code)

┌
└

656
19
149018
1

NOTE: Changes in labeling formula differing in substance from that accepted in connection with this registration must be submitted to and accepted by the Registration Division prior to use of the label in commerce. In any correspondence on this product always refer to the above U.S. EPA registration number.

On the basis of information furnished by the registrant, the above named pesticide is hereby Registered/Reregistered under the Federal Insecticide, Fungicide, and Rodenticide Act.

A copy of the labeling accepted in connection with this Registration/Reregistration is returned herewith.

Registration is in no way to be construed as an indorsement or approval of this product by this Agency. In order to protect health and the environment, the Administrator, on his motion, may at any time suspend or cancel the registration of a pesticide in accordance with the Act. The acceptance of any name in connection with the registration of a product under this Act is not to be construed as giving the registrant a right to exclusive use of the name or to its use if it has been covered by others.

Based on your response to the sulfur registration Standard Guidance Packet, EPA has reregistered the product listed above.

The labeling referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide, and Rodenticide Act, is acceptable subject to the comments listed below. Five copies of the finished label(s) must be submitted.

1. Change the spelling to "sulfur" wherever it appears on the label.
2. Add the following statements to the Human hazards Section:

Wash thoroughly with soap and water after handling.
Remove contaminated clothing and wash before reuse.

This registration is granted under the authority of section 3(q) of the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) as amended.

Serry M. Jacoby
Product Manager (21)
Fungicide-Insecticide Branch
Registration Division (75-76)

ATTACHMENT IS APPLICABLE

SIGNATURE OF APPROVING OFFICIAL

Serry M. Jacoby

DATE

5/14/85

Front Panel



LILLY/MILLER

S U L P H U R

Illustration

ACTIVE INGREDIENT

Sulphur.....	90%
INERT INGREDIENTS.....	10%
Total	100%

Fungicide-Insecticide
Dust or Spray

FOR CONTROL OF CERTAIN FRUIT DISEASES SUCH AS BROWN ROT OF PEACHES, CHERRIES, PLUMS AND PRUNES; POWDERY MILDEW OF STRAWBERRIES, ROSES, CHRYSANTHEMUMS; RUST OF BEANS, HOLLY-HOCKS AND SNAPDRAGONS; AND FOR CONTROL OF CERTAIN MITES OF CITRUS FRUITS.

Keep out of reach of children

CAUTION

See right side panel for additional cautions

NET WT. 1 LB. 4 OZ.

ACCEPTED
with COMMENTS
in EPA Letter Dated
REREGISTERED

MAY 14 1985

Under the Federal Insecticide,
Fungicide, and Rodenticide Act
as amended, for the pesticide
registered under EPA Reg. No.

802-16

Left Side Panel

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION

Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists.

Hand thoroughly with soap & water after handling. Remove contaminated clothing & wash before reuse.

ENVIRONMENTAL HAZARDS


Do not apply directly to water. Do not contaminate water by cleaning of equipment or disposal of wastes.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

STORAGE AND DISPOSAL

Do not reuse empty container. Wrap and put in trash collection.



Please read the entire label.
Use only as directed,
noting all use precautions.

GENERAL DIRECTIONS

1. Before dusting, shake duster vigorously to mix air with the sulphur. Dust lightly. Frequent light dustings are more effective than single heavy dustings.
2. As a spray, mix the required amount of SULFUR with sufficient water to make a thin, creamy paste before diluting with the balance of the required amount of water.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangeloes, Tangerines) - for control of Six-Spotted, Citrus Rust, Citrus Flat and Yuma Spider Mites, dust or spray (2 level tablespoons per gallon of water) at the following times:

1. For Six-Spotted and Citrus Rust Mites, apply as mites appear.
2. For Citrus Flat and Yuma Spider Mites, make applications during October to March 15.
3. Repeat as necessary for control.
4. Avoid application during or preceding high temperatures.
5. Do not apply within two months of an oil application.

GRAPES - for control of Powdery Mildew, thoroughly dust or spray (1 level tablespoon per gallon of water) during the growing season at the following times:

1. When new shoot growth is 6 to 10 inches long.
2. Just before blossoming.
3. After the berries are set and not later than 1/4 grown.
4. Between the time the berries are 1/2 and fully grown.
5. Just as the grapes begin to ripen.

Right Side Panel

PEACHES, CHERRIES - for control of Brown Rot Blossom Blight, thoroughly dust or spray (3 level tablespoons per gallon of water) at the following times:

1. Just before petals begin to open.
2. When trees are in full blossom.
3. When most of the petals are falling.

PEACHES, CHERRIES, PLUMS, PRUNES - for control of Brown Rot of Fruit, thoroughly dust or spray (3 level tablespoons per gallon of water) at the following times:

1. First application 5 weeks before harvest, or earlier if rot appears on fruit.
2. Repeat at weekly intervals, making a last application just before harvest.

STRAWBERRIES - for control of Powdery Mildew dust or spray (3 level tablespoons per gallon of water) at the following times:

1. First application when disease is first noticed.
2. Repeat at 4 to 5 day intervals until mildew is under control.

VEGETABLES

BEANS - for the control of Rust, dust at the following times:

1. Start applications in early spring before any rust is noticed.
2. Repeat applications at 7 day intervals until blossom time.

FLOWERS

CHRYSANTHEMUMS, CLEMATIS, RHODODENDRONS -
for the control of Powdery Mildew, dust or
spray (3 tablespoons per gallon of water)
at the following times:

1. Start applications at first sign of mildew.
2. Repeat at 4 to 5 day intervals until mildew is under control. See Use Precautions.

HOLLYHOCKS, SNAPDRAGONS - for control of Rust,
dust or spray (1 level tablespoon per gallon of
water) at the following times:

1. Start control program in early spring.
2. Repeat as needed until disease is under control - applications twice a week may be necessary. See Use Precautions.

ROSES - for control of Powdery Mildew, dust or
spray (8 level tablespoons per gallon of water)
at the following times:

1. Start applications in spring when young growth first appears.
2. Repeat at 4 to 5 day intervals until mildew is under control. See Use Precautions.

USE PRECAUTIONS

Do not apply sulphur dusts or sprays when shade temperatures during the day are likely to exceed 85-90°F. as injury to foliage and fruit may result.

Do not apply to open Chrysanthemum flowers as bleaching may result.

Do not use sulphur within 30 days of an oil spray, nor combine with an oil spray.

Do not enter treated area for 24 hours until spray has dried or dust has settled.

Right Side Panel - Cont'd

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixtures with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

LC-125

THE CHAS. H. LILLY CO.
Portland, Oregon 97218

EPA Est. 802-OR-1

EPA Reg. No. 802-16