

following times:

1. Start applications at first sign of mildew.
2. Repeat at 4 to 5 day intervals until mildew is under control. See Use Precautions.

HOLLYHOCKS, SNAPDRAGONS—for control of Rust, dust or spray (1 level tablespoon per gallon of water) at the following times:

1. Start control program in early spring.
2. Repeat as needed until disease is under control—applications twice a week may be necessary. See Use Precautions.

ROSES—for control of Powdery Mildew, dust or spray (8 level tablespoons per gallon of water) at the following times:

1. Start applications in spring when young growth first appears.
2. Repeat at 4 to 5 day intervals until mildew is under control. See Use Precautions.

USE PRECAUTIONS: Do not apply SULFUR dusts or sprays when shade temperatures during the day are likely to exceed 85-90° F as injury to foliage and fruit may result. Do not apply to open Chrysanthemum flowers as bleaching may result. Do not use SULFUR within 30 days of an oil spray, nor combine with an oil spray.

STORAGE AND DISPOSAL

Store this product in its original container and keep in a locked storage area out of reach of children and domestic animals. Do not reuse container. Securely wrap container in several layers of newspaper and discard in trash.

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: Harmful if swallowed, inhaled or absorbed through the skin. Avoid breathing dust. Avoid contact with skin, eyes or clothing. In case of contact, immediately flush eyes or skin with plenty of water. Get medical attention if irritation persists. Wash thoroughly with soap and water after handling. Remove contaminated clothing and wash before reuse.

ENVIRONMENTAL HAZARDS

Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters.

NOTICE: Follow directions carefully. Timing and method of application, weather and crop conditions, mixes with other chemicals not specifically recommended, and other influencing factors in the use of this product are beyond the control of the seller. Buyer assumes all risks of use, storage or handling of this material not in strict accordance with directions given herewith.

Filled by weight not by volume. Product may settle after packaging.

THE CHAS. H. LILLY CO.
PORTLAND, OREGON 97218

EPA Est. 822-CP-1

81428-4

EPA Reg. No. 822-15

BEST AVAILABLE COPY




LILLY/MILLER.

Sulfur

ACCEPTED

JUN 7 1988


Under the Federal Insecticide,
Fungicide, and Rodenticide Act,
as amended, for the pesticide
registered under
EPA Reg. No. 802-16

ACTIVE INGREDIENT:

Sulfur 90%
INERT INGREDIENTS 10%

Fungicide-Insecticide Dust or Spray

FOR CONTROL OF CERTAIN
FRUIT DISEASES SUCH AS BROWN ROT
OF PEACHES, CHERRIES, PLUMS AND
PRUNES; POWDERY MILDEW OF
STRAWBERRIES, ROSES,
CHRYSANTHEMUMS, RUST OF BEANS,
HOLLYHOCKS AND SNAPDRAGONS;
FOR CONTROL OF CERTAIN MITES
OF CITRUS FRUITS.



Keep out of reach of children
CAUTION

See right side panel for additional cautions.

NET WT. 1 LB. 4 OZ.

with the sulfur. Dust lightly. Frequent light dustings are more effective than single heavy dustings.

2. As a spray, mix the required amount of SULFUR with sufficient water to make a thin, creamy paste before diluting with the balance of the required amount of water.

CITRUS (Grapefruit, Kumquats, Lemons, Limes, Oranges, Tangeloes, Tangerines)—for control of Six-Spotted, Citrus Rust, Citrus Flat and Yuma Spider Mites, dust or spray (2 level tablespoons per gallon of water) at the following times:

1. For Six-Spotted and Citrus Rust Mites, apply as mites appear.
2. For Citrus Flat and Yuma Spider Mites, make applications during October to March 15.
3. Repeat as necessary for control.
4. Avoid application during or preceding high temperatures.
5. Do not apply within two months of an oil application.

GRAPES—for control of Powdery Mildew, thoroughly dust or spray (1 level tablespoon per gallon of water) during the growing season at the following times:

1. When new shoot growth is 6 to 10 inches long.
2. Just before blossoming.
3. After the berries are set and not later than 1/4 grown.
4. Between the time the berries are 1/2 and fully grown.
5. Just as the grapes begin to ripen.

PEACHES, CHERRIES—for control of Brown Rot Blossom Blight, thoroughly dust or spray (3 level tablespoons per gallon of water) at the following times:

1. Just before petals begin to open.
2. When trees are in full blossom.
3. When most of the petals are falling.

PEACHES, CHERRIES, PLUMS, PRUNES—for control of Brown Rot of Fruit, thoroughly dust or spray (3 level tablespoons per gallon of water) at the following times:

1. First application 5 weeks before harvest, or earlier if rot appears on fruit.
2. Repeat at weekly intervals, making a last application just before harvest.

STRAWBERRIES—for control of Powdery Mildew dust or spray (3 level tablespoons per gallon of water) at the following times:

1. First application when disease is first noticed.
2. Repeat at 4 to 5 day intervals until mildew is under control.

VEGETABLES

BEANS—for the control of Rust, dust at the following times:

1. Start applications in early spring before any rust is noticed.
2. Repeat applications at 7 day intervals until blossom time.

FLOWERS

CHRYSANTHEMUMS, CLEMATIS, RHODODENDRONS—for the control of Powdery Mildew, dust or

BEST AVAILABLE COPY