



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, DC 20460

OFFICE OF CHEMICAL SAFETY
AND POLLUTION PREVENTION

December 23, 2020

Rebecca M. Horton
Agent
Lilly Miller Brands
Suite 700, 1000 Parkwood Circle
Atlanta GA 30339

Subject: Label Amendment – Minor Label Revisions
Product Name: Miller’s Garden Sulphur
EPA Registration Number: 802-16
Application Date: 12/3/2020
Decision Number: 568682

Dear Ms. Horton:

The amended label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act, as amended, is acceptable. This approval does not affect any conditions that were previously imposed on this registration. You continue to be subject to existing conditions on your registration and any deadlines connected with them.

A stamped copy of your labeling is enclosed for your records. This labeling supersedes all previously accepted labeling. You must submit one copy of the final printed labeling before you release the product for shipment with the new labeling. In accordance with 40 CFR 152.130(c), you may distribute or sell this product under the previously approved labeling for 18 months from the date of this letter. After 18 months, you may only distribute or sell this product if it bears this new revised labeling or subsequently approved labeling. “To distribute or sell” is defined under FIFRA section 2(gg) and its implementing regulation at 40 CFR 152.3.

Should you wish to add/retain a reference to the company’s website on your label, then please be aware that the website becomes labeling under the Federal Insecticide Fungicide and Rodenticide Act and is subject to review by the Agency. If the website is false or misleading, the product would be misbranded and unlawful to sell or distribute under FIFRA section 12(a)(1)(E). 40 CFR 156.10(a)(5) list examples of statements EPA may consider false or misleading. In addition, regardless of whether a website is referenced on your product’s label, claims made on the website may not substantially differ from those claims approved through the registration process. Therefore, should the Agency find or if it is brought to our attention that a website contains false or misleading statements or claims substantially differing from the EPA approved registration, the website will be referred to the EPA’s Office of Enforcement and Compliance.

Your release for shipment of the product constitutes acceptance of these conditions. If these conditions are not complied with, the registration will be subject to cancellation in accordance

Page 2 of 2
EPA Reg. No. 802-16
Decision No. 568682

with FIFRA section 6. If you have any questions, please contact Manjula Unnikrishnan by phone at 703-347-8520, or via email at unnikrishnan.manjula@epa.gov.

Sincerely,

A handwritten signature in black ink, appearing to read "Jordan Page", is written over a light gray rectangular background.

Jordan Page, (Acting) Product Manager 24
Fungicide and Herbicide Branch
Registration Division (7505P)
Office of Pesticide Programs

Enclosure

{NOTE TO REVIEWER [] Indicates alternate or optional wording. {} Indicates notes to Reviewer that will not appear on the final label}

MILLER'S GARDEN SULPHUR

[Additional Brand Name: Lilly Miller Sulfur Dust]
Intended For [Outdoor] Residential Use Only


Active Ingredient:

Sulfur* 90%

Other Ingredients: 10%

Total:.....100%

*Contains 0.54 lb active ingredient per gallon.



KEEP OUT OF REACH OF CHILDREN

CAUTION

See [side][back] panel for additional Precautionary Statements and First Aid.

NET WT ____

EPA Reg. No. 802-16

EPA Est. No 67572-GA-1A, 67572-GA-2B, 69703-GA-1C, 69703-GA-2D

Subscript used is first letter of run code on container.

FIRST AID	
Have product container or label with you when calling a poison control center or doctor, or going for treatment.	
If in eyes	<ul style="list-style-type: none"> • Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. • Call a poison control center or doctor for treatment advice.
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • Have person sip a glass of water if able to swallow. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give anything by mouth to an unconscious person
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth if possible. • Call a poison control center or doctor for further treatment advice
If on skin or on clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15-20 minutes. • Call a poison control center or doctor for treatment advice.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals

CAUTION: Causes moderate eye irritation. Harmful if inhaled, absorbed through skin, or swallowed. Avoid breathing dust. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet. Remove contaminated clothing and wash clothing before reuse.

ENVIRONMENTAL HAZARDS

DO NOT apply directly to water. **DO NOT** contaminate water when disposing of rinsate or equipment washwater.

[If you have a medical emergency, need consultation on diagnosis, therapy and analysis please call (800) 420-9347]

[If you have questions relating to product use or application, please call (866) 945-5033]

[If you have a product spill or transportation emergency call (800) 424-9300]

[For Consumer Refund: TechPac, LLC PO Box 95437 Palatine, IL 60095-0437]

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Intended for residential use only.

This product is a combination fungicide and insecticide that can be used as a dust or spray. It can be used on listed flowers, ornamentals, fruits, and vegetables. It kills listed pests and controls diseases including powdery mildew, leaf spot, black spot, scab, rust, and more listed below.

HOW TO USE

Please read the entire label. Use only as directed, noting all use restrictions. Consult State Agricultural Experiment Station or the State Agricultural Extension Service for sensitive plants and additional information as the timing, number and rate of application needed may vary with local conditions.

AS A DUST: Before dusting, shake duster vigorously to mix air with the sulfur. Dust lightly. Frequent light dustings are more effective than single heavy dustings.

AS A SPRAY: Remove cap with dip tube to measure dust. Mix the required amount of MILLER'S GARDEN SULPHUR with sufficient water to make a thin, creamy paste before diluting with the balance of the required amount of water. Use about 1 gallon of diluted spray per 1000 square feet.

tsp = teaspoon tbs = tablespoon

Fruits & Nuts

Make full coverage applications on both sides of leaves. Apply to plant foliage, stems, and flowers to leave a “barely visible” dust. As a preventive measure or for light infestations, use the lower dosage rates.

USE RESTRICTIONS: DO NOT apply MILLER’S GARDEN SULPHUR dusts or sprays when shade temperatures during the day are likely to exceed 85° - 90° F. Injury to foliage and fruit may result. **DO NOT** use within 30 days of an oil spray (60 days on citrus). **DO NOT** combine with an oil spray.

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Almonds	Brown Rot, Scab, Powdery Mildew, Leaf Spot, Mites (Blister, European red, Flat, Silver, Twospotted), Rust, Twig Blight	Dust or spray (2½-7¾ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods.	DO NOT exceed 0.6 lb. [0.54 lbs. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.2 lbs. [6.5 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 12 applications per year. Minimum retreatment interval: 30 days.
Apples, Pears	Scab, Powdery Mildew, Mites (Blister, European red, Silver, Twospotted), Apple Cedar Rust, Black Rot, Frog Eye, Quince Rust, Sooty Blotch	Spray (2½-7¾ tbsp per gallon of water) when buds begin to swell and thereafter until 2 weeks after the petals fall. Treat Delicious Apples and Anjou Pears only before bloom or injury may occur	DO NOT exceed 0.6 lb. [0.54 lbs. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.5 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days.
Avocados	Mites, Avocado Thrips*	Dust or spray (2½-7¾ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.6 lb. [0.54 lbs. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.5 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per

			year. Minimum retreatment interval: 7 days
Bushberries (Blueberries, Currants, Huckleberries)	Powdery Mildew	Dust or spray (1½-3¾ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.3 lb. [0.27 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.8 lbs. [7.02 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Canefruits (Blackberries, Boysenberries, Dewberries, Gooseberries, Loganberries, Raspberries)	Powdery Mildew	Dust or spray (1½-3¾ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.3 lb. [0.27 lb. a.i.] per product per 1000 sq. ft. per application. DO NOT exceed a total of 7.8 lbs. [7.02 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Strawberries	Powdery Mildew	Dust or spray (1⅓-2½ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Cherries	Powdery Mildew, Silver Mite, Brown Rot, Cherry Leaf Spot*, Rust, Brown rot	Dust or spray (2½-7¾ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods. The first three applications will	DO NOT exceed 0.6 lb. [0.54 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.5 lbs. a.i.] of product

	Blossom Blight, Leaf spot, Mites, Red Spider, Scab, and Stem Blight.	also provide control of Brown Rot Blossom Blight. Weekly applications from 5 weeks before and continuing up to the day of harvest will provide control of Brown Rot of Fruit.	per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 10 days
Citrus (Oranges, Tangelos, Tangerines, Grapefruit, Lemons, Limes, Kumquats, Citron-Citrus)	Citrus Thrips, Citrus Flat Mite, Yuma Spider Mite, Six-spotted Mite, Citrus Rust Mite, Mites, Scales (Soft, Red, Yellow, Purple, Black, and Citricola), Brown Rot, Rust Mite, Two-Spotted Mite, Citrus Red Spider, Thrips	Dust or spray (2¼ - 5¼ tbsp per gallon of water) when pests first appear. Avoid application during or preceding high temperatures. DO NOT apply within two months of an oil application.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 6.8 lbs. [6.12 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 17 applications per year. Minimum retreatment interval: 21 days
Dates	Banks Grass Mites	Dust or spray (5¼ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Figs	Mites	Dust or spray (1⅓ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 1.7 lbs. [1.53 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 17 applications per year. Minimum

			retreatment interval: 21 days
Olives	Mites	Dust or spray (2½ - 7¾ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Macadamias, Walnuts	Mites	Dust or spray (2½ - 5¼ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Pecans	Mites	Dust or spray (1⅓ - 2⅔ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 4.2 lbs. [3.78 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 21 applications per year. Minimum retreatment interval: 10 days
Pomegranates	Mites	Dust or spray (¾ - 2⅔ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 3.4 lbs. [3.06 lbs. a.i.] of product per 1000 sq. ft. per year.

			DO NOT exceed a total of 17 applications per year. Minimum retreatment interval: 21 days
Grapes	Powdery Mildew, Mites, Rust, Phomopsis	Dust or spray ($\frac{1}{2}$ - $2\frac{2}{3}$ tbsp per gallon of water) during the growing season when new shoot growth is 6 to 10 inches long. Concord and other labrusca type grapes may be injured by MILLER'S GARDEN SULPHUR	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.8 lbs. [2.52 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 14 applications per year. Minimum retreatment interval: 26 days
Mangoes	Powdery Mildew	Dust or spray ($2\frac{2}{3}$ - $7\frac{3}{4}$ tbsp per gallon of water) when pests first appear. Repeat at bloom, at fruit set, and 3 weeks later.	DO NOT exceed 0.6 lb. [0.54 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.4 lbs. [2.16 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 4 applications per year. Make four applications, as directed in Application Directions.
Nectarines, Peaches	Brown rot blossom blight, Coryneum blight, Mites, Powdery mildew, Silver Mite, Brown Rot, Cherry Leaf Spot, Rust, Scab, Leaf Spot, Peach Canker, Shothole, Red Spider	Dust or spray ($2\frac{2}{3}$ - $7\frac{3}{4}$ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods. The first three applications will also provide control of Brown Rot Blossom Blight. Weekly applications from 5 weeks before and continuing up to the day of harvest will provide control of Brown Rot of Fruit.	DO NOT exceed 0.6 lb. [0.54 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.5 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 10 days

Papaya	Mites (except Carmine mites)	Dust or spray ($2\frac{2}{3}$ - $7\frac{3}{4}$ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.6 lb. [0.54 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.5 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days
Peanuts	Leaf spot, Powdery mildew, Rust	Dust or spray ($1\frac{1}{3}$ - 2 tbsp per gallon of water) when disease first appears. Apply when disease threatens during rainy periods.	DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Pistachios	Citrus Flat Mite, Mites	Dust or spray ($2\frac{2}{3}$ - $5\frac{1}{4}$ tbsp per gallon of water) when mites first appear.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Plums, Prunes	Brown rot blossom blight, Leaf spot, Mites, Powdery mildew, Silver Mite, Brown Rot, Cherry Leaf Spot,	Dust or spray ($2\frac{2}{3}$ - $7\frac{3}{4}$ tbsp per gallon of water) before blossoms open, during bloom, and after bloom. Apply when disease threatens during rainy periods. Weekly applications from 5 weeks before and continuing up to the day of harvest will	DO NOT exceed 0.6 lb. [0.54 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.5 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per

	Rust, Scab, Shothole, Red Spider	provide control of Brown Rot of Fruit.	year. See Application Directions for retreatment schedule to control Brown Rot of Fruit. For other uses, minimum retreatment interval: 10 days.
Quince	Brown rot blossom blight, Powdery mildew	Dust or spray (2 $\frac{2}{3}$ - 7 $\frac{3}{4}$ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Sugar beets	Powdery mildew, Red spider mites*	Dust or spray (2 $\frac{2}{3}$ - 5 $\frac{1}{4}$ tbsp per gallon of water) when disease first appears. For spider mites, dust or spray (2-11 tbsp/gallon) when pests first appear.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

Vegetables

Make full coverage applications on both sides of leaves. Apply to plant foliage, stems, and flowers to leave a “barely visible” dust. As a preventive measure or for light infestations, use the lower dosage rates.

USE RESTRICTIONS: DO NOT apply MILLER’S GARDEN SULPHUR dusts or sprays when shade temperatures during the day are likely to exceed 85° - 90° F. Injury to foliage and fruit may result. **DO NOT** use within 30 days of an oil spray. **DO NOT** combine with an oil spray.

Garden Crop	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Artichoke	Leaf Spot	Dust or spray (2 $\frac{2}{3}$ - 5 $\frac{1}{4}$ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Asparagus	Rust, Mites	Dust or spray (2 $\frac{2}{3}$ - 7 $\frac{3}{4}$ tbsp per gallon of water) when pests first appear and repeat through fern growth season.	DO NOT exceed 0.6 lb. [0.56 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 15 lbs. [13.8 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 25 applications per year. Minimum retreatment interval: 7 days
Beans including Adzuki, Broad, Dry, Jackbean, Kidney, Lablab, Lima, Moth, Mung, Navy, Pink, Pinto, Rice, Runner, Snap, Tepary, Urd, Wax, Yardlong Peas including Blackeyed, Chickpea, Cowpea, Crowder, Dry, Dwarf,	Powdery Mildew, Rust, Thrips, Spider Mites, Leaf Spot, Mites	Dust or spray ($\frac{3}{4}$ - 2 $\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed. For Rust control, start applications in early spring before any rust is noticed. Repeat applications at 10 day intervals until bloom.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days

<p>Edible Podded, English, Field, Garden, Green, Pigeon, Shelled, Snow, Southern, Succulent, Sugar Snap, Unshelled</p>			
<p>Beets, Table</p>	<p>Mites, Powdery Mildew</p>	<p>Dust or spray ($\frac{3}{4}$ - $1\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.</p>	<p>DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days</p>
<p>Carrots, Garlic, Onions, Shallots</p>	<p>Petrobia Mite, Powdery Mildew, Downy Mildew*</p>	<p>Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.</p>	<p>DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days</p>
<p>Celery, Endive, Escarole, Swiss Chard, Salsify</p>	<p>Mites, Powdery mildew, Rust</p>	<p>Dust or spray (1 – $1\frac{1}{2}$ tbsp per gallon of water) when pests first appear.</p>	<p>DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year.</p>

			Minimum retreatment interval: 10 days
Cucumbers	Powdery Mildew	Dust or spray ($\frac{1}{2}$ - 1 tbsp per gallon of water) when pests first appear. May cause injury at when used at higher rates during excessive temperatures. Use lower rate ($\frac{1}{2}$ tbsp) when temperatures exceed 85°F.	DO NOT exceed 0.08 lb. [0.07 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.1 lbs. [1.9 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Cole crops including Broccoli, Brussels sprouts, Cabbage, Cauliflower	Mites, Powdery Mildew, Septoria Leaf Spot	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Eggplant	Powdery Mildew	Dust or spray (1 – $1\frac{1}{2}$ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Greens* including Collards, Kale, Mustard Greens	Mites, Powdery Mildew	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year.

			Minimum retreatment interval: 14 days
Horseradish*, Parsnips*, Radishes*, Rutabagas*, Turnips*	Powdery Mildew, Mites	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Lettuce* (Head & Leaf)	Powdery Mildew, Red Spider Mite, Rust	Dust or spray ($1\frac{1}{3}$ - $2\frac{2}{3}$ tbsp per gallon of water) when at early leaf stage.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Melons including Watermelon, Cantaloupe, Muskmelon	Powdery Mildew	Dust or spray ($5\frac{1}{4}$ - $6\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed $\frac{1}{2}$ lb. [0.45 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 13 lbs. [11.7 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Okra	Leaf Spot, Mites, Powdery Mildew	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year.

			Minimum retreatment interval: 14 days
Peppers (Anaheim, Banana, Bell, Cayenne, Chili, Ghost, Habanero, Jalapeno, Pepperoncini, Poblano, Scotch Bonnet, Serrano, Sweet)	Leaf Spot, Powdery Mildew	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Peppermint Tops, Spearmint Tops	Powdery Mildew	Dust or spray (1 – $1\frac{1}{2}$ tbsp per gallon of water) when pests first appear.	DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 15 days
Potatoes	Leaf Spot, Powdery Mildew, Two-spotted mites	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 10 days
Pumpkins*, Squash* (Summer & Winter)	Powdery Mildew	Dust or spray ($1\frac{1}{3}$ - $2\frac{2}{3}$ tbsp per gallon of water) when disease first appears. Avoid application during or preceding high temperatures.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of

			26 applications per year. Minimum retreatment interval: 7 days
Soybeans	Leaf Spot, Powdery Mildew, Rust	Dust or spray (2½-3¼ tbsp per gallon of water) when disease first appears.	DO NOT exceed 0.3 lb. [0.27 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 7.8 lbs. [7.02 lb. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Spinach	Powdery Mildew, Rust	Dust or spray (1 – 1½ tbsp per gallon of water) when pests are first noticed. May cause injury to certain varieties of spinach. Apply to a small area first to test for phytotoxicity before making a full application.	DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Sweet Potatoes	Leaf Spot, Mites, Powdery Mildew, Red Spider Mites	Dust or spray (1½ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.1 lb. [0.09 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 2.6 lbs. [2.34 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 14 days
Tomatoes	Tomato Russet Mite, Mites, Powdery Mildew	Dust or spray (2½ - 5¼ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.4 lb. [0.36 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 10.4 lbs. [9.36 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year.

			Minimum retreatment interval: 7 days
--	--	--	--------------------------------------

*Not for use in California.

Ornamental Flowers, Shrubs, & Trees

To control black spot and red spiders on roses, ornamental shrubs, and perennials (except in California).

To control rust and powdery mildew on ornamental flowers, shrubs and trees, dust or spray as directed to completely cover foliage. As a preventive measure or for light infestations, use the lower dosage rates.

USE RESTRICTIONS: DO NOT apply MILLER'S GARDEN SULPHUR dusts or sprays when shade temperatures during the day are likely to exceed 85° - 90° F. Injury to foliage and fruit may result. **DO NOT** use within 30 days of an oil spray. **DO NOT** combine with an oil spray.

Follow directions given for specific plants.

Ornamental Flowers, Shrubs & Trees	Target Pests/Diseases	Application Directions	Application Restrictions Per 1000 sq. ft.
Alder, Azalea, Buttonbush, Catalpa, Cosmos, Dahlia, Daisy, Delphinium, Dogwood, Euonymus, Golden Fleece, Hibiscus, Honeysuckle, Horse chestnut, Lady's mantle, Ligustrum, Lilac, Linden, Matrimony vine, Crepe myrtle, Poplar, Rhododendron, Staghorn sumac, Silver vine, Spiraea, Sweetpea,	Powdery Mildew	Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

Sunflower, Sycamore, Trumpetvine, Verbena, Willow, Witch hazel, Zinnia			
Arborvitae, Cedar, Sweetpea	Red Spider Mites	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Aster	Powdery Mildew, Rust*	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Calendula, Cherry laurel, Columbine, English ivy, Foxglove, Petunia, Sage, Smilax	Leaf Spot	Dust or spray ($\frac{3}{4}$ - $2\frac{2}{3}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

Carnation	Red Spider Mites, Rust*	Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Chrysanthemum	Powdery Mildew, Red Spider Mites, Rust*	Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Clematis	Leaf Spot, Red Spider Mites	Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Hydrangea	Botrytis Blight, Leaf Spot, Powdery Mildew, Red Spider Mites	Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per

			year. Minimum retreatment interval: 7 days
Phlox	Leaf Spot, Powdery Mildew	Dust or spray ($\frac{3}{4}$ - $2\frac{1}{2}$ tbsp per gallon of water) when pests are first noticed.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Hollyhocks, Snapdragons	Rust	Dust or spray (1 tablespoon per gallon of water) in early spring.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days
Roses	Powdery Mildew, Black Spot, Red Spider Mites, Rust*, Brown Canker*	Dust or spray (4 tbsp per gallon of water) in spring when young growth first appears. See USE RESTRICTIONS.	DO NOT exceed 0.2 lb. [0.18 lb. a.i.] of product per 1000 sq. ft. per application. DO NOT exceed a total of 5.2 lbs. [4.68 lbs. a.i.] of product per 1000 sq. ft. per year. DO NOT exceed a total of 26 applications per year. Minimum retreatment interval: 7 days

*Not for use in California.

STORAGE AND DISPOSAL

Storage: Store product in original container in a cool, dry, locked place out of the reach of children and pets.

Pesticide Disposal & Container Handling: Nonrefillable container. **DO NOT** reuse or refill this container. *If empty* – Place in trash or offer for recycling if available. *If partly filled* – Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

Filled by weight not by volume. Product may settle after packaging.

NOTICE: To the extent consistent with applicable law, seller makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. To the extent consistent with applicable law, buyer assumes all risk of use and/or handling of this material when such use and/or handling is contrary to label instructions.

OR:

NOTICE: To the extent consistent with applicable law, buyer assumes all responsibility for safety and use not in accordance with directions

[Lot no. to be added at production]

LILLY MILLER BRANDS
1000 Parkwood Circle, Suite 700
Atlanta, GA 30339

[company website]

[INSERT TRADEMARK STATEMENT(S) AS APPLICABLE]

{Optional Marketing Statements}

[Fungicide - Insecticide Dust or Spray

Controls fruit diseases on peaches, cherries, plums and prunes; powdery mildew on strawberries, roses, chrysanthemums; rust on beans, hollyhocks and snapdragons; and controls certain mites on citrus fruits.

For use on roses, grapes, citrus, fruits, berries, nuts, vegetables, flowers, shrubs, trees.

Controls powdery mildew, rust, scab, mites.]

Quick [Reference Table][Facts] {NOTE to Reviewer: Table below is optional}

{Icons may be used for any of the below; e.g.:}



[Product Facts]	[Insert listed claim(s)]
[Where to Use]	Fruit and Vegetable Gardens, Ornamental Plants and Shrubs, Flowers and Ground Covers [Insert listed claim(s)]
[Fruits & Vegetables]	[Insert listed fruits & vegetables] [And other listed fruits and vegetables]
[Kills] [&] [Controls]	[Insert listed pests] [And other listed [pests] [diseases] [and] [insects] [bugs]] [Insert listed clam(s)]
[People & Pets]	[Insert listed claim(s)]

[For use on {may list any one or combination of use sites from those featured in the label Directions for Use}]

For use on Flowers, Ornamentals, and Home Fruit & Vegetable Gardens

For use on roses, flowers, ground cover, bulbs, annual and perennial flowers

Controls [insert listed site] pests

Protect your [roses][and][flowers] from [damaging][harmful] pests.

Not harmful to [roses][flowers]

Won't harm [your][plants][shrubs] [flowers] [ornamentals] [blooms][roses][azaleas]

Tough on [insert listed pest], gentle on [your] [flowers] [ornamentals][insert listed site][roses]

Strong [insect][pest] protection for your [insert listed site]

Ideal for [outdoor] container gardening

Great for [azaleas] [camellias] [hollies] [roses] [bulbs] [annual] [and] [perennial] [flowers]

Intended for outdoor residential use only

Controls Rust, Scab and Powdery Mildew [on][insert listed site]

Controls [insert listed disease]

Hate the Bugs. Love the Garden. Get [insert brand name]!

Plant a love for gardening

Home & Garden insecticide

Broad spectrum insect control

Protects against listed damaging insects

Broad spectrum

Apply at first signs of insect damage or activity

Insect Killer

Bug Killer

Pest Killer

Kills listed [bugs] [pests] [insects] by contact

Starts killing immediately upon contact

Replace cap for safety

Remove liner before use

Cut here to open

[Fungicide][Insecticide][Dust][or][Spray]

Easy to Use

2-in-1 [action]: Disease and Insect [Protection][Control]

Dust lightly to control [insert listed pest] [listed insects][insert listed disease][listed diseases][listed insects and diseases]

Effective control [of] [rose] [pests] [listed insects][listed diseases]

Visible results

2-in-1 Formula: Disease & Insect Control

Dual action: Disease & Insect [control][protection]

Protects over XX plants from insects and diseases

Kills over XX listed pests

Protects plants from over XX [diseases][insects]

Kills by contact

Kills [insert listed pest]

Got [insert listed pest]? Get [insert brand]!

Kills chewing and sucking insects

Kills leaf eating insects

Controls [insert listed disease]

Trusted Brand since [insert year]
Established brand since [insert year]
Est. [insert year]
Trusted brand for over XX years
Formulated and Packaged in the U.S.A.
Guaranteed results[*] [*]or your money back
Guaranteed to work[*] [*]or your money back
100% Guarantee Will Work

People and pets may re-enter once the dust settles or product dries.
People and pets can return to the [lawn][treated area][garden] once the [product][dust] has settled or product has dried.
Once product has settled and dried, people and pets can return to the [lawn][treated area][garden].

Buy X Get X Free
Special Value! X% More (than X [lb]/[oz] [bag][can])

PEST LIST: {NOTE to Reviewer: This pest list is optional}

[Insects:

1. Silver Mite
2. Citrus Flat Mite
3. Citrus Rust Mite
4. Citrus Thrips
5. Petrobia Mite
6. Sixspotted Mite
7. Spider Mites
8. Thrips
9. Tomato Russet Mite
10. Yuma Spider Mite
11. Red Spider Mite
12. Mites
13. Twospotted mites
14. Blister mites
15. European red mites
16. Scale – Soft
17. Scale – Red
18. Scale – Yellow
19. Scale – Purple
20. Scale – Black
21. Scale – Citricola
22. Rust Mite
23. Citrus Red Spider

24. Red Spider

Diseases:

1. Brown Rot
2. Cherry Leaf Spot
3. Powdery Mildew
4. Rust
5. Scab
6. Leaf Spot
7. Botrytis Blight
8. Black Spot
9. Brown Rot Blossom Blight
10. Coryneum blight
11. Stem blight
12. Peach Canker
13. Shot Hole
14. Twig Blight
15. Apple Cedar Rust
16. Black Rot
17. Frog Eye
18. Quince Rust
19. Sooty Blotch
20. Phomopsis
21. Downy Mildew
22. Septoria Leaf Spot
23. Brown Canker]